

On Quantum Spaces of Lie Algebras

Lieven le Bruyn*
Michel Van den Bergh

*Dept. of Mathematics and Computer Science,
University of Antwerp, UIA
B-2610 Wilrijk (Belgium)*

September 1991 - 91-45

Abstract

The homogenization $H(\mathfrak{g})$ of the enveloping algebra of a finite dimensional Lie algebra \mathfrak{g} is an Artin-Schelter regular algebra. We characterize d -dimensional linear subspaces in the corresponding quantum space $\mathbb{P}_q(\mathfrak{g})$ as homogenizations of induced representations from codimension d Lie subalgebras. Furthermore we prove that the point variety has an embedded component iff there is a line, not contained in this point variety.

Key Words

Enveloping Algebras, Quantum Spaces

AMS Subject Classification

16S30

* Research Associate of the NFWO (Belgium)

On quantum spaces of Lie algebras

Lieven Le Bruyn* and Michel Van den Bergh
Departement Wiskunde en Informatica
Universitaire Instelling Antwerpen
B-2610 Wilrijk (Belgium)

September 10, 1991

Abstract

The homogenization $H(\mathfrak{g})$ of the enveloping algebra of a finite dimensional Lie algebra \mathfrak{g} is an Artin-Schelter regular algebra. We characterize d -dimensional linear subspaces in the corresponding quantum space $\mathbb{P}_q(\mathfrak{g})$ as homogenizations of induced representations from codimension d Lie subalgebras. Furthermore we prove that the point variety has an embedded component iff there is a line, not contained in this point variety.

1 Introduction and notations

To any finite dimensional Lie algebra

$$\mathfrak{g} = \mathbb{C}X_1 \oplus \cdots \oplus \mathbb{C}X_n \quad \text{with} \quad [X_i, X_j] = \sum_{k=1}^n \alpha_{ij,k} X_k$$

one classically associates an associative algebra $U(\mathfrak{g})$, the so-called “enveloping algebra” of \mathfrak{g} . The enveloping algebra carries a natural filtration

$$U_0(\mathfrak{g}) \subset U_1(\mathfrak{g}) \subset \cdots \subset U_n(\mathfrak{g}) \subset \cdots$$

*Research associate of the NFWO (Belgium)

such that $\text{gr } U(\mathfrak{g}) = S(\mathfrak{g})$, the symmetric algebra on \mathfrak{g} . This is the famous Poincaré-Birkhoff-Witt-theorem.

By homogenizing the defining relations of $U(\mathfrak{g})$ one can also associate a regular algebra in the sense of Artin and Schelter [1] to \mathfrak{g} . This algebra, which will be denoted by $H(\mathfrak{g})$ in the sequel, is the positively graded quadratic algebra generated by X_0, \dots, X_n where X_0 is taken to be central and the remaining defining relations are

$$X_i X_j - X_j X_i - \sum_{k=1}^n \alpha_{ij,k} X_k X_0$$

Observe that $H(\mathfrak{g})/(X_0 - 1)H(\mathfrak{g}) \cong U(\mathfrak{g})$, the enveloping algebra of \mathfrak{g} and that $H(\mathfrak{g})/X_0 H(\mathfrak{g}) \cong \text{gr}(U(\mathfrak{g})) \cong S(\mathfrak{g})$. From these facts one derives as usual (see e.g. [8] or [9]) that $H(\mathfrak{g})$ is a Noetherian maximal order domain, it is Auslander-regular of dimension $n + 1$ and satisfies the Cohen-Macaulay property.

Following M. Artin [3] we define the quantum space $\mathbb{P}_q(\mathfrak{g})$ of \mathfrak{g} to be $\text{Proj}(H(\mathfrak{g}))$ which is the quotient category of the category of all finitely generated graded left $H(\mathfrak{g})$ -modules modulo the subcategory of finite length modules. Via the identification $S(\mathfrak{g}) = H(\mathfrak{g})/X_0 H(\mathfrak{g})$, $\mathbb{P}(\mathfrak{g}^*) = \text{Proj } S(\mathfrak{g})$ will be thought of as the hyperplane at infinity of $\mathbb{P}_q(\mathfrak{g})$ with equation $X_0 = 0$.

As in [2] one can define points and lines in $\mathbb{P}_q(\mathfrak{g})$. This note originated from our attempts to understand the following example, taken from [7].

Example 1.1 *Let $\mathfrak{sl}(2, \mathbb{C}) = \mathbb{C}e \oplus \mathbb{C}f \oplus \mathbb{C}h$ with $[e, f] = h$, $[h, e] = 2e$ and $[h, f] = -2f$. Then the lines in $\mathbb{P}_q(\mathfrak{sl}(2, \mathbb{C}))$ are precisely the lines lying on the pencil of quadrics in $\mathbb{P}^3 = \text{Proj}(\mathbb{C}[e, f, h, t])$*

$$Q(\delta) = \mathcal{V}((h^2 + 4ef) + \delta^2 t^2)$$

In particular, all lines intersect the hyperplane $V(t)$ at infinity in a point of the conic $(t, h^2 + 4ef)$ which is precisely the embedded component of the point-variety of $\mathbb{P}_q(\mathfrak{sl}(2, \mathbb{C}))$.

Moreover all three dimensional Lie algebras behave in a similar way. That is, the point-variety has an embedded component lying in the plane at infinity and all lines not lying on the point-variety intersect with the hyperplane at infinity in a point of this component.

In this note we partially generalize Example 1.1 by showing that there always is a subvariety of the hyperplane at infinity in $\mathbb{P}_q(\mathfrak{g})$, closely related with a possible embedded component of the point variety, such that all lines, not lying in the hyperplane at infinity intersect this variety. In case $[\mathfrak{g}, \mathfrak{g}] = \mathfrak{g}$ this variety is the embedded component of the point variety.

We also show that the existence of such an embedded component is caused by the fact that there may be lines in $\mathbb{P}_q(\mathfrak{g})$, not lying in the point variety of $\mathbb{P}_q(\mathfrak{g})$.

The proofs of these results, given in Section 3, rely on the classification of linear subspaces in $\mathbb{P}_q(\mathfrak{g})$, given in the next section.

2 Linear subspaces, Lie subalgebras and polarizations

We first state a few generalities. To any finitely generated filtered $U(\mathfrak{g})$ -module

$$\cdots \subset M_{-1} \subset M_0 \subset M_1 \subset \cdots \subset \bigcup_{i=-\infty}^{\infty} M_i = M$$

we can associate a graded $H(\mathfrak{g})$ -module (and hence an object in $\mathbb{P}_q(\mathfrak{g})$) :

$$\tilde{M} = \bigoplus_{i=-\infty}^{\infty} M_i X_0^i \subset M[X_0, X_0^{-1}]$$

We observe that $\tilde{M}/(X_0 - 1)\tilde{M} \cong M$ and conversely, any X_0 -torsion free graded left $H(\mathfrak{g})$ -module arises in this way, see [4] or [6].

In the sequel there will be a few occasions where we use the notation \tilde{M} for a graded $H(\mathfrak{g})$ -module, even if it is not a priori obtained from some filtered $U(\mathfrak{g})$ -module M . I.e. \tilde{M} could have X_0 -torsion.

In this note we concentrate on a specific class of graded left $H(\mathfrak{g})$ -modules :

Definition 2.1 *A d -dimensional linear subspace of $\mathbb{P}_q(\mathfrak{g})$ is the object in $\mathbb{P}_q(\mathfrak{g})$ associated to a cyclic graded left $H(\mathfrak{g})$ -module $\tilde{M} = \bigoplus_{i=0}^{\infty} M_i$ with Hilbert series $\mathcal{H}(\tilde{M}, t) = \sum_{i=0}^{\infty} \dim_{\mathbb{C}}(M_i)t^i = 1/(1-t)^{d+1}$.*

Theorem 2.2 *The d -dimensional linear subspaces of $\mathbb{P}_q(\mathfrak{g})$ are :*

1. *The d -dimensional subspaces in $\mathbb{P}^{n-1} = \text{Proj}(S(\mathfrak{g}))$*

2. The homogenizations \tilde{M} of induced modules

$$M = U(\mathfrak{g}) \otimes_{U(\mathfrak{h})} \mathbb{C}_f$$

where \mathfrak{h} is a Lie subalgebra of \mathfrak{g} of codimension d and \mathbb{C}_f is the one-dimensional representation of \mathfrak{h} determined by $f \in \mathfrak{h}^*$ such that $f([\mathfrak{h}, \mathfrak{h}]) = 0$. Here the pair (\mathfrak{h}, f) is uniquely determined by \tilde{M} .

Proof Let \tilde{M} be a graded $H(\mathfrak{g})$ -module as in Definition 2.1. Then it is easy to see that \tilde{M} is either annihilated by X_0 or X_0 acts without torsion. In the first case \tilde{M} will be a quotient of $H(\mathfrak{g})/X_0H(\mathfrak{g}) = S(\mathfrak{g})$. I.e. \tilde{M} will represent a linear subspace in $\text{Proj } S(\mathfrak{g})$.

In the second case $M = \tilde{M}/(X_0 - 1)\tilde{M}$ is a filtered $H(\mathfrak{g})/(X_0 - 1)H(\mathfrak{g}) = U(\mathfrak{g})$ -module. I.e. there is map $\phi : U(\mathfrak{g}) \rightarrow M$ such that

$$\dim \phi(U_i(\mathfrak{g})) = \binom{d+i}{i}$$

The result now follows from the following proposition ■

Proposition 2.3 *Assume that M is a left $U(\mathfrak{g})$ -module and there is a surjective map $\phi : U(\mathfrak{g}) \rightarrow M$ such that*

$$\dim \phi(U_i(\mathfrak{g})) = \binom{d+i}{i} \quad \text{for } i \in \mathbb{N}. \quad (1)$$

Then there is a unique pair (\mathfrak{h}, f) such that

$$M \cong U(\mathfrak{g}) \otimes_{U(\mathfrak{h})} \mathbb{C}_f \quad (2)$$

with notation as in Theorem 2.2.

Furthermore the filtration induced on M via ϕ makes (2) into a filtered isomorphism.

Proof The uniqueness of the pair (\mathfrak{h}, f) is easy to see. Therefore we concentrate on its existence. First define

$$\mathfrak{h}' = U_1(\mathfrak{g}) \cap \ker \phi$$

Clearly $\dim \mathfrak{h}' = n - d$.

Using the fact that $U(\mathfrak{g})$ has a PBW-basis, it is easy to see that

$$U_1(\mathfrak{g})\mathfrak{h}' \cap U_1(\mathfrak{g}) = [\mathfrak{h}', \mathfrak{h}'] + \mathfrak{h}'$$

Furthermore if V is the image of $U_1(\mathfrak{g})\mathfrak{h}'$ in $U_2(\mathfrak{g})/U_1(\mathfrak{g}) = S_2(\mathfrak{g})$ then we compute that

$$\dim V = \binom{n+1}{2} - \binom{d+1}{2}$$

Hence we obtain

$$\begin{aligned} \dim U_1(\mathfrak{g})\mathfrak{h}' &= \dim V + \dim U_1(\mathfrak{g})\mathfrak{h}' \cap U_1(\mathfrak{g}) \\ &= \binom{n+1}{2} - \binom{d+1}{2} + \dim([\mathfrak{h}', \mathfrak{h}'] + \mathfrak{h}') \end{aligned} \quad (3)$$

Also, using (1) and the fact that $U_1(\mathfrak{g})\mathfrak{h}' \subset \ker \phi$ we find that

$$\dim U_1(\mathfrak{g})\mathfrak{h}' + \binom{d+2}{2} \leq \binom{n+2}{2} \quad (4)$$

Combining (3) and (4) we find

$$\dim([\mathfrak{h}', \mathfrak{h}'] + \mathfrak{h}') \leq n - d = \dim \mathfrak{h}'$$

Hence

$$[\mathfrak{h}', \mathfrak{h}'] \subset \mathfrak{h}'$$

and thus we find that \mathfrak{h}' is a Lie subalgebra of $U_1(\mathfrak{g})$.

Let \mathfrak{h} be the image of \mathfrak{h}' in \mathfrak{g} under the canonical decomposition $U_1(\mathfrak{g}) = \mathbb{C} \oplus \mathfrak{g}$. Since $1 \notin \mathfrak{h}'$, there will be a linear map $f \in \mathfrak{h}^*$ such that

$$\mathfrak{h}' = \{u - f(u) \mid u \in \mathfrak{h}\}$$

Let $u, v \in \mathfrak{h}$. Then

$$[u - f(u), v - f(v)] = [u, v] \in \mathfrak{h}'$$

and therefore $f([u, v]) = 0$.

We have now a surjective map

$$U(\mathfrak{g})/U(\mathfrak{g})\mathfrak{h}' \cong U(\mathfrak{g}) \otimes_{U(\mathfrak{h})} \mathbb{C}_f \rightarrow M$$

Applying (1) yields that this map must be a filtered isomorphism. ■

To any d -dimensional linear subspace in $\mathbb{P}_q(\mathfrak{g})$ we can associate a unique d -dimensional linear subspace in $\mathbb{P}^n = \mathbb{P}(\mathfrak{g}^* \oplus \mathbb{C}X_0^*)$ as follows. Let \tilde{M} be a graded quotient $H(\mathfrak{g}) \rightarrow \tilde{M}$. Then restriction to degree 1 defines a linear map $\mathfrak{g} \oplus \mathbb{C}X_0 \rightarrow \tilde{M}_1$, hence a linear subspace in \mathbb{P}^n . Conversely, a linear subspace in \mathbb{P}^n can at most correspond to one linear subspace in $\text{Proj } H(\mathfrak{g})$.

If we identify the open affine piece $X_0 = 1$ with \mathfrak{g}^* we would like to know the minimal dimension $d(f)$ of a linear subspace in $\mathbb{P}_q(\mathfrak{g})$ containing $f \in \mathfrak{g}^*$.

If $f \in \mathfrak{g}^*$ then we denote by B_f the bilinear form $(x, y) \rightarrow f([x, y])$ and with $\mathfrak{g}^f = \{x \in \mathfrak{g} \mid f([x, \mathfrak{g}]) = 0\}$ the radical which is a Lie subalgebra of \mathfrak{g} and as B_f induces a non-degenerate alternating bilinear form on $\mathfrak{g}/\mathfrak{g}^f$ we have that $n - \dim(\mathfrak{g}^f)$ is even.

In view of Theorem 2.2 we see that a d -dimensional linear subspace of $\mathbb{P}_q(\mathfrak{g})$ containing f is the homogenization of an induced one-dimensional representation of a codimension d Lie subalgebra \mathfrak{h} such that $f([\mathfrak{h}, \mathfrak{h}]) = 0$. In the terminology of [5, Ch I, §12] \mathfrak{h} is subordinate to f . One knows that the maximal dimension of such a Lie subalgebra is $\frac{1}{2}(n + \dim(\mathfrak{g}^f))$ meaning that

$$d(f) \geq \frac{1}{2}(n - \dim(\mathfrak{g}^f))$$

and if equality occurs, \mathfrak{h} is said to be a polarization of \mathfrak{g} at f . Note that polarizations exist for every f if \mathfrak{g} is solvable or $\mathfrak{g} = \mathfrak{sl}(n)$ and for arbitrary \mathfrak{g} a (solvable) polarization exists whenever f is a regular form, cfr. [5, Ch I, §11].

Corollary 2.4 *The minimal dimension d such that every point in $\mathbb{P}(\mathfrak{g}^* \oplus \mathbb{C}X_0^*)$ is contained in a d -dimensional linear subspace of $\mathbb{P}_q(\mathfrak{g})$ is equal to $\frac{1}{2}(n - r)$ where r is the index of \mathfrak{g} .*

If \mathfrak{g} is semi-simple, the index of \mathfrak{g} is equal to its rank. Moreover, in this case all linear subspaces in $\mathbb{P}_q(\mathfrak{g})$, not lying in the hyperplane at infinity and of minimal dimension, are obtained by parabolic induction.

3 Points, lines and embedded components

Recall from [2] that the bihomogeneous equations for the point-variety of $H(\mathfrak{g})$ are given by

$$x_i^{(1)}x_j^{(2)} - x_j^{(1)}x_i^{(2)} = \left(\sum \alpha_{ij,k}x_k^{(1)}\right)x_0^{(2)}$$

$$x_0^{(1)}x_i^{(2)} = x_i^{(1)}x_0^{(2)} \quad (5)$$

where $1 \leq i, j, k \leq n$.

To simplify some of the notations in the sequel, let us make the following conventions. If $U = \sum u_i X_i \in \mathfrak{g}$ then $u = \sum u_i x_i$, $u^{(1)} = \sum u_i x_i^{(1)}$, etc.... Similarly, if $U, V \in \mathfrak{g}$ and $[U, V] = \sum w_i X_i$ then $[u, v] = \sum w_i x_i$. As above, U and V may be equipped with the superscripts (1) and (2).

Using this notation we may rewrite (5) more succinctly as

$$x_i^{(1)}x_j^{(2)} - x_j^{(1)}x_i^{(2)} = [x_i^{(1)}, x_j^{(1)}]x_0^{(2)} \quad (6)$$

$$x_0^{(1)}x_i^{(2)} = x_i^{(1)}x_0^{(2)} \quad (7)$$

These equations determine a variety $X \subset \mathbb{P}(\mathfrak{g}^* \oplus \mathbb{C}x_0^*) \times \mathbb{P}(\mathfrak{g}^* \oplus \mathbb{C}x_0^*)$.

Let $\sigma : Y = pr_1(X) \rightarrow Y' = pr_2(Y)$ be the induced automorphism, which is given by shifting point modules one place to the right (we are working with left modules, instead of with right modules as in [2]). From the description (Theorem 2.2) of points as one-dimensional representations of $U(\mathfrak{g})$ or points at infinity it follows that σ is the identity on Y_{red} .

Theorem 3.1 *If we set $x_i = x_i^{(1)}$, then the defining relations for Y are*

$$\begin{aligned} (x_i[x_j, x_k] + x_j[x_k, x_i] + x_k[x_i, x_j])x_0 &= 0 \\ [x_i, x_j]x_0^2 &= 0 \end{aligned} \quad (8)$$

Proof We will cover Y by affine opens. First assume that $x_0 = x_0^{(1)} = 1$. Then from $\sigma = \text{id}$ on Y_{red} we may assume that $x_0^{(2)} = 1$. Then from (7) we obtain that $x_i^{(1)} = x_i^{(2)}$. Substituting this in (6) yields that $[x_i, x_j] = 0$ for $i, j \in \{1, \dots, n\}$. This set of equations clearly defines the same variety as (8) when $x_0 = 1$.

Assume now that $x_s = x_s^{(1)} = 1$ for some $s \in \{1, \dots, n\}$. Again we may assume that $x_s^{(2)} = 1$. Put $\epsilon_j = x_j^{(1)} - x_j^{(2)}$. Clearly $\epsilon_s = 0$, and from (7) it follows that $\epsilon_0 = 0$.

(6) and (7) may now be rewritten as

$$x_j\epsilon_k - x_k\epsilon_j = [x_j, x_k]x_0 \quad (9)$$

$$x_0\epsilon_j = 0 \quad (10)$$

$j, k \in \{1, \dots, n\}$.

Applying (9) with $j = s$ yields

$$\epsilon_k = [x_s, x_k]x_0$$

Resubstituting yields the equations

$$\begin{aligned} (x_s[x_j, x_k] + x_k[x_s, x_j] + x_j[x_k, x_s])x_0 &= 0 \\ [x_s, x_j]x_0^2 &= 0 \end{aligned} \quad (11)$$

where $j, k \in \{1, \dots, n\}$. Using the fact that $x_s = 1$, we find that the equations (11) are equivalent with the equations (8). ■

Now we give an interpretation for the two sets of equations occurring in (8).

Lemma 3.2 1. $A = V([x_i, x_j]; 1 \leq i, j \leq n) \subset \mathbb{P}(\mathfrak{g}^*)$ is the linear space of one-dimensional representations of \mathfrak{g}

2. $B = V(x_i[x_j, x_k] + x_j[x_k, x_i] + x_k[x_i, x_j]; 1 \leq i, j, k \leq n) \subset \mathbb{P}(\mathfrak{g}^*)$ classifies codimension one Lie subalgebras of \mathfrak{g}

Proof (1) is clear. A point of B is a surjective map $\phi : \mathfrak{g} \rightarrow \mathbb{C}$ satisfying $\phi(u)\phi([v, w]) + \phi(v)\phi([w, u]) + \phi(w)\phi([u, v]) = 0$. Let $\mathfrak{h} = \ker(\phi)$ and take $u, v \in \mathfrak{h}$ and $w \notin \mathfrak{h}$, it follows that $\phi([u, v]) = 0$ i.e. \mathfrak{h} is a codimension one Lie subalgebra. Conversely the quotient map $\mathfrak{g} \rightarrow \mathfrak{g}/\mathfrak{h} \cong \mathbb{C}$ for any codimension one Lie subalgebra \mathfrak{h} of \mathfrak{g} gives a point of B . ■

Corollary 3.3 Every line in $\mathbb{P}_q(\mathfrak{g})$, not lying in the hyperplane at infinity, intersects B .

Proof Such a line is given by a module of the form

$$(U(\mathfrak{g}) \otimes_{U(\mathfrak{h})} \mathbb{C}_f)^\sim$$

where \mathfrak{h} is a codimension one Lie subalgebra of \mathfrak{g} , and $f \in \mathfrak{h}^*$ is a linear form such that $f([\mathfrak{h}, \mathfrak{h}]) = 0$.

The intersection with the hyperplane at infinity is represented by

$$(U(\mathfrak{g}) \otimes_{U(\mathfrak{h})} \mathbb{C}_f)^\sim \otimes H(\mathfrak{g})/X_0H(\mathfrak{g}) \cong S(\mathfrak{g})/S(\mathfrak{g})\mathfrak{h}$$

The corresponding point in $\mathbb{P}(\mathfrak{g}^*)$ is given by the quotient map $\mathfrak{g} \rightarrow \mathfrak{g}/\mathfrak{h} \cong \mathbb{C}$. Hence it must be a point of B . ■

Corollary 3.4 *If $[\mathfrak{g}, \mathfrak{g}] = \mathfrak{g}$ then the point variety of $\mathbb{P}_q(\mathfrak{g})$ consists of*

1. *The origin.*
2. *The hyperplane at infinity.*
3. *An embedded component at infinity given by B .*

Proof Clear from the equations (8). ■

Theorem 3.5 *The following are equivalent*

1. *The point variety of $H(\mathfrak{g})$ is reduced, i.e. $Y = Y_{\text{red}} = V([x_i, x_j]x_0; 1 \leq i, j \leq n)$*
2. *All lines in $\mathbb{P}_q(\mathfrak{g})$ lie on the point variety*
3. *For every codimension one Lie subalgebra \mathfrak{h} of \mathfrak{g} holds $[\mathfrak{h}, \mathfrak{h}] = [\mathfrak{g}, \mathfrak{g}]$*

Proof From the results of the foregoing section it is clear that (2) and (3) are equivalent. Let us prove that (1) implies (3) :

First, there can be no codimension one Lie subalgebra \mathfrak{h} such that $[\mathfrak{g}, \mathfrak{g}]$ is not contained in \mathfrak{h} for otherwise there is a point in A not lying on B meaning that there would be an embedded point at infinity. Hence, assume there is a decomposition $\mathfrak{g} = \mathfrak{h} \oplus \mathbb{C}e$ where \mathfrak{h} is a codimension one Lie algebra and $[\mathfrak{h}, \mathfrak{h}]$ is properly contained in $[\mathfrak{g}, \mathfrak{g}] \subset \mathfrak{h}$. Let $\chi_{\mathfrak{h}} : \mathfrak{g} \rightarrow \mathbb{C}$ be the projection on the second factor. Then, there exists a $\phi \in \mathfrak{g}^*$ s.t. $\phi([\mathfrak{h}, \mathfrak{h}]) = 0$ whereas $\phi([\mathfrak{g}, \mathfrak{g}]) \neq 0$. But then $\psi = \chi_{\mathfrak{h}} + \epsilon \cdot \phi$ is a point of B over the dual ring $\mathbb{C}[\epsilon]$ which is not a point of A over $\mathbb{C}[\epsilon]$ meaning again that the point variety has an embedded point.

Finally, let us prove that (3) implies (1) : we have to show that $A = B$ and as A is smooth it suffices to prove that any point of B over $\mathbb{C}[\epsilon]$ lies in A . So, take a surjection $\psi : \mathfrak{g} \rightarrow \mathbb{C}[\epsilon]$ with the property that

$$\psi(u)\psi([v, w]) + \psi(v)\psi([w, u]) + \psi(w)\psi([u, v]) = 0$$

Decompose $\psi = \psi_1 + \epsilon\psi_2$. Then the kernel of ψ_1 is a codimension one Lie subalgebra \mathfrak{h} . Put $\mathfrak{g} = \mathfrak{h} \oplus \mathbb{C}e$ where $\psi(e) = 1$. Substitute $u, v \in \mathfrak{h}$ and $w = e$ in the above relation and as $[\mathfrak{g}, \mathfrak{g}] = [\mathfrak{h}, \mathfrak{h}] \subset \mathfrak{h}$ we obtain $\psi([u, v]) = 0$, i.e. $\psi([\mathfrak{h}, \mathfrak{h}]) = \psi([\mathfrak{g}, \mathfrak{g}]) = 0$ so it belongs to B . ■

With this result in mind, we pose the question whether d -dimensional linear subspaces (for $d > 1$) of $\mathbb{P}_q(\mathfrak{g})$ arise in a similar way from embedded points in varieties describing lower-dimensional subspaces.

References

- [1] Artin M., Schelter W., Graded algebras of global dimension 3, *Adv.Math.* 66 (1987) 171-216
- [2] Artin M., Tate J., Van den Bergh M., Some algebras associated to automorphisms of elliptic curves, *Grothendieck Festschrift*, Vol. 1, 33-85, Birkhäuser (1990)
- [3] Artin M., *Geometry of quantum planes*, MIT preprint (1990)
- [4] Ascensio M., Van den Bergh M., Van Oystaeyen F., A new algebraic approach to microlocalization, *Trans. Amer. Math. Soc.*, 316, 2, 537-553, (1989).
- [5] Dixmier J., "Enveloping algebras", North-Holland (1977)
- [6] Ginsburg V., Characteristic varieties and vanishing cycles, *Inv.Math.* 84 (1986) 327-402
- [7] Le Bruyn L., Smith S.P., Homogenized $\mathfrak{sl}(2)$, *UIA-preprint* (1991)
- [8] Li Huishi, Van Oystaeyen F., *Zariskian filtrations*, monograph, to appear.
- [9] Levasseur Th., Some properties of non-commutative regular graded rings, *Proc. Edinb. Math. Soc.*, to appear.