

# noncommutative geometry @<sub>n</sub>

Lieven Le Bruyn

UNIVERSITEIT ANTWERPEN (UA), B-2020 ANTWERP (BELGIUM)

*E-mail address:* [lieven.lebruy@ua.ac.be](mailto:lieven.lebruy@ua.ac.be)

*URL:* <http://www.math.ua.ac.be/~lebruy>

2000 *Mathematics Subject Classification*. Primary 14A22,16A30 ; Secondary  
13A50, 14D20, 14L24, 16G20, 53D30

Research Director of the FWO (Belgium).

ABSTRACT. This text will never be published!

# Contents

Preface	11
<b>alg @<sub>n</sub></b>	1
Chapter 1. Generic Matrices.	3
1.1. Conjugacy classes of matrices	3
1.2. Simultaneous conjugacy classes.	13
1.3. Matrix invariants and necklaces	16
1.4. The trace algebra.	22
1.5. The symmetric group.	26
1.6. Necklace relations.	28
1.7. Trace relations.	35
1.8. Cayley-Hamilton algebras.	39
References.	44
Chapter 2. Reconstructing Algebras.	47
2.1. Representation spaces.	47
2.2. Some algebraic geometry.	49
2.3. The Hilbert criterium.	54
2.4. Semisimple modules	58
2.5. Some invariant theory.	62
2.6. Geometric reconstruction	68
2.7. The Gerstenhaber-Hesselink theorem.	73
2.8. The real moment map.	82
References.	88
<b>top @<sub>n</sub></b>	89
Chapter 3. Etale Cohomology.	91
3.1. Etale topology.	91
3.2. Central simple algebras	98
3.3. Quiver orders.	101
3.4. Simple roots.	114
3.5. Spectral sequences	119
3.6. Tsen and Tate fields	122
3.7. Coniveau spectral sequence	125
3.8. The Artin-Mumford exact sequence	129
References.	135
Chapter 4. Etale Slices.	137
4.1. $C^\infty$ slices.	137

4.2. Normal spaces.	139
4.3. Knop-Luna slices.	145
4.4. Smoothness.	149
4.5. Local structure.	155
4.6. $A_\infty$ -algebras.	161
4.7. Indecomposable roots.	165
4.8. Canonical decomposition.	173
References.	183
<b>loc @<sub>n</sub></b>	185
Chapter 5. Classification.	187
5.1. Type stratification.	187
5.2. Cayley-smooth locus.	193
5.3. Local classification.	200
5.4. Low dimensional orders.	202
5.5. Noncommutative smooth surfaces.	210
5.6. Complex moment map.	215
5.7. Preprojective algebras.	220
5.8. Quantum groups.	222
References.	229
Chapter 6. Nullcones.	231
6.1. Cornering matrices.	231
6.2. General subrepresentations.	236
6.3. Semistable representations.	240
6.4. Optimal corners.	246
6.5. Hesselink stratification.	250
6.6. Cornering quiver representations.	257
6.7. Simultaneous conjugacy classes.	262
6.8. Representation fibers.	269
References.	275
<b>geo @<sub>n</sub></b>	277
Chapter 7. Noncommutative Geometry	279
7.1. Formal structure.	279
7.2. Semi invariants.	285
7.3. Universal localization.	294
7.4. Compact manifolds.	301
7.5. Differential forms.	308
7.6. deRham cohomology.	321
7.7. Symplectic structure.	327
7.8. Necklace Lie algebras.	333
References.	336
Chapter 8. Moduli Spaces	337
8.1. Moment maps.	337
8.2. Dynamical systems.	341
8.3. Deformed preprojective algebras.	347

8.4. Hilbert schemes.	351
8.5. Hyper Kähler structure.	361
8.6. Calogero particles.	366
8.7. Coadjoint orbits.	370
8.8. Adelic Grassmannian.	374
References.	378
Bibliography	379
Index	383



## Preface

This is an example of an unnumbered chapter which can be used for a Preface or Foreword.

The purpose of this paper is to establish a relationship between an infinite-dimensional Grassmannian and arbitrary algebraic vector bundles of any rank defined over an arbitrary complete irreducible algebraic curve, which generalizes the known connection between the Grassmannian and line bundles on algebraic curves.

Author Name





## alg @<sub>n</sub>

noncommutative geometry @<sub>n</sub> is the study of *families* of (commutative) algebraic varieties (with specified connecting morphisms) which are locally controlled by noncommutative algebras. In the next two chapters we will clarify the latter part of this dogmatic statement.

Define the category  $\mathbf{alg}^{\text{tr}}$  of (not necessarily commutative)  $\mathbb{C}$ -algebras equipped with a  $\mathbb{C}$ -linear map  $tr : A \rightarrow \mathbb{C}$  which is a formal trace, that is, satisfies for all  $a, b \in A$

$$tr(a)b = btr(a) \quad tr(ab) = tr(ba) \quad \text{and} \quad tr(tr(a)b) = tr(a)tr(b).$$

Morphisms in  $\mathbf{alg}^{\text{tr}}$  are  $\mathbb{C}$ -algebra morphisms compatible with the trace structure. The forgetful functor to the category of *all*  $\mathbb{C}$ -algebras has a left adjoint

$$\mathbf{alg} \xrightarrow{tr} \mathbf{alg}^{\text{tr}} \quad A \mapsto A^{tr}$$

where  $A^{tr}$  is obtained by formally adjoining traces to cyclic words in a generating set of  $A$ . For  $a \in A$  and a fixed natural number  $n$ , we can define the formal Cayley-Hamilton polynomial  $\chi_a^{(n)}(t) \in A^{tr}[t]$  of degree  $n$  by expressing  $f(t) = \prod_{i=1}^n (t - \lambda_i)$  (in the commuting variables  $\lambda_i$ ) as a polynomial in  $t$  with all coefficients polynomials in the Newton functions  $\sum_{i=1}^n \lambda_i^k$  and replacing each occurrence of  $\sum_{i=1}^n \lambda_i^k$  by  $tr(a^k)$ . The *level  $n$  approximation* of the algebra  $A$  is defined to be the quotient algebra

$$A @_n = \frac{A^{tr}}{(tr(1) - n, \chi_a^{(n)}(a) \quad \forall a \in A^{tr})}$$

The main results, theorems 1.28 and 2.38, assert that we can recover the algebra  $A @_n$  and its central subalgebra  $tr A @_n$  from geometric data. If  $A$  is an affine  $\mathbb{C}$ -algebra with generating set  $\{a_1, \dots, a_m\}$ , then the representation space  $\underline{rep}_n A$  of all  $n$ -dimensional representations of  $A$  is a closed subscheme of  $m$ -tuples of  $n \times n$  matrices  $M_n^m = M_n \oplus \dots \oplus M_n$  and is stable under the  $GL_n$ -action on  $M_n^m$  by simultaneous conjugation. The  $GL_n$ -orbits in  $\underline{rep}_n A$  are the isomorphism classes of representations and an orbit is closed if and only if the corresponding representation is semisimple. Geometric invariant theory asserts the existence of an algebraic quotient  $\underline{iss}_n A = \underline{rep}_n A / GL_n$  classifying the isomorphism classes of  $n$ -dimensional semisimple representations of  $A$ . We recover  $A @_n$  as the algebra of  $GL_n$ -equivariant polynomial maps from  $\underline{rep}_n A$  to  $M_n$  and the central subalgebra  $tr A @_n$  as the coordinate ring of the quotient variety  $\underline{iss}_n A$ .

A family  $(X_i)_{i \in I}$  of (commutative) algebraic varieties is said to be locally controlled by a set of noncommutative algebras  $\mathcal{A}$  if every  $X_i$  is locally isomorphic to an irreducible component of  $\underline{iss}_n A$  for some  $n \in \mathbb{N}$  and  $A \in \mathcal{A}$ . When  $\mathcal{A}$  is

a collection of Quillen-smooth algebras (which implies that all  $\underline{rep}_n A$  are smooth varieties) we will determine the étale local structure of the  $X_i$  in the next part.

## Generic Matrices.

### 1.1. Conjugacy classes of matrices

We denote by  $M_n$  the space of all  $n \times n$  matrices  $M_n(\mathbb{C})$  and by  $GL_n$  the general linear group  $GL_n(\mathbb{C})$ . A matrix  $A \in M_n$  determines by left multiplication a linear operator on the  $n$ -dimensional vectorspace  $V_n = \mathbb{C}^n$  of *column vectors*. If  $g \in GL_n$  is the matrix describing the *base change* from the canonical basis of  $V_n$  to a new basis, then the linear operator expressed in this new basis is represented by the matrix  $gAg^{-1}$ . For a given matrix  $A$  we want to find an suitable basis such that the *conjugated matrix*  $gAg^{-1}$  has a simple form.

Consider the linear action of  $GL_n$  on the  $n^2$ -dimensional vectorspace  $M_n$


$$GL_n \times M_n \longrightarrow M_n \quad (g, A) \mapsto g.A = gAg^{-1}.$$

The *orbit*  $\mathcal{O}(A) = \{gAg^{-1} \mid g \in GL_n\}$  of  $A$  under this action is called the *conjugacy class* of  $A$ . We look for a particularly nice representative in a given conjugacy class. The answer to this problem is, of course, given by the *Jordan normal form* of the matrix.

With  $e_{ij}$  we denote the matrix whose unique non-zero entry is 1 at entry  $(i, j)$ . Recall that the group  $GL_n$  is generated by the following three classes of matrices :

- the *permutation* matrices  $p_{ij} = \mathbb{1}_n + e_{ij} + e_{ji} - e_{ii} - e_{jj}$  for all  $i \neq j$ ,
- the *addition* matrices  $a_{ij}(\lambda) = \mathbb{1}_n + \lambda e_{ij}$  for all  $i \neq j$  and  $0 \neq \lambda$ , and
- the *multiplication* matrices  $m_i(\lambda) = \mathbb{1}_n + (\lambda - 1)e_{ii}$  for all  $i$  and  $0 \neq \lambda$ .

Conjugation by these matrices determine the three types of *Jordan moves* on  $n \times n$  matrices, where the altered rows and columns are dashed.



Therefore, it suffices to consider sequences of these moves on a given  $n \times n$  matrix  $A \in M_n$ . The *characteristic polynomial* of  $A$  is defined to be the polynomial of

degree  $n$  in the variable  $t$

$$\chi_A(t) = \det(t\mathbb{1}_n - A) \in \mathbb{C}[t].$$

As  $\mathbb{C}$  is algebraically closed,  $\chi_A(t)$  decomposes as a product of linear terms

$$\prod_{i=1}^e (t - \lambda_i)^{d_i}$$

where the  $\{\lambda_1, \dots, \lambda_e\}$  are called the *eigenvalues* of the matrix  $A$ . Observe that  $\lambda_i$  is an eigenvalue of  $A$  if and only if there is a non-zero *eigenvector*  $v \in V_n = \mathbb{C}^n$  with eigenvalue  $\lambda_i$ , that is,  $A.v = \lambda_i v$ . In particular, the *rank*  $r_i$  of the matrix  $A_i = \lambda_i \mathbb{1}_n - A$  satisfies  $n - d_i \leq r_i < n$ . A nice inductive procedure using Jordan moves given in [23] gives a proof of the following *Jordan-Weierstrass theorem*.

**THEOREM 1.1 (Jordan-Weierstrass).** *Let  $A \in M_n$  with characteristic polynomial  $\chi_A(t) = \prod_{i=1}^e (t - \lambda_i)^{d_i}$ . Then,  $A$  determines unique partitions*

$$p_i = (a_{i1}, a_{i2}, \dots, a_{im_i}) \quad \text{of } d_i$$

*associated to the eigenvalues  $\lambda_i$  of  $A$  such that  $A$  is conjugated to a unique (up to permutation of the blocks) block-diagonal matrix*

$$J_{(p_1, \dots, p_e)} = \begin{bmatrix} \boxed{B_1} & & & & \\ & \boxed{B_2} & & & \\ & & \ddots & & \\ & & & \ddots & \\ & & & & \boxed{B_m} \end{bmatrix}$$

*with  $m = m_1 + \dots + m_e$  and exactly one block  $B_l$  of the form  $J_{a_{ij}}(\lambda_i)$  for all  $1 \leq i \leq e$  and  $1 \leq j \leq m_i$  where*

$$J_{a_{ij}}(\lambda_i) = \begin{bmatrix} \lambda_i & 1 & & & \\ & \lambda_i & \ddots & & \\ & & \ddots & \ddots & \\ & & & \ddots & 1 \\ & & & & \lambda_i \end{bmatrix} \in M_{a_{ij}}(\mathbb{C})$$

Let us prove uniqueness of the partitions  $p_i$  of  $d_i$  corresponding to the eigenvalue  $\lambda_i$  of  $A$ . Assume  $A$  is conjugated to another Jordan block matrix  $J_{(q_1, \dots, q_e)}$ , necessarily with partitions  $q_i = (b_{i1}, \dots, b_{im'_i})$  of  $d_i$ . To begin, observe that for a Jordan block of size  $k$  we have that

$$rk J_k(0)^l = k - l \quad \text{for all } l \leq k \text{ and if } \mu \neq 0 \text{ then } rk J_k(\mu)^l = k$$


for all  $l$ . As  $J_{(p_1, \dots, p_e)}$  is conjugated to  $J_{(q_1, \dots, q_e)}$  we have for all  $\lambda \in \mathbb{C}$  and all  $l$

$$rk (\lambda \mathbb{1}_n - J_{(p_1, \dots, p_e)})^l = rk (\lambda \mathbb{1}_n - J_{(q_1, \dots, q_e)})^l$$

Now, take  $\lambda = \lambda_i$  then only the Jordan blocks with eigenvalue  $\lambda_i$  are important in the calculation and one obtains for the ranks

$$n - \sum_{h=1}^l \#\{j \mid a_{ij} \geq h\} \quad \text{respectively} \quad n - \sum_{h=1}^l \#\{j \mid b_{ij} \geq h\}.$$

Now, for any partition  $p = (c_1, \dots, c_u)$  and any natural number  $h$  we see that the number  $z = \#\{j \mid c_j \geq h\}$



is the number of blocks in the  $h$ -th row of the dual partition  $p^*$  which is defined to be the partition obtained by interchanging rows and columns in the *Young diagram* of  $p$ . Therefore, the above rank equality implies that  $p_i^* = q_i^*$  and hence that  $p_i = q_i$ . As we can repeat this argument for the other eigenvalues we have the required uniqueness. Hence, the Jordan normal form shows that the classification of  $GL_n$ -orbits in  $M_n$  consists of two parts : a discrete part choosing

- a partition  $p = (d_1, d_2, \dots, d_e)$  of  $n$ , and for each  $d_i$ ,
- a partition  $p_i = (a_{i1}, a_{i2}, \dots, a_{im_i})$  of  $d_i$ ,

determining the sizes of the Jordan blocks and a continuous part choosing

- an  $e$ -tuple of distinct complex numbers  $(\lambda_1, \lambda_2, \dots, \lambda_e)$ .

fixing the eigenvalues. Moreover, this  $e$ -tuple  $(\lambda_1, \dots, \lambda_e)$  is determined only up to permutations of the subgroup of all permutations  $\pi$  in the symmetric group  $S_e$  such that  $p_i = p_{\pi(i)}$  for all  $1 \leq i \leq e$ . Whereas this gives a satisfactory set-theoretical description of the orbits we cannot put an Hausdorff topology on this set due to the existence of non-closed orbits in  $M_n$ . For example, if  $n = 2$ , consider the matrices

$$A = \begin{bmatrix} \lambda & 1 \\ 0 & \lambda \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} \lambda & 0 \\ 0 & \lambda \end{bmatrix}$$

which are in different normal form so correspond to distinct orbits. For any  $\epsilon \neq 0$  we have that


$$\begin{bmatrix} \epsilon & 0 \\ 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} \lambda & 1 \\ 0 & \lambda \end{bmatrix} \cdot \begin{bmatrix} \epsilon^{-1} & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} \lambda & \epsilon \\ 0 & \lambda \end{bmatrix}$$

belongs to the orbit of  $A$ . Hence if  $\epsilon \rightarrow 0$ , we see that  $B$  lies in the closure of  $\mathcal{O}(A)$ . As any matrix in  $\mathcal{O}(A)$  has trace  $2\lambda$ , the orbit is contained in the 3-dimensional subspace

$$\begin{bmatrix} \lambda + x & y \\ z & \lambda - x \end{bmatrix} \hookrightarrow M_2$$

In this space, the orbit-closure  $\overline{\mathcal{O}(A)}$  is the set of points satisfying  $x^2 + yz = 0$  (the determinant has to be  $\lambda^2$ ), which is a cone having the origin as its top : The orbit  $\mathcal{O}(B)$  is the top of the cone and the orbit  $\mathcal{O}(A)$  is the complement, see figure 1.

Still, for general  $n$  we can try to find the best separated topological quotient space for the action of  $GL_n$  on  $M_n$ . We will prove that this space coincide with the quotient variety determined by the invariant polynomial functions.

FIGURE 1. Orbit closure for  $2 \times 2$  matrices

If two matrices are conjugated  $A \sim B$ , then  $A$  and  $B$  have the same *unordered*  $n$ -tuple of eigenvalues  $\{\lambda_1, \dots, \lambda_n\}$  (occurring with multiplicities). Hence any symmetric function in the  $\lambda_i$  will have the same values in  $A$  as in  $B$ . In particular this is the case for the *elementary symmetric functions*  $\sigma_l$

$$\sigma_l(\lambda_1, \dots, \lambda_l) = \sum_{i_1 < i_2 < \dots < i_l} \lambda_{i_1} \lambda_{i_2} \dots \lambda_{i_l}.$$

Observe that for every  $A \in M_n$  with eigenvalues  $\{\lambda_1, \dots, \lambda_n\}$  we have

$$\prod_{j=1}^n (t - \lambda_j) = \chi_A(t) = \det(t\mathbb{1}_n - A) = t^n + \sum_{i=1}^n (-1)^i \sigma_i(A) t^{n-i}$$

Developing the determinant  $\det(t\mathbb{1}_n - A)$  we see that each of the coefficients  $\sigma_i(A)$  is in fact a *polynomial function* in the entries of  $A$ . A fortiori,  $\sigma_i(A)$  is a complex valued continuous function on  $M_n$ . The above equality also implies that the functions  $\sigma_i : M_n \rightarrow \mathbb{C}$  are constant along orbits. We now construct the continuous map

$$M_n \xrightarrow{\pi} \mathbb{C}^n$$

sending a matrix  $A \in M_n$  to the point  $(\sigma_1(A), \dots, \sigma_n(A))$  in  $\mathbb{C}^n$ . Clearly, if  $A \sim B$  then they map to the same point in  $\mathbb{C}^n$ . We claim that  $\pi$  is surjective. Take any point  $(a_1, \dots, a_n) \in \mathbb{C}^n$  and consider the matrix  $A \in M_n$

$$(1.1) \quad A = \begin{bmatrix} 0 & & & & a_n \\ -1 & 0 & & & a_{n-1} \\ & \ddots & \ddots & & \vdots \\ & & -1 & 0 & a_2 \\ & & & -1 & a_1 \end{bmatrix}$$

then we will show that  $\pi(A) = (a_1, \dots, a_n)$ , that is,

$$\det(t\mathbb{1}_n - A) = t^n - a_1 t^{n-1} + a_2 t^{n-2} - \dots + (-1)^n a_n.$$

Indeed, developing the determinant of  $tI_n - A$  along the first column we obtain

$$\begin{vmatrix} \textcircled{t} & 0 & 0 & \cdots & 0 & -a_n \\ 1 & t & 0 & & 0 & -a_{n-1} \\ 0 & 1 & t & & 0 & -a_{n-2} \\ \vdots & & \ddots & \ddots & \vdots & \vdots \\ 0 & 0 & & 1 & t & -a_2 \\ 0 & 0 & & & 1 & t - a_1 \end{vmatrix} - \begin{vmatrix} t & 0 & 0 & \cdots & 0 & -a_n \\ \textcircled{1} & t & 0 & & 0 & -a_{n-1} \\ 0 & 1 & t & & 0 & -a_{n-2} \\ \vdots & & \ddots & \ddots & \vdots & \vdots \\ 0 & 0 & & 1 & t & -a_2 \\ 0 & 0 & & & 1 & t - a_1 \end{vmatrix}$$

Here, the second determinant is equal to  $(-1)^{n-1}a_n$  and by induction on  $n$  the first determinant is equal to  $t.(t^{n-1} - a_1t^{n-2} + \dots + (-1)^{n-1}a_{n-1})$ , proving the claim.

Next, we will determine which  $n \times n$  matrices can be conjugated to a matrix in the canonical form  $A$  as above. We call a matrix  $B \in M_n$  *cyclic* if there is a (column) vector  $v \in \mathbb{C}^n$  such that  $\mathbb{C}^n$  is spanned by the vectors  $\{v, B.v, B^2.v, \dots, B^{n-1}.v\}$ . Let  $g \in GL_n$  be the basechange transforming the standard basis to the ordered basis

$$(v, -B.v, B^2.v, -B^3.v, \dots, (-1)^{n-1}B^{n-1}.v).$$

In this new basis, the linear map determined by  $B$  (or equivalently,  $g.B.g^{-1}$ ) is equal to the matrix in canonical form

$$\begin{bmatrix} 0 & & & & b_n \\ -1 & 0 & & & b_{n-1} \\ & \ddots & \ddots & & \vdots \\ & & -1 & 0 & b_2 \\ & & & -1 & b_1 \end{bmatrix}$$

where  $B^n.v$  has coordinates  $(b_n, \dots, b_2, b_1)$  in the new basis. Conversely, any matrix in this form is a cyclic matrix.

We claim that the set of all cyclic matrices in  $M_n$  is a *dense* open subset. To see this take  $v = (x_1, \dots, x_n)^T \in \mathbb{C}^n$  and compute the determinant of the  $n \times n$  matrix

$$\begin{bmatrix} | & | & & | \\ \hline v & Bv & \cdots & B^{n-1}.v \\ \hline | & | & & | \end{bmatrix}$$

This gives a polynomial of total degree  $n$  in the  $x_i$  with all its coefficients polynomial functions  $c_j$  in the entries  $b_{kl}$  of  $B$ . Now,  $B$  is a cyclic matrix if and only if at least one of these coefficients is non-zero. That is, the set of non-cyclic matrices is exactly the intersection of the finitely many *hypersurfaces*

$$V_j = \{B = (b_{kl})_{k,l} \in M_n \mid c_j(b_{11}, b_{12}, \dots, b_{nn}) = 0\}$$

in the vectorspace  $M_n$ .

**THEOREM 1.2.** *The best continuous approximation to the orbit space is given by the surjection*

$$M_n \xrightarrow{\pi} \mathbb{C}^n$$

mapping a matrix  $A \in M_n(\mathbb{C})$  to the  $n$ -tuple  $(\sigma_1(A), \dots, \sigma_n(A))$ .

Let  $f : M_n \rightarrow \mathbb{C}$  be a continuous function which is constant along conjugacy classes. We will show that  $f$  factors through  $\pi$ , that is,  $f$  is really a continuous function in the  $\sigma_i(A)$ . Consider the diagram

$$\begin{array}{ccc} M_n & \xrightarrow{f} & \mathbb{C} \\ \uparrow s & \searrow \pi & \nearrow f' = f \circ s \\ \mathbb{C}^n & & \end{array}$$

where  $s$  is the *section* of  $\pi$  (that is,  $\pi \circ s = id_{\mathbb{C}^n}$ ) determined by sending a point  $(a_1, \dots, a_n)$  to the cyclic matrix in canonical form  $A$  as in equation (1.1). Clearly,  $s$  is continuous, hence so is  $f' = f \circ s$ . The approximation property follows if we prove that  $f = f' \circ \pi$ . By continuity, it suffices to check equality on the dense open set of cyclic matrices in  $M_n$ .

There it is a consequence of the following three facts we have proved before : (1) : any cyclic matrix lies in the same orbit as one in standard form, (2) :  $s$  is a section of  $\pi$  and (3) :  $f$  is constant along orbits.

**EXAMPLE 1.3.** Orbits in  $M_2$ .

A  $2 \times 2$  matrix  $A$  can be conjugated to an upper triangular matrix with diagonal entries the eigenvalues  $\lambda_1, \lambda_2$  of  $A$ . As the trace and determinant of both matrices are equal we have


$$\sigma_1(A) = tr(A) \text{ and } \sigma_2(A) = det(A).$$

The best approximation to the orbit space is therefore given by the surjective map

$$M_2 \xrightarrow{\pi} \mathbb{C}^2 \quad \begin{bmatrix} a & b \\ c & d \end{bmatrix} \mapsto (a+d, ad-bc)$$

The matrix  $A$  has two equal eigenvalues if and only if the discriminant of the characteristic polynomial  $t^2 - \sigma_1(A)t + \sigma_2(A)$  is zero, that is when  $\sigma_1(A)^2 - 4\sigma_2(A) = 0$ . This condition determines a closed curve  $C$  in  $\mathbb{C}^2$  where

$$C = \{(x, y) \in \mathbb{C}^2 \mid x^2 - 4y = 0\}.$$








FIGURE 2. Orbit closures of  $2 \times 2$  matrices

Observe that  $C$  is a smooth 1-dimensional submanifold of  $\mathbb{C}^2$ . We will describe the *fibers* (that is, the inverse images of points) of the surjective map  $\pi$ .

If  $p = (x, y) \in \mathbb{C}^2 - C$ , then  $\pi^{-1}(p)$  consists of precisely one orbit (which is then necessarily closed in  $M_2$ ) namely that of the diagonal matrix

$$\begin{bmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{bmatrix} \quad \text{where} \quad \lambda_{1,2} = \frac{-x \pm \sqrt{x^2 - 4y}}{2}$$

If  $p = (x, y) \in C$  then  $\pi^{-1}(p)$  consists of two orbits,

$$\mathcal{O} \begin{bmatrix} \lambda & 1 \\ 0 & \lambda \end{bmatrix} \quad \text{and} \quad \mathcal{O} \begin{bmatrix} \lambda & 0 \\ 0 & \lambda \end{bmatrix}$$

where  $\lambda = \frac{1}{2}x$ . We have seen that the second orbit lies in the closure of the first. Observe that the second orbit reduces to one point in  $M_2$  and hence is closed. Hence, also  $\pi^{-1}(p)$  contains a unique closed orbit.

To describe the fibers of  $\pi$  as closed subsets of  $M_2$  it is convenient to write any matrix  $A$  as a linear combination

$$A = u(A) \begin{bmatrix} \frac{1}{2} & 0 \\ 0 & \frac{1}{2} \end{bmatrix} + v(A) \begin{bmatrix} \frac{1}{2} & 0 \\ 0 & -\frac{1}{2} \end{bmatrix} + w(A) \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} + z(A) \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}.$$


Expressed in the coordinate functions  $u, v, w$  and  $z$  the fibers  $\pi^{-1}(p)$  of a point  $p = (x, y) \in \mathbb{C}^2$  are the common zeroes of

$$\begin{cases} u & = x \\ v^2 + 4wz & = x^2 - 4y \end{cases}$$

The first equation determines a three dimensional affine subspace of  $M_2$  in which the second equation determines a *quadric*. If  $p \notin C$  this quadric is non-degenerate and thus  $\pi^{-1}(p)$  is a smooth 2-dimensional submanifold of  $M_2$ . If  $p \in C$ , the quadric is a cone with top lying in the point  $\frac{x}{2}\mathbb{1}_2$ . Under the  $GL_2$ -action, the unique singular point of the cone must be clearly fixed giving us the closed orbit of dimension 0 corresponding to the diagonal matrix. The other orbit is the complement of the top and hence is a smooth 2-dimensional (non-closed) submanifold of  $M_2$ . The graphs in figure 2 represent the orbit-closures and the dimensions of the orbits.

EXAMPLE 1.4. Orbits in  $M_3$ .

We will describe the fibers of the surjective map  $M_3 \xrightarrow{\pi} \mathbb{C}^3$ . If a  $3 \times 3$  matrix has multiple eigenvalues then the *discriminant*  $d = (\lambda_1 - \lambda_2)^2(\lambda_2 - \lambda_3)^2(\lambda_3 - \lambda_1)^2$

FIGURE 3. Representation strata for  $3 \times 3$  matrices.

is zero. Clearly,  $d$  is a symmetric polynomial and hence can be expressed in terms of  $\sigma_1, \sigma_2$  and  $\sigma_3$ . More precisely,

$$d = 4\sigma_1^3\sigma_3 + 4\sigma_2^3 + 27\sigma_3^2 - \sigma_1^2\sigma_2^2 - 18\sigma_1\sigma_2\sigma_3$$

The set of points in  $\mathbb{C}^3$  where  $d$  vanishes is a surface  $S$  with *singularities*. These singularities are the common zeroes of the  $\frac{\partial d}{\partial \sigma_i}$  for  $1 \leq i \leq 3$ . One computes that these singularities form a *twisted cubic curve*  $C$  in  $\mathbb{C}^3$ , that is,

$$C = \{(3c, 3c^2, c^3) \mid c \in \mathbb{C}\}.$$

The description of the fibers  $\pi^{-1}(p)$  for  $p = (x, y, z) \in \mathbb{C}^3$  is as follows. When  $p \notin S$ , then  $\pi^{-1}(p)$  consists of a unique orbit (which is therefore closed in  $M_3$ ), the conjugacy class of a matrix with paired distinct eigenvalues. If  $p \in S - C$ , then  $\pi^{-1}(p)$  consists of the orbits of

$$A_1 = \begin{bmatrix} \lambda & 1 & 0 \\ 0 & \lambda & 0 \\ 0 & 0 & \mu \end{bmatrix} \quad \text{and} \quad A_2 = \begin{bmatrix} \lambda & 0 & 0 \\ 0 & \lambda & 0 \\ 0 & 0 & \mu \end{bmatrix}$$


Finally, if  $p \in C$ , then the matrices in the fiber  $\pi^{-1}(p)$  have a single eigenvalue  $\lambda = \frac{1}{3}x$  and the fiber consists of the orbits of the matrices

$$B_1 = \begin{bmatrix} \lambda & 1 & 0 \\ 0 & \lambda & 1 \\ 0 & 0 & \lambda \end{bmatrix} \quad B_2 = \begin{bmatrix} \lambda & 1 & 0 \\ 0 & \lambda & 0 \\ 0 & 0 & \lambda \end{bmatrix} \quad B_3 = \begin{bmatrix} \lambda & 0 & 0 \\ 0 & \lambda & 0 \\ 0 & 0 & \lambda \end{bmatrix}$$

We observe that the *strata* with distinct fiber behavior (that is,  $\mathbb{C}^3 - S$ ,  $S - C$  and  $C$ ) are all submanifolds of  $\mathbb{C}^3$ , see figure 3.

The dimension of an orbit  $\mathcal{O}(A)$  in  $M_n$  is computed as follows. Let  $C_A$  be the subspace of all matrices in  $M_n$  commuting with  $A$ . Then, the *stabilizer* subgroup of  $A$  is a dense open subset of  $C_A$  whence the dimension of  $\mathcal{O}(A)$  is equal to  $n^2 - \dim C_A$ .

Performing these calculations for the matrices given above, we obtain the following graphs representing orbit-closures and the dimensions of orbits



Returning to  $M_n$ , the set of cyclic matrices is a Zariski open subset of  $M_n$ . For, consider the generic matrix of coordinate functions and generic column vector

$$X = \begin{bmatrix} x_{11} & \dots & x_{1n} \\ \vdots & & \vdots \\ x_{n1} & \dots & x_{nn} \end{bmatrix} \quad \text{and} \quad v = \begin{bmatrix} v_1 \\ \vdots \\ v_n \end{bmatrix}$$

and form the square matrix


$$[v \quad X.v \quad X^2.v \quad \dots \quad X^{n-1}.v] \in M_n(\mathbb{C}[x_{11}, x_{12}, \dots, x_{nn}, v_1, \dots, v_n])$$

Then its determinant can be written as  $\sum_{l=1}^z p_l(x_{ij})q_l(v_k)$  where the  $q_l$  are polynomials in the  $v_k$  and the  $p_l$  polynomials in the  $x_{ij}$ . Let  $A \in M_n$  be such that at least one of the  $p_l(A) \neq 0$ , then the polynomial  $d = \sum_l p_l(A)q_l(v_k) \in \mathbb{C}[v_1, \dots, v_k]$  is non-zero. But then there is a  $c = (c_1, \dots, c_n) \in \mathbb{C}^n$  such that  $d(c) \neq 0$  and hence  $c^\tau$  is a cyclic vector for  $A$ . The converse implication is obvious.

**THEOREM 1.5.** *Let  $f : M_n \longrightarrow \mathbb{C}$  is a regular (that is, polynomial) function on  $M_n$  which is constant along conjugacy classes, then*

$$f \in \mathbb{C}[\sigma_1(X), \dots, \sigma_n(X)]$$

**PROOF.** Consider again the diagram



The function  $f' = f \circ s$  is a regular function on  $\mathbb{C}^n$  whence is a polynomial in the coordinate functions of  $\mathbb{C}^n$  (which are the  $\sigma_i(X)$ ), so

$$f' \in \mathbb{C}[\sigma_1(X), \dots, \sigma_n(X)] \hookrightarrow \mathbb{C}[M_n].$$

Moreover,  $f$  and  $f'$  are equal on a Zariski open (dense) subset of  $M_n$  whence they are equal as polynomials in  $\mathbb{C}[M_n]$ .  $\square$

The ring of polynomial functions on  $M_n$  which are constant along conjugacy classes can also be viewed as a ring of invariants. The group  $GL_n$  acts as algebra automorphisms on the polynomial ring  $\mathbb{C}[M_n]$ . The automorphism  $\phi_g$  determined by  $g \in GL_n$  sends the variable  $x_{ij}$  to the  $(i, j)$ -entry of the matrix  $g^{-1}.X.g$  which

is a linear form in  $\mathbb{C}[M_n]$ . This action is determined by the property that for all  $g \in GL_n$ ,  $A \in A$  and  $f \in \mathbb{C}[M_n]$  we have that

$$\phi_g(f)(A) = f(g.A.g^{-1})$$

The *ring of polynomial invariants* is the algebra of polynomials left invariant under this action

$$\mathbb{C}[M_n]^{GL_n} = \{f \in \mathbb{C}[M_n] \mid \phi_g(f) = f \text{ for all } g \in GL_n\}$$

and hence is the ring of polynomial functions on  $M_n$  which are constant along orbits. The foregoing theorem determines the ring of polynomial invariants

$$\mathbb{C}[M_n]^{GL_n} = \mathbb{C}[\sigma_1(X), \dots, \sigma_n(X)]$$

We will give an equivalent description of this ring below.

Consider the variables  $\lambda_1, \dots, \lambda_n$  and consider the polynomial

$$f_n(t) = \prod_{i=1}^n (t - \lambda_i) = t^n + \sum_{i=1}^n (-1)^i \sigma_i t^{n-i}$$

then  $\sigma_i$  is the  $i$ -th elementary symmetric polynomial in the  $\lambda_j$ . We know that these polynomials are algebraically independent and generate the *ring of symmetric polynomials* in the  $\lambda_j$ , that is,

$$\mathbb{C}[\sigma_1, \dots, \sigma_n] = \mathbb{C}[\lambda_1, \dots, \lambda_n]^{S_n}$$

where  $S_n$  is the *symmetric group* on  $n$  letters acting by automorphisms on the polynomial ring  $\mathbb{C}[\lambda_1, \dots, \lambda_n]$  via  $\pi(\lambda_i) = \lambda_{\pi(i)}$  and the algebra of polynomials which are fixed under these automorphisms are precisely the symmetric polynomials in the  $\lambda_j$ .

Consider the symmetric Newton functions  $s_i = \lambda_1^i + \dots + \lambda_n^i$ , then we claim that this is another generating set of symmetric polynomials, that is,

$$\mathbb{C}[\sigma_1, \dots, \sigma_n] = \mathbb{C}[s_1, \dots, s_n].$$

To prove this it suffices to express each  $\sigma_i$  as a polynomial in the  $s_j$ . More precisely, we claim that the following identities hold for all  $1 \leq j \leq n$

$$(1.2) \quad s_j - \sigma_1 s_{j-1} + \sigma_2 s_{j-2} - \dots + (-1)^{j-1} \sigma_{j-1} s_1 + (-1)^j \sigma_j = 0$$

For  $j = n$  this identity holds because we have

$$0 = \sum_{i=1}^n f_n(\lambda_i) = s_n + \sum_{i=1}^n (-1)^i \sigma_i s_{n-i}$$

if we take  $s_0 = n$ . Assume now  $j < n$  then the left hand side of equation 1.2 is a symmetric function in the  $\lambda_i$  of degree  $\leq j$  and is therefore a polynomial  $p(\sigma_1, \dots, \sigma_j)$  in the first  $j$  elementary symmetric polynomials. Let  $\phi$  be the algebra epimorphism

$$\mathbb{C}[\lambda_1, \dots, \lambda_n] \xrightarrow{\phi} \mathbb{C}[\lambda_1, \dots, \lambda_j]$$

defined by mapping  $\lambda_{j+1}, \dots, \lambda_j$  to zero. Clearly,  $\phi(\sigma_i)$  is the  $i$ -th elementary symmetric polynomial in  $\{\lambda_1, \dots, \lambda_j\}$  and  $\phi(s_i) = \lambda_1^i + \dots + \lambda_j^i$ . Repeating the above  $j = n$  argument (replacing  $n$  by  $j$ ) we have

$$0 = \sum_{i=1}^j f_j(\lambda_i) = \phi(s_j) + \sum_{i=1}^j (-1)^i \phi(\sigma_i) \phi(s_{n-i})$$

(this time with  $s_0 = j$ ). But then,  $p(\phi(\sigma_1), \dots, \phi(\sigma_j)) = 0$  and as the  $\phi(\sigma_k)$  for  $1 \leq k \leq j$  are algebraically independent we must have that  $p$  is the zero polynomial finishing the proof of the claimed identity.

If  $\lambda_1, \dots, \lambda_n$  are the eigenvalues of an  $n \times n$  matrix  $A$ , then  $A$  can be conjugated to an upper triangular matrix  $B$  with diagonal entries  $(\lambda_1, \dots, \lambda_n)$ . Hence, the trace  $tr(A) = tr(B) = \lambda_1 + \dots + \lambda_n = s_1$ . In general,  $A^i$  can be conjugated to  $B^i$  which is an upper triangular matrix with diagonal entries  $(\lambda_1^i, \dots, \lambda_n^i)$  and hence the traces of  $A^i$  and  $B^i$  are equal to  $\lambda_1^i + \dots + \lambda_n^i = s_i$ . Concluding, we have

**THEOREM 1.6.** *Consider the action of conjugation by  $GL_n$  on  $M_n$ . Let  $X$  be the generic matrix of coordinate functions on  $M_n$*

$$X = \begin{bmatrix} x_{11} & \dots & x_{nn} \\ \vdots & & \vdots \\ x_{n1} & \dots & x_{nn} \end{bmatrix}$$

*Then, the ring of polynomial invariants is generated by the traces of powers of  $X$ , that is,*

$$\mathbb{C}[M_n]^{GL_n} = \mathbb{C}[tr(X), tr(X^2), \dots, tr(X^n)]$$

**PROOF.** The result follows from theorem 1.5 and the fact that

$$\mathbb{C}[\sigma_1(X), \dots, \sigma_n(X)] = \mathbb{C}[tr(X), \dots, tr(X^n)]$$

□

## 1.2. Simultaneous conjugacy classes.

It will be crucial to extend what we have done for conjugacy classes of matrices to simultaneous conjugacy classes of  $m$ -tuples of matrices. Consider the  $mn^2$ -dimensional complex vectorspace

$$M_n^m = \underbrace{M_n \oplus \dots \oplus M_n}_m$$

of  $m$ -tuples  $(A_1, \dots, A_m)$  of  $n \times n$ -matrices  $A_i \in M_n$ . On this space we let the group  $GL_n$  act by simultaneous conjugation, that is

$$g.(A_1, \dots, A_m) = (g.A_1.g^{-1}, \dots, g.A_m.g^{-1})$$

for all  $g \in GL_n$  and all  $m$ -tuples  $(A_1, \dots, A_m)$ . Unfortunately, there is no substitute for the Jordan normalform result in this more general setting. Still, for small  $m$  and  $n$  one can work out the  $GL_n$ -orbits by ad hoc methods.

**EXAMPLE 1.7.** Orbits in  $M_2^2 = M_2 \oplus M_2$ .

We can try to mimic the geometric approach to the conjugacy class problem, that is, we will try to approximate the orbitspace via polynomial functions on  $M_2^2$  which are constant along orbits. For  $(A, B) \in M_2^2 = M_2 \oplus M_2$  clearly the polynomial functions we have encountered before  $tr(A), det(A)$  and  $tr(B), det(B)$  are constant along orbits. However, there are more: for example  $tr(AB)$ . Later, we will show that these five functions generate all polynomial functions which are constant along orbits. Here, we will show that the map  $M_2^2 = M_2 \oplus M_2 \xrightarrow{\pi} \mathbb{C}^5$  defined by

$$(A, B) \mapsto (tr(A), det(A), tr(B), det(B), tr(AB))$$

is surjective such that each fiber contains precisely one closed orbit. In the next chapter, we will see that this property characterizes the best polynomial approximation to the (non-existent) orbit space.

First, we will show surjectivity of  $\pi$ , that is, for every  $(x_1, \dots, x_5) \in \mathbb{C}^5$  we will construct a couple of  $2 \times 2$  matrices  $(A, B)$  (or rather its orbit) such that  $\pi(A, B) = (x_1, \dots, x_5)$ . Consider the open set where  $x_1^2 \neq 4x_2$ . We have seen that this property characterizes those  $A \in M_2$  such that  $A$  has distinct eigenvalues and hence diagonalizable. Hence, we can take a representative of the orbit  $\mathcal{O}(A, B)$  to be a couple

$$\left( \begin{bmatrix} \lambda & 0 \\ 0 & \mu \end{bmatrix}, \begin{bmatrix} c_1 & c_2 \\ c_3 & c_4 \end{bmatrix} \right)$$

with  $\lambda \neq \mu$ . We need a solution to the set of equations

$$\begin{cases} x_3 = c_1 + c_4 \\ x_4 = c_1 c_4 - c_2 c_3 \\ x_5 = \lambda c_1 + \mu c_4 \end{cases}$$

Because  $\lambda \neq \mu$  the first and last equation uniquely determine  $c_1, c_4$  and substitution in the second gives us  $c_2 c_3$ . Analogously, points of  $\mathbb{C}^5$  lying in the open set  $x_3^2 \neq x_4$  lie in the image of  $\pi$ . Finally, for a point in the complement of these open sets, that is when  $x_1^2 = x_2$  and  $x_3^2 = 4x_4$  we can consider a couple  $(A, B)$

$$\left( \begin{bmatrix} \lambda & 1 \\ 0 & \lambda \end{bmatrix}, \begin{bmatrix} \mu & 0 \\ c & \mu \end{bmatrix} \right)$$

where  $\lambda = \frac{1}{2}x_1$  and  $\mu = \frac{1}{2}x_3$ . Observe that the remaining equation  $x_5 = \text{tr}(AB) = 2\lambda\mu + c$  has a solution in  $c$ .

Now, we will describe the fibers of  $\pi$ . Assume  $(A, B)$  is such that  $A$  and  $B$  have a common eigenvector  $v$ . Simultaneous conjugation with a  $g \in GL_n$  expressing a basechange from the standard basis to  $\{v, w\}$  for some  $w$  shows that the orbit  $\mathcal{O}(A, B)$  contains a couple of upper-triangular matrices. We want to describe the image of these matrices under  $\pi$ . Take an upper triangular representative in  $\mathcal{O}(A, B)$

$$\left( \begin{bmatrix} a_1 & a_2 \\ 0 & a_3 \end{bmatrix}, \begin{bmatrix} b_1 & b_2 \\ 0 & b_3 \end{bmatrix} \right).$$

with  $\pi$ -image  $(x_1, \dots, x_5)$ . The coordinates  $x_1, x_2$  determine the eigenvalues  $a_1, a_3$  of  $A$  only as an unordered set (similarly,  $x_3, x_4$  only determine the set of eigenvalues  $\{b_1, b_3\}$  of  $B$ ). Hence,  $\text{tr}(AB)$  is one of the following two expressions

$$a_1 b_1 + a_3 b_3 \quad \text{or} \quad a_1 b_3 + a_3 b_1$$

and therefore satisfies the equation

$$(\text{tr}(AB) - a_1 b_1 - a_3 b_3)(\text{tr}(AB) - a_1 b_3 - a_3 b_1) = 0.$$

Recall that  $x_1 = a_1 + a_3, x_2 = a_1 a_3, x_3 = b_1 + b_3, x_4 = b_1 b_3$  and  $x_5 = \text{tr}(AB)$  we can express this equation as

$$x_5^2 - x_1 x_3 x_5 + x_1^2 x_4 + x_3^2 x_2 - 4x_2 x_4 = 0.$$

This determines an hypersurface  $H \hookrightarrow \mathbb{C}^5$ . If we view the left-hand side as a polynomial  $f$  in the coordinate functions of  $\mathbb{C}^5$  we see that  $H$  is a four dimensional

subset of  $\mathbb{C}^5$  with singularities the common zeroes of the partial derivatives

$$\frac{\partial f}{\partial x_i} \text{ for } 1 \leq i \leq 5$$

These singularities for the 2-dimensional submanifold  $S$  of points of the form  $(2a, a^2, 2b, b^2, 2ab)$ . We now claim that the smooth submanifolds  $\mathbb{C}^5 - H$ ,  $H - S$  and  $S$  of  $\mathbb{C}^5$  describe the different types of fiber behavior. In chapter 6 we will see that the subsets of points with different fiber behavior (actually, of different representation type) are manifolds for  $m$ -tuples of  $n \times n$  matrices.

If  $p \notin H$  we claim that  $\pi^{-1}(p)$  is a unique orbit, which is therefore closed in  $M_2^2$ . Let  $(A, B) \in \pi^{-1}$  and assume first that  $x_1^2 \neq 4x_2$  then there is a representative in  $\mathcal{O}(A, B)$  of the form

$$\left( \begin{bmatrix} \lambda & 0 \\ 0 & \mu \end{bmatrix}, \begin{bmatrix} c_1 & c_2 \\ c_3 & c_4 \end{bmatrix} \right)$$

with  $\lambda \neq \mu$ . Moreover,  $c_2 c_3 \neq 0$  (for otherwise  $A$  and  $B$  would have a common eigenvector whence  $p \in H$ ) hence we may assume that  $c_2 = 1$  (eventually after simultaneous conjugation with a suitable diagonal matrix  $\text{diag}(t, t^{-1})$ ). The value of  $\lambda, \mu$  is determined by  $x_1, x_2$ . Moreover,  $c_1, c_3, c_4$  are also completely determined by the system of equations

$$\begin{cases} x_3 &= c_1 + c_4 \\ x_4 &= c_1 c_4 - c_3 \\ x_5 &= \lambda c_1 + \mu c_4 \end{cases}$$

and hence the point  $p = (x_1, \dots, x_5)$  completely determines the orbit  $\mathcal{O}(A, B)$ . Remains to consider the case when  $x_1^2 = 4x_2$  (that is, when  $A$  has a single eigenvalue). Consider the couple  $(uA + vB, B)$  for  $u, v \in \mathbb{C}^*$ . To begin,  $uA + vB$  and  $B$  do not have a common eigenvalue. Moreover,  $p = \pi(A, B)$  determines  $\pi(uA + vB, B)$  as

$$\begin{cases} \text{tr}(uA + vB) &= \text{utr}(A) + \text{vtr}(B) \\ \det(uA + vB) &= u^2 \det(A) + v^2 \det(B) + uv(\text{tr}(A)\text{tr}(B) - \text{tr}(AB)) \\ \text{tr}((uA + vB)B) &= \text{utr}(AB) + v(\text{tr}(B)^2 - 2\det(B)) \end{cases}$$

Assume that for all  $u, v \in \mathbb{C}^*$  we have the equality  $\text{tr}(uA + vB)^2 = 4\det(uA + vB)$  then comparing coefficients of this equation expressed as a polynomial in  $u$  and  $v$  we obtain the conditions  $x_1^2 = 4x_2$ ,  $x_3^2 = 4x_4$  and  $2x_5 = x_1 x_3$  whence  $p \in S \hookrightarrow H$ , a contradiction. So, fix  $u, v$  such that  $uA + vB$  has distinct eigenvalues. By the above argument  $\mathcal{O}(uA + vB, B)$  is the unique orbit lying over  $\pi(uA + vB, B)$ , but then  $\mathcal{O}(A, B)$  must be the unique orbit lying over  $p$ .

Let  $p \in H - S$  and  $(A, B) \in \pi^{-1}(p)$ , then  $A$  and  $B$  are simultaneous upper triangularizable, with eigenvalues  $a_1, a_2$  respectively  $b_1, b_2$ . Either  $a_1 \neq a_2$  or  $b_1 \neq b_2$  for otherwise  $p \in S$ . Assume  $a_1 \neq a_2$ , then there is a representative in the orbit  $\mathcal{O}(A, B)$  of the form

$$\left( \begin{bmatrix} a_i & 0 \\ 0 & a_j \end{bmatrix}, \begin{bmatrix} b_k & b \\ 0 & b_l \end{bmatrix} \right)$$

for  $\{i, j\} = \{1, 2\} = \{k, l\}$ . If  $b \neq 0$  we can conjugate with a suitable diagonal matrix to get  $b = 1$  hence we get at most 9 possible orbits. Checking all possibilities we see that only three of them are distinct, those corresponding to the couples

$$\left( \begin{bmatrix} a_1 & 0 \\ 0 & a_2 \end{bmatrix}, \begin{bmatrix} b_1 & 1 \\ 0 & b_2 \end{bmatrix} \right) \quad \left( \begin{bmatrix} a_1 & 0 \\ 0 & a_2 \end{bmatrix}, \begin{bmatrix} b_1 & 0 \\ 0 & b_2 \end{bmatrix} \right) \quad \left( \begin{bmatrix} a_2 & 0 \\ 0 & a_1 \end{bmatrix}, \begin{bmatrix} b_1 & 1 \\ 0 & b_2 \end{bmatrix} \right)$$

Clearly, the first and last orbit have the middle one lying in its closure. Observe that the case assuming that  $b_1 \neq b_2$  is handled similarly. Hence, if  $p \in H - S$  then  $\pi^{-1}(p)$  consists of three orbits, two of dimension three whose closures intersect in a (closed) orbit of dimension two.

Finally, consider the case when  $p \in S$  and  $(A, B) \in \pi^{-1}(p)$ . Then, both  $A$  and  $B$  have a single eigenvalue and the orbit  $\mathcal{O}(A, B)$  has a representative of the form

$$\left( \begin{bmatrix} a & x \\ 0 & a \end{bmatrix}, \begin{bmatrix} b & y \\ 0 & b \end{bmatrix} \right)$$

for certain  $x, y \in \mathbb{C}$ . If either  $x$  or  $y$  are non-zero, then the subgroup of  $GL_2$  fixing this matrix consists of the matrices of the form


$$\text{Stab} \begin{bmatrix} c & 1 \\ 0 & c \end{bmatrix} = \left\{ \begin{bmatrix} u & v \\ 0 & u \end{bmatrix} \mid u \in \mathbb{C}^*, v \in \mathbb{C} \right\}$$

but these matrices also fix the second component. Therefore, if either  $x$  or  $y$  is nonzero, the orbit is fully determined by  $[x : y] \in \mathbb{P}^1$ . That is, for  $p \in S$ , the fiber  $\pi^{-1}(p)$  consists of an infinite family of orbits of dimension 2 parameterized by the points of the projective line  $\mathbb{P}^1$  together with the orbit of

$$\left( \begin{bmatrix} a & 0 \\ 0 & a \end{bmatrix}, \begin{bmatrix} b & 0 \\ 0 & b \end{bmatrix} \right)$$

which consists of one point (hence is closed in  $M_2^2$ ) and lies in the closure of each of the 2-dimensional orbits.

Concluding, we see that each fiber  $\pi^{-1}(p)$  contains a unique closed orbit (that of minimal dimension). The orbitclosure and dimension diagrams have the following shapes



### 1.3. Matrix invariants and necklaces

In this section we will determine the ring of all polynomial maps

$$M_n^m = \underbrace{M_n \oplus \dots \oplus M_n}_m \xrightarrow{f} \mathbb{C}$$

which are constant along orbits under the action of  $GL_n$  on  $M_n^m$  by simultaneous conjugation. The strategy we will use is classical in invariant theory.

- First, we will determine the *multilinear* maps which are constant along orbits, equivalently, the *linear* maps

$$M_n^{\otimes m} = \underbrace{M_n \otimes \dots \otimes M_n}_m \longrightarrow \mathbb{C}$$



which are constant along  $GL_n$ -orbits where  $GL_n$  acts by the diagonal action, that is,

$$g.(A_1 \otimes \dots \otimes A_m) = gA_1g^{-1} \otimes \dots \otimes gA_mg^{-1}.$$

- Afterwards, we will be able to obtain from them all polynomial invariant maps by using *polarization* and *restitution* operations.

First, we will translate our problem into one studied in classical invariant theory of  $GL_n$ .

Let  $V_n \simeq \mathbb{C}^n$  be the  $n$ -dimensional vectorspace of column vectors on which  $GL_n$  acts naturally by left multiplication

$$V_n = \begin{bmatrix} \mathbb{C} \\ \mathbb{C} \\ \vdots \\ \mathbb{C} \end{bmatrix} \quad \text{with action} \quad g. \begin{bmatrix} \nu_1 \\ \nu_2 \\ \vdots \\ \nu_n \end{bmatrix}$$

In order to define an action on the dual space  $V_n^* = \text{Hom}(V_n, \mathbb{C}) \simeq \mathbb{C}^n$  of *covectors* (or, row vectors) we have to use the *contragradient* action

$$V_n^* = [\mathbb{C} \quad \mathbb{C} \quad \dots \quad \mathbb{C}] \quad \text{with action} \quad [\phi_1 \quad \phi_2 \quad \dots \quad \phi_n] . g^{-1}$$

Observe, that we have an *evaluation* map  $V_n^* \times V_n \longrightarrow \mathbb{C}$  which is given by the scalar product  $f(v)$  for all  $f \in V_n^*$  and  $v \in V_n$

$$[\phi_1 \quad \phi_2 \quad \dots \quad \phi_n] \cdot \begin{bmatrix} \nu_1 \\ \nu_2 \\ \vdots \\ \nu_n \end{bmatrix} = \phi_1\nu_1 + \phi_2\nu_2 + \dots + \phi_n\nu_n$$

which is invariant under the diagonal action of  $GL_n$  on  $V_n^* \times V_n$ . Further, we have the natural identification

$$M_n = V_n \otimes V_n^* = \begin{bmatrix} \mathbb{C} \\ \mathbb{C} \\ \vdots \\ \mathbb{C} \end{bmatrix} \otimes [\mathbb{C} \quad \mathbb{C} \quad \dots \quad \mathbb{C}].$$

Under this identification, a *pure tensor*  $v \otimes f$  corresponds to the rank one matrix or rank one endomorphism of  $V_n$  defined by

$$v \otimes f : V_n \longrightarrow V_n \quad \text{with} \quad w \mapsto f(w)v$$

and observe that the rank one matrices span  $M_n$ . The diagonal action of  $GL_n$  on  $V_n \otimes V_n^*$  is then determined by its action on the pure tensors where it is equal to

$$g. \begin{bmatrix} \nu_1 \\ \nu_2 \\ \dots \\ \nu_n \end{bmatrix} \otimes [\phi_1 \quad \phi_2 \quad \dots \quad \phi_n] . g^{-1}$$

and therefore coincides with the action of conjugation on  $M_n$ . Now, let us consider the identification

$$(V_n^{*\otimes m} \otimes V_n^{\otimes m})^* \simeq \text{End}(V_n^{\otimes m})$$

obtained from the nondegenerate pairing

$$\text{End}(V_n^{\otimes m}) \times (V_n^{*\otimes m} \otimes V_n^{\otimes m}) \longrightarrow \mathbb{C}$$

given by the formula

$$\langle \lambda, f_1 \otimes \dots \otimes f_m \otimes v_1 \otimes \dots \otimes v_m \rangle = f_1 \otimes \dots \otimes f_m(\lambda(v_1 \otimes \dots \otimes v_m))$$

$GL_n$  acts diagonally on  $V_n^{\otimes m}$  and hence again by conjugation on  $\text{End}(V_n^{\otimes m})$  after embedding  $GL_n \hookrightarrow GL(V_n^{\otimes m}) = GL_{mn}$ . Thus, the above identifications are isomorphism as vectorspaces with  $GL_n$ -action. But then, the space of  $GL_n$ -invariant linear maps

$$V_n^{*\otimes m} \otimes V_n^{\otimes m} \longrightarrow \mathbb{C}$$

can be identified with the space  $\text{End}_{GL_n}(V_n^{\otimes m})$  of  $GL_n$ -linear endomorphisms of  $V_n^{\otimes m}$ . We will now give a different presentation of this vector space relating it to the symmetric group.

Apart from the diagonal action of  $GL_n$  on  $V_n^{\otimes m}$  given by

$$g.(v_1 \otimes \dots \otimes v_m) = g.v_1 \otimes \dots \otimes g.v_m$$

we have an action of the symmetric group  $S_m$  on  $m$  letters on  $V_n^{\otimes m}$  given by

$$\sigma.(v_1 \otimes \dots \otimes v_m) = v_{\sigma(1)} \otimes \dots \otimes v_{\sigma(m)}$$

These two actions commute with each other and give embeddings of  $GL_n$  and  $S_m$  in  $\text{End}(V_n^{\otimes m})$ . The subspace of  $V_n^{\otimes m}$  spanned by the image of  $GL_n$  will be denoted by  $\langle GL_n \rangle$ . Similarly, with  $\langle S_m \rangle$  we denote the subspace spanned by the image of  $S_m$ .

**THEOREM 1.8.** *With notations as above we have :*

- (1)  $\langle GL_n \rangle = \text{End}_{S_m}(V_n^{\otimes m})$
- (2)  $\langle S_m \rangle = \text{End}_{GL_n}(V_n^{\otimes m})$

**PROOF.** (1) : Under the identification  $\text{End}(V_n^{\otimes m}) = \text{End}(V_n)^{\otimes m}$  an element  $g \in GL_n$  is mapped to the symmetric tensor  $g \otimes \dots \otimes g$ . On the other hand, the image of  $\text{End}_{S_m}(V_n^{\otimes m})$  in  $\text{End}(V_n)^{\otimes m}$  is the subspace of all symmetric tensors in  $\text{End}(V)^{\otimes m}$ . We can give a basis of this subspace as follows. Let  $\{e_1, \dots, e_{n^2}\}$  be a basis of  $\text{End}(V_n)$ , then the vectors  $e_{i_1} \otimes \dots \otimes e_{i_m}$  form a basis of  $\text{End}(V_n)^{\otimes m}$  which is stable under the  $S_m$ -action. Further, any  $S_m$ -orbit contains a unique representative of the form

$$e_1^{\otimes h_1} \otimes \dots \otimes e_{n^2}^{\otimes h_{n^2}}$$

with  $h_1 + \dots + h_{n^2} = m$ . If we denote by  $r(h_1, \dots, h_{n^2})$  the sum of all elements in the corresponding  $S_m$ -orbit then these vectors are a basis of the symmetric tensors in  $\text{End}(V_n)^{\otimes m}$ .

The claim follows if we can show that every linear map  $\lambda$  on the symmetric tensors which is zero on all  $g \otimes \dots \otimes g$  with  $g \in GL_n$  is the zero map. Write  $e = \sum x_i e_i$ , then

$$\lambda(e \otimes \dots \otimes e) = \sum x_1^{h_1} \dots x_{n^2}^{h_{n^2}} \lambda(r(h_1, \dots, h_{n^2}))$$

is a polynomial function on  $\text{End}(V_n)$ . As  $GL_n$  is a Zariski open subset of  $\text{End}(V)$  on which by assumption this polynomial vanishes, it must be the zero polynomial. Therefore,  $\lambda(r(h_1, \dots, h_{n^2})) = 0$  for all  $(h_1, \dots, h_{n^2})$  finishing the proof.

(2) : Recall that the group algebra  $\mathbb{C}S_m$  of  $S_m$  is a *semisimple algebra*. Any epimorphic image of a semisimple algebra is semisimple. Therefore,  $\langle S_m \rangle$  is a

semisimple subalgebra of the matrixalgebra  $End(V_n^{\otimes m}) \simeq M_{nm}$ . By the *double centralizer theorem* (see for example [63]), it is therefore equal to the centralizer of  $End_{S_m}(V_m^{\otimes m})$ . By the first part, it is the centralizer of  $\langle GL_n \rangle$  in  $End(V_n^{\otimes m})$  and therefore equal to  $End_{GL_n}(V_n^{\otimes m})$ .  $\square$

Because  $End_{GL_n}(V_n^{\otimes m}) = \langle S_m \rangle$ , every  $GL_n$ -endomorphism of  $V_n^{\otimes m}$  can be written as a linear combination of the morphisms  $\lambda_\sigma$  describing the action of  $\sigma \in S_m$  on  $V_n^{\otimes m}$ . Our next job is to trace back these morphisms  $\lambda_\sigma$  through the canonical identifications until we can express them in terms of matrices.

To start let us compute the linear invariant

$$\mu_\sigma : V_n^{*\otimes m} \otimes V_n^{\otimes m} \longrightarrow \mathbb{C}$$

corresponding to  $\lambda_\sigma$  under the identification  $(V_n^{*\otimes m} \otimes V_n^{\otimes m})^* \simeq End(V_n^{\otimes m})$ . By the identification we know that  $\mu_\sigma(f_1 \otimes \dots \otimes f_m \otimes v_1 \otimes \dots \otimes v_m)$  is equal to

$$\begin{aligned} \langle \lambda_\sigma, f_1 \otimes \dots \otimes f_m \otimes v_1 \otimes \dots \otimes v_m \rangle &= f_1 \otimes \dots \otimes f_m(v_{\sigma(1)} \otimes \dots \otimes v_{\sigma(m)}) \\ &= \prod_i f_i(v_{\sigma(i)}) \end{aligned}$$

That is, we have proved

PROPOSITION 1.9. *Any multilinear  $GL_n$ -invariant map*

$$\gamma : V_n^{*\otimes m} \otimes V_n^{\otimes m} \longrightarrow \mathbb{C}$$

*is a linear combination of the invariants*

$$\mu_\sigma(f_1 \otimes \dots \otimes f_m \otimes v_1 \otimes \dots \otimes v_m) = \prod_i f_i(v_{\sigma(i)})$$

for  $\sigma \in S_m$ .

Using the identification  $M_n(\mathbb{C}) = V_n \otimes V_n^{*\otimes}$  a multilinear  $GL_n$ -invariant map

$$(V_n^* \otimes V_n)^{\otimes m} = V_n^{*\otimes m} \otimes V_n^{\otimes m} \longrightarrow \mathbb{C}$$

corresponds to a multilinear  $GL_n$ -invariant map

$$M_n(\mathbb{C}) \otimes \dots \otimes M_n(\mathbb{C}) \longrightarrow \mathbb{C}$$

We will now give a description of the generating maps  $\mu_\sigma$  in terms of matrices. Under the identification, matrix multiplication is induced by composition on rank one endomorphisms and here the rule is given by

$$v \otimes f.v' \otimes f' = f(v')v \otimes f'$$

$$\begin{bmatrix} \nu_1 \\ \vdots \\ \nu_n \end{bmatrix} \otimes [\phi_1 \quad \dots \quad \phi_n] \cdot \begin{bmatrix} \nu'_1 \\ \vdots \\ \nu'_n \end{bmatrix} \otimes [\phi'_1 \quad \dots \quad \phi'_n] = \begin{bmatrix} \nu_1 \\ \vdots \\ \nu_n \end{bmatrix} f(v') \otimes [\phi'_1 \quad \dots \quad \phi'_n].$$

Moreover, the trace map on  $M_n$  is induced by that on rank one endomorphisms where it is given by the rule

$$tr(v \otimes f) = f(v)$$

$$tr\left(\begin{bmatrix} \nu_1 \\ \vdots \\ \nu_n \end{bmatrix} \otimes [\phi_1 \quad \dots \quad \phi_n]\right) = tr\left(\begin{bmatrix} \nu_1\phi_1 & \dots & \nu_1\phi_n \\ \vdots & \ddots & \vdots \\ \nu_n\phi_1 & \dots & \nu_n\phi_n \end{bmatrix}\right) = \sum_i \nu_i\phi_i = f(v)$$

With these rules we can now give a matrix-interpretation of the  $GL_n$ -invariant maps  $\mu_\sigma$ .

PROPOSITION 1.10. *Let  $\sigma = (i_1 i_2 \dots i_\alpha)(j_1 j_2 \dots j_\beta) \dots (z_1 z_2 \dots z_\zeta)$  be a decomposition of  $\sigma \in S_m$  into cycles (including those of length one). Then, under the above identification we have*

$$\mu_\sigma(A_1 \otimes \dots \otimes A_m) = \text{tr}(A_{i_1} A_{i_2} \dots A_{i_\alpha}) \text{tr}(A_{j_1} A_{j_2} \dots A_{j_\beta}) \dots \text{tr}(A_{z_1} A_{z_2} \dots A_{z_\zeta})$$

PROOF. Both sides are multilinear hence it suffices to verify the equality for rank one matrices. Write  $A_i = v_i \otimes f_i$ , then we have that

$$\begin{aligned} \mu_\sigma(A_1 \otimes \dots \otimes A_m) &= \mu_\sigma(v_1 \otimes \dots \otimes v_m \otimes f_1 \otimes \dots \otimes f_m) \\ &= \prod_i f_i(v_{\sigma(i)}) \end{aligned}$$

Consider the subproduct

$$f_{i_1}(v_{i_2}) f_{i_2}(v_{i_3}) \dots f_{i_{\alpha-1}}(v_{i_\alpha}) = S$$

Now, look at the matrixproduct

$$v_{i_1} \otimes f_{i_1} \cdot v_{i_2} \otimes f_{i_2} \cdot \dots \cdot v_{i_\alpha} \otimes f_{i_\alpha}$$

which is by the product rule equal to

$$f_{i_1}(v_{i_2}) f_{i_2}(v_{i_3}) \dots f_{i_{\alpha-1}}(v_{i_\alpha}) v_{i_1} \otimes f_{i_\alpha}$$

Hence, by the trace rule we have that

$$\text{tr}(A_{i_1} A_{i_2} \dots A_{i_\alpha}) = \prod_{j=1}^{\alpha} f_{i_j}(v_{\sigma(i_j)}) = S$$

□

Having found a description of the multilinear invariant polynomial maps

$$M_n^m = \underbrace{M_n \oplus \dots \oplus M_n}_m \longrightarrow \mathbb{C}$$

we will now describe all polynomial maps which are constant along orbits by polarization. The coordinate algebra  $\mathbb{C}[M_n^m]$  is the polynomial ring in  $mn^2$  variables  $x_{ij}(k)$  where  $1 \leq k \leq m$  and  $1 \leq i, j \leq n$ . Consider the  $m$  generic  $n \times n$  matrices

$$\boxed{k} = X_k = \begin{bmatrix} x_{11}(k) & \dots & x_{1n}(k) \\ \vdots & & \vdots \\ x_{n1}(k) & \dots & x_{nn}(k) \end{bmatrix} \in M_n(\mathbb{C}[M_n^m]).$$

The action of  $GL_n$  on polynomial maps  $f \in \mathbb{C}[M_n^m]$  is fully determined by the action on the coordinate functions  $x_{ij}(k)$ . As in the case of one  $n \times n$  matrix we see that this action is given by

$$g \cdot x_{ij}(k) = (g^{-1} \cdot X_k \cdot g)_{ij}.$$

We see that this action preserves the subspaces spanned by the entries of any of the generic matrices. Hence, we can define a gradation on  $\mathbb{C}[M_n^m]$  by  $\text{deg}(x_{ij}(k)) = (0, \dots, 0, 1, 0, \dots, 0)$  (with 1 at place  $k$ ) and decompose

$$\mathbb{C}[M_n^m] = \bigoplus_{(d_1, \dots, d_m) \in \mathbb{N}^m} \mathbb{C}[M_n^m]_{(d_1, \dots, d_m)}$$

where  $\mathbb{C}[M_n^m]_{(d_1, \dots, d_m)}$  is the subspace of all multihomogeneous forms  $f$  in the  $x_{ij}(k)$  of degree  $(d_1, \dots, d_m)$ , that is, in each monomial term of  $f$  there are exactly

$d_k$  factors coming from the entries of the generic matrix  $X_k$  for all  $1 \leq k \leq m$ . The action of  $GL_n$  stabilizes each of these subspaces, that is,

$$\text{if } f \in \mathbb{C}[M_n^m]_{(d_1, \dots, d_m)} \quad \text{then } g.f \in \mathbb{C}[M_n^m]_{(d_1, \dots, d_m)} \quad \text{for all } g \in GL_n.$$

In particular, if  $f$  determines a polynomial map on  $M_n^m$  which is constant along orbits, that is, if  $f$  belongs to the ring of invariants  $\mathbb{C}[M_n^m]^{GL_n}$  then each of its multihomogeneous components is also an invariant and therefore it suffices to determine all multihomogeneous invariants.

Let  $f \in \mathbb{C}[M_n^m]_{(d_1, \dots, d_m)}$  and take for each  $1 \leq k \leq m$   $d_k$  new variables  $t_1(k), \dots, t_{d_k}(k)$ . Expand

$$f(t_1(1)A_1(1) + \dots + t_{d_1}A_{d_1}(1), \dots, t_1(m)A_1(m) + \dots + t_{d_m}(m)A_{d_m}(m))$$

as a polynomial in the variables  $t_i(k)$ , then we get an expression

$$\sum t_1(1)^{s_1(1)} \dots t_{d_1}^{s_{d_1}(1)} \dots t_1(m)^{s_1(m)} \dots t_{d_m}(m)^{s_{d_m}(m)} \cdot f_{(s_1(1), \dots, s_{d_1}(1), \dots, s_1(m), \dots, s_{d_m}(m))}(A_1(1), \dots, A_{d_1}(1), \dots, A_1(m), \dots, A_{d_m}(m))$$

such that for all  $1 \leq k \leq m$  we have  $\sum_{i=1}^{d_k} s_i(k) = d_k$ . Moreover, each of the  $f_{(s_1(1), \dots, s_{d_1}(1), \dots, s_1(m), \dots, s_{d_m}(m))}$  is a multi-homogeneous polynomial function on

$$\underbrace{M_n \oplus \dots \oplus M_n}_{d_1} \oplus \underbrace{M_n \oplus \dots \oplus M_n}_{d_2} \oplus \dots \oplus \underbrace{M_n \oplus \dots \oplus M_n}_{d_m}$$

of multi-degree  $(s_1(1), \dots, s_{d_1}(1), \dots, s_1(m), \dots, s_{d_m}(m))$ . Observe that if  $f$  is an invariant polynomial function on  $M_n^m$ , then each of these multi homogeneous functions is an invariant polynomial function on  $M_n^D$  where  $D = d_1 + \dots + d_m$ .

In particular, we consider the multi-linear function

$$f_{1, \dots, 1} : M_n^D = M_n^{d_1} \oplus \dots \oplus M_n^{d_m} \longrightarrow \mathbb{C}$$

which we call the *polarization* of the polynomial  $f$  and denote with  $Pol(f)$ . Observe that  $Pol(f)$  is symmetric in each of the entries belonging to a block  $M_n^{d_k}$  for every  $1 \leq k \leq m$ . If  $f$  is invariant under  $GL_n$ , then so is the multilinear function  $Pol(f)$  and we know the form of all such functions by the results given before (replacing  $M_n^m$  by  $M_n^D$ ).

Finally, we want to recover  $f$  back from its polarization. We claim to have the equality

$$Pol(f)(\underbrace{A_1, \dots, A_1}_{d_1}, \dots, \underbrace{A_m, \dots, A_m}_{d_m}) = d_1! \dots d_m! f(A_1, \dots, A_m)$$

and hence we recover  $f$ . This process is called *restitution*. The claim follows from the observation that

$$\begin{aligned} f(t_1(1)A_1 + \dots + t_{d_1}(1)A_1, \dots, t_1(m)A_m + \dots + t_{d_m}(m)A_m) &= \\ f((t_1(1) + \dots + t_{d_1}(1))A_1, \dots, (t_1(m) + \dots + t_{d_m}(m))A_m) &= \\ (t_1(1) + \dots + t_{d_1}(1))^{d_1} \dots (t_1(m) + \dots + t_{d_m}(m))^{d_m} f(A_1, \dots, A_m) & \end{aligned}$$

and the definition of  $Pol(f)$ . Hence we have proved that any multi-homogeneous invariant polynomial function  $f$  on  $M_n^m$  of multidegree  $(d_1, \dots, d_m)$  can be obtained by restitution of a multilinear invariant function

$$Pol(f) : M_n^D = M_n^{d_1} \oplus \dots \oplus M_n^{d_m} \longrightarrow \mathbb{C}$$

If we combine this fact with our description of all multilinear invariant functions on  $M_n \oplus \dots \oplus M_n$  we finally obtain :

**THEOREM 1.11** (First fundamental theorem of matrix invariants). *Any polynomial function  $M_n^m \xrightarrow{f} \mathbb{C}$  which is constant along orbits under the action of  $GL_n$  by simultaneous conjugation is a polynomial in the invariants*


$$\text{tr}(X_{i_1} \dots X_{i_l})$$

where  $X_{i_1} \dots X_{i_l}$  run over all possible noncommutative polynomials in the generic matrices  $\{X_1, \dots, X_m\}$ .

We will call the algebra  $\mathbb{C}[M_n^m]$  generated by these invariants the *necklace algebra*  $\mathbb{N}_n^m = \mathbb{C}[M_n^m]^{GL_n}$ . The terminology is justified by the observation that the generators

$$\text{tr}(X_{i_1} X_{i_2} \dots X_{i_l})$$

are only determined up to cyclic permutation of the factors  $X_j$ . They correspond to a *necklace word*  $w$



where each  $i$ -colored bead  $\boxed{i}$  corresponds to a generic matrix  $X_i$ . To obtain an invariant, these bead-matrices are cyclicly multiplied to obtain an  $n \times n$  matrix with coefficients in  $M_n(\mathbb{C}[M_n^m])$ . The trace of this matrix is called  $\text{tr}(w)$  and theorem 1.11 asserts that these elements generate the ring of polynomial invariants.

#### 1.4. The trace algebra.

In this section we will prove a bound on the length of the necklace words  $w$  necessary for the  $\text{tr}(w)$  to generate  $\mathbb{N}_n^m$ . In the last section, after we have determined the relations between these necklaces  $\text{tr}(w)$ , we will be able to improve this bound.

First, we will characterize all  $GL_n$ -equivariant maps from  $M_n^m$  to  $M_n$ , that is all polynomial maps  $M_n^m \xrightarrow{f} M_n$  such that for all  $g \in GL_n$  the diagram below is commutative

$$\begin{array}{ccc} M_n^m & \xrightarrow{f} & M_n \\ \downarrow g.g^{-1} & & \downarrow g.g^{-1} \\ M_n^m & \xrightarrow{f} & M_n \end{array}$$

With pointwise addition and multiplication in the target algebra  $M_n$ , these polynomial maps form a noncommutative algebra  $\mathbb{T}_n^m$  called the *trace algebra*. Obviously,

the trace algebra is a subalgebra of the algebra of *all* polynomial maps from  $M_n^m$  to  $M_n$ , that is,

$$\mathbb{T}_n^m \hookrightarrow M_n(\mathbb{C}[M_n^m])$$

Clearly, using the diagonal embedding of  $\mathbb{C}$  in  $M_n$  any invariant polynomial on  $M_n^m$  determines a  $GL_n$ -equivariant map. Equivalently, using the diagonal embedding of  $\mathbb{C}[M_n^m]$  in  $M_n(\mathbb{C}[M_n^m])$  we can embed the necklace algebra

$$\mathbb{N}_n^m = \mathbb{C}[M_n^m]^{GL_n} \hookrightarrow \mathbb{T}_n^m$$

Another source of  $GL_n$ -equivariant maps are the *coordinate maps*

$$X_i : M_n^m = M_n \oplus \dots \oplus M_n^m \longrightarrow M_n \quad (A_1, \dots, A_m) \mapsto A_i$$

Observe that the coordinate map  $X_i$  is represented by the generic matrix  $\boxed{i} = X_i$  in  $M_n(\mathbb{C}[M_n^m])$ .

**PROPOSITION 1.12.** *As an algebra over the necklace algebra  $\mathbb{N}_n^m$ , the trace algebra  $\mathbb{T}_n^m$  is generated by the elements  $\{X_1, \dots, X_m\}$ .*

**PROOF.** Consider a  $GL_n$ -equivariant map  $M_n^m \xrightarrow{f} M_n$  and associate to it the polynomial map

$$M_n^{m+1} = M_n^m \oplus M_n \xrightarrow{tr(fX_{m+1})} \mathbb{C}$$

defined by sending  $(A_1, \dots, A_m, A_{m+1})$  to  $tr(f(A_1, \dots, A_m) \cdot A_{m+1})$ . For all  $g \in GL_n$  we have that  $f(g \cdot A_1 \cdot g^{-1}, \dots, g \cdot A_m \cdot g^{-1})$  is equal to  $g \cdot f(A_1, \dots, A_m) \cdot g^{-1}$  and hence

$$\begin{aligned} tr(f(g \cdot A_1 \cdot g^{-1}, \dots, g \cdot A_m \cdot g^{-1}) \cdot g \cdot A_{m+1} \cdot g^{-1}) &= tr(g \cdot f(A_1, \dots, A_m) \cdot g^{-1} \cdot g \cdot A_{m+1} \cdot g^{-1}) \\ &= tr(g \cdot f(A_1, \dots, A_m) \cdot A_{m+1} \cdot g^{-1}) \\ &= tr(f(A_1, \dots, A_m) \cdot A_{m+1}) \end{aligned}$$

so  $tr(fX_{m+1})$  is an invariant polynomial function on  $M_n^{m+1}$  which is *linear* in  $X_{m+1}$ . By theorem 1.11 we can therefore write

$$tr(fX_{m+1}) = \sum_{\substack{g_{i_1 \dots i_m} \\ \in \mathbb{N}_n^m}} tr(X_{i_1} \dots X_{i_m} X_{m+1})$$

Here, we used the necklace property allowing to permute cyclicly the trace terms in which  $X_{m+1}$  occurs such that  $X_{m+1}$  occurs as the last factor. But then,  $tr(fX_{m+1}) = tr(gX_{m+1})$  where

$$g = \sum g_{i_1 \dots i_m} X_{i_1} \dots X_{i_m}.$$

Finally, using the *nondegeneracy* of the trace map on  $M_n$  (that is, if  $A, B \in M_n$  such that  $tr(AC) = tr(BC)$  for all  $C \in M_n$ , then  $A = B$ ) it follows that  $f = g$ .  $\square$

If we give each of the generic matrices  $X_i$  degree one, we see that the trace algebra  $\mathbb{T}_n^m$  is a *connected positively graded algebra*

$$\mathbb{T}_n^m = T_0 \oplus T_1 \oplus T_2 \oplus \dots \quad \text{with } T_0 = \mathbb{C}.$$

Our aim is to bound the length of the monomials in the  $X_i$  necessary to generate  $\mathbb{T}_n^m$  as a module over the necklace algebra  $\mathbb{N}_n^m$ . Before we can do this we need to make a small detour in one of the more exotic realms of noncommutative algebra : the *Nagata-Higman problem* .

THEOREM 1.13 (Nagata-Higman). *Let  $R$  be an associative algebra without a unit element. Assume there is a fixed natural number  $n$  such that  $x^n = 0$  for all  $x \in R$ . Then,  $R^{2^n-1} = 0$ , that is*

$$x_1 \cdot x_2 \cdot \dots \cdot x_{2^n-1} = 0$$

for all  $x_j \in R$ .

PROOF. We use induction on  $n$ , the case  $n = 1$  being obvious. Consider for all  $x, y \in R$

$$f(x, y) = yx^{n-1} + xyx^{n-2} + x^2yx^{n-3} + \dots + x^{n-2}yx + x^{n-1}y.$$

Because for all  $c \in \mathbb{C}$  we must have that

$$0 = (y + cx)^n = x^n c^n + f(x, y)c^{n-1} + \dots + y^n$$

it follows that all the coefficients of the  $c^i$  with  $1 \leq i < n$  must be zero, in particular  $f(x, y) = 0$ . But then we have for all  $x, y, z \in R$  that

$$\begin{aligned} 0 &= f(x, z)y^{n-1} + f(x, zy)y^{n-2} + f(x, zy^2)y^{n-3} + \dots + f(x, zy^{n-1}) \\ &= nx^{n-1}zy^{n-1} + zf(y, x^{n-1}) + xzf(y, x^{n-2}) + x^2zf(y, x^{n-3}) + \dots + x^{n-2}zf(y, x) \end{aligned}$$

and therefore  $x^{n-1}zy^{n-1} = 0$ . Let  $I \triangleleft R$  be the twosided ideal of  $R$  generated by all elements  $x^{n-1}$ , then we have that  $I.R.I = 0$ . In the quotient algebra  $\bar{R} = R/I$  every element  $\bar{x}$  satisfies  $\bar{x}^{n-1} = 0$ .

By induction we may assume that  $\bar{R}^{2^{n-1}-1} = 0$ , or equivalently that  $R^{2^{n-1}-1}$  is contained in  $I$ . But then,

$$R^{2^n-1} = R^{2(2^{n-1}-1)+1} = R^{2^{n-1}-1}.R.R^{2^{n-1}-1} \hookrightarrow I.R.I = 0$$

finishing the proof.  $\square$

PROPOSITION 1.14. *The trace algebra  $\mathbb{T}_n^m$  is spanned as a module over the necklace algebra  $\mathbb{N}_n^m$  by all monomials in the generic matrices*

$$X_{i_1} X_{i_2} \dots X_{i_l}$$

of degree  $l \leq 2^n - 1$ .

PROOF. By the diagonal embedding of  $\mathbb{N}_n^m$  in  $M_n(\mathbb{C}[M_n^m])$  it is clear that  $\mathbb{N}_n^m$  commutes with any of the  $X_i$ . Let  $\mathbb{T}_+$  and  $\mathbb{N}_+$  be the strict positive degrees of  $\mathbb{T}_n^m$  and  $\mathbb{N}_n^m$  and form the graded associative algebra (without unit element)

$$R = \mathbb{T}_+ / \mathbb{N}_+ \cdot \mathbb{T}_+$$

Observe that any element  $t \in \mathbb{T}_+$  satisfies an equation of the form

$$t^n + c_1 t^{n-1} + c_2 t^{n-2} + \dots + c_n = 0$$

with all of the  $c_i \in \mathbb{N}_+$ . Indeed we have seen that all the coefficients of the characteristic polynomial of a matrix can be expressed as polynomials in the traces of powers of the matrix. But then, for any  $x \in R$  we have that  $x^n = 0$ .

By the Nagata-Higman theorem we know that  $R^{2^n-1} = (R_1)^{2^n-1} = 0$ . Let  $\mathbb{T}'$  be the graded  $\mathbb{N}_n^m$ -submodule of  $\mathbb{T}_n^m$  spanned by all monomials in the generic matrices  $X_i$  of degree at most  $2^n - 1$ , then the above can be reformulated as

$$\mathbb{T}_n^m = \mathbb{T}' + \mathbb{N}_+ \cdot \mathbb{T}_n^m.$$



We claim that  $\mathbb{T}_n^m = \mathbb{T}'$ . Assume not, then there is a homogeneous  $t \in \mathbb{T}_n^m$  of minimal degree  $d$  not contained in  $\mathbb{T}'$  but still we have a description

$$t = t' + c_1.t_1 + \dots + c_s.t_s$$

with  $t'$  and all  $c_i, t_i$  homogeneous elements. As  $\deg(t_i) < d$ ,  $t_i \in \mathbb{T}'$  for all  $i$  but then is  $t \in \mathbb{T}'$  a contradiction.  $\square$

Finally we are in a position to bound the length of the necklaces generating  $\mathbb{N}_n^m$  as an algebra.

**THEOREM 1.15.** *The necklace algebra  $\mathbb{N}_n^m$  is generated by all necklaces  $tr(w)$  where  $w$  is a necklace word in the bead-matrices  $\{X_1, \dots, X_m\}$  of length  $l \leq 2^n$ .*

**PROOF.** Let  $\mathbb{T}'$  be the  $\mathbb{C}$ -subalgebra of  $\mathbb{T}_n^m$  generated by the generic matrices  $X_i$ . Then,  $tr(\mathbb{T}'_+)$  generates the ideal  $\mathbb{N}_+$ . Let  $\mathbb{S}$  be the set of all monomials in the  $X_i$  of degree at most  $2^n - 1$ . By the foregoing proposition we know that  $\mathbb{T}' \hookrightarrow \mathbb{N}_n^m \cdot \mathbb{S}$ . The trace map

$$tr : \mathbb{T}_n^m \longrightarrow \mathbb{N}_n^m$$

is  $\mathbb{N}_n^m$ -linear and therefore, because  $\mathbb{T}'_+ \subset \mathbb{T}' \cdot (\mathbb{C}X_1 + \dots + \mathbb{C}X_m)$  we have

$$tr(\mathbb{T}'_+) \subset tr(\mathbb{N}_n^m \cdot \mathbb{S} \cdot (\mathbb{C}X_1 + \dots + \mathbb{C}X_m)) \subset \mathbb{N}_n^m \cdot tr(\mathbb{S}')$$

where  $\mathbb{S}'$  is the set of monomials in the  $X_i$  of degree at most  $2^n$ . If  $\mathbb{N}'$  is the  $\mathbb{C}$ -subalgebra of  $\mathbb{N}_n^m$  generated by all  $tr(\mathbb{S}')$ , then we have  $tr(\mathbb{T}'_+) \subset \mathbb{N}_n^m \cdot \mathbb{N}'_+$ . But then, we have

$$\mathbb{N}_+ = \mathbb{N}_n^m tr(\mathbb{T}'_+) \subset \mathbb{N}_n^m \mathbb{N}'_+ \quad \text{and thus} \quad \mathbb{N}_n^m = \mathbb{N}' + \mathbb{N}_n^m \mathbb{N}'_+$$

from which it follows that  $\mathbb{N}_n^m = \mathbb{N}'$  by a similar argument as in the foregoing proof.  $\square$

**EXAMPLE 1.16.** The algebras  $\mathbb{N}_2^2$  and  $\mathbb{T}_2^2$ .

When working with  $2 \times 2$  matrices, the following identities are often helpful

$$0 = A^2 - tr(A)A + det(A)$$

$$A.B + B.A = tr(AB) - tr(A)tr(B) + tr(A)B + tr(B)A$$

for all  $A, B \in M_2$ . Let  $\mathbb{N}'$  be the subalgebra of  $\mathbb{N}_2^2$  generated by  $tr(X_1), tr(X_2), det(X_1), det(X_2)$  and  $tr(X_1X_2)$ . Using the two formulas above and  $\mathbb{N}_2^2$ -linearity of the trace on  $\mathbb{T}_2^2$  we see that the trace of any monomial in  $X_1$  and  $X_2$  of degree  $d \geq 3$  can be expressed in elements of  $\mathbb{N}'$  and traces of monomials of degree  $\leq d - 1$ . Hence, we have

$$\mathbb{N}_2^2 = \mathbb{C}[tr(X_1), tr(X_2), det(X_1), det(X_2), tr(X_1X_2)].$$

Observe that there can be no algebraic relations between these generators as we have seen that the induced map  $\pi : M_2^2 \longrightarrow \mathbb{C}^5$  is surjective. Another consequence of the above identities is that over  $\mathbb{N}_2^2$  any monomial in the  $X_1, X_2$  of degree  $d \geq 3$  can be expressed as a linear combination of  $1, X_1, X_2$  and  $X_1X_2$  and so these elements generate  $\mathbb{T}_2^2$  as a  $\mathbb{N}_2^2$ -module. In fact, they are a basis of  $\mathbb{T}_2^2$  over  $\mathbb{N}_2^2$ . Assume otherwise, there would be a relation say

$$X_1X_2 = \alpha I_2 + \beta X_1 + \gamma X_2$$

with  $\alpha, \beta, \gamma \in \mathbb{C}(tr(X_1), tr(X_2), det(X_1), det(X_2), tr(X_1X_2))$ . Then this relation has to hold for all matrix couples  $(A, B) \in M_2^2$  and we obtain a contradiction if we take the couple

$$A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \quad B = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \quad \text{whence} \quad AB = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}.$$

Concluding, we have the following description of  $\mathbb{N}_2^2$  and  $\mathbb{T}_2^2$  as a subalgebra of  $\mathbb{C}[M_2^2]$  respectively  $M_2(\mathbb{C}[M_2^2])$

$$\begin{cases} \mathbb{N}_2^2 = \mathbb{C}[tr(X_1), tr(X_2), det(X_1), det(X_2), tr(X_1X_2)] \\ \mathbb{T}_2^2 = \mathbb{N}_2^2 \cdot I_2 \oplus \mathbb{N}_2^2 \cdot X_1 \oplus \mathbb{N}_2^2 \cdot X_2 \oplus \mathbb{N}_2^2 \cdot X_1X_2 \end{cases}$$

Observe that we might have taken the generators  $tr(X_i^2)$  rather than  $det(X_i)$  because  $det(X_i) = \frac{1}{2}(tr(X_i)^2 - tr(X_i^2))$  as follows from taking the trace of characteristic polynomial of  $X_i$ .

### 1.5. The symmetric group.

Let  $S_d$  be the symmetric group of all permutations on  $d$  letters. The group algebra  $\mathbb{C} S_d$  is a semisimple algebra. In particular, any simple  $S_d$ -representation is isomorphic to a minimal left ideal of  $\mathbb{C} S_d$  which is generated by an *idempotent*. We will now determine these idempotents.

To start, conjugacy classes in  $S_d$  correspond naturally to *partitions*  $\lambda = (\lambda_1, \dots, \lambda_k)$  of  $d$ , that is, decompositions in natural numbers

$$d = \lambda_1 + \dots + \lambda_k \quad \text{with} \quad \lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_k \geq 1$$

The correspondence associates to a partition  $\lambda = (\lambda_1, \dots, \lambda_k)$  the conjugacy class of a permutation consisting of disjoint cycles of lengths  $\lambda_1, \dots, \lambda_k$ . It is traditional to assign to a partition  $\lambda = (\lambda_1, \dots, \lambda_k)$  a *Young diagram* with  $\lambda_i$  boxes in the  $i$ -th row, the rows of boxes lined up to the left. The *dual partition*  $\lambda^* = (\lambda_1^*, \dots, \lambda_r^*)$  to  $\lambda$  is defined by interchanging rows and columns in the Young diagram of  $\lambda$ . For example, to the partition  $\lambda = (3, 2, 1, 1)$  of 7 we assign the Young diagram

$$\lambda = \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \\ \hline \square & & \\ \hline \square & & \\ \hline \end{array} \quad \lambda^* = \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & & \\ \hline \square & & & \\ \hline \square & & & \\ \hline \end{array}$$

with dual partition  $\lambda^* = (4, 2, 1)$ . A *Young tableau* is a numbering of the boxes of a Young diagram by the integers  $\{1, 2, \dots, d\}$ . For example, two distinct Young tableaux of type  $\lambda$  are

$$\begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline 4 & 5 & \\ \hline 6 & & \\ \hline 7 & & \\ \hline \end{array} \quad \begin{array}{|c|c|c|} \hline 1 & 3 & 5 \\ \hline 2 & 4 & \\ \hline 6 & & \\ \hline 7 & & \\ \hline \end{array}$$

Now, fix a Young tableau  $T$  of type  $\lambda$  and define subgroups of  $S_d$  by

$$P_\lambda = \{ \sigma \in S_d \mid \sigma \text{ preserves each row} \}$$

$$Q_\lambda = \{ \sigma \in S_d \mid \sigma \text{ preserves each column} \}$$

For example, for the second Young tableaux given above we have that

$$\begin{cases} P_\lambda &= S_{\{1,3,5\}} \times S_{\{2,4\}} \times \{(6)\} \times \{(7)\} \\ Q_\lambda &= S_{\{1,2,6,7\}} \times S_{\{3,4\}} \times \{(5)\} \end{cases}$$

Observe that different Young tableaux for the same  $\lambda$  define different subgroups and different elements to be defined below. Still, the simple representations we will construct from them turn out to be isomorphic.

Using these subgroups, we define the following elements in the group algebra  $\mathbb{C}S_d$

$$a_\lambda = \sum_{\sigma \in P_\lambda} e_\sigma \quad , \quad b_\lambda = \sum_{\sigma \in Q_\lambda} \text{sgn}(\sigma) e_\sigma \quad \text{and} \quad c_\lambda = a_\lambda \cdot b_\sigma$$

The element  $c_\lambda$  is called a *Young symmetrizer*. The next result gives an explicit one-to-one correspondence between the simple representations of  $\mathbb{C}S_d$  and the conjugacy classes in  $S_d$  (or, equivalently, Young diagrams).

**THEOREM 1.17.** *For every partition  $\lambda$  of  $d$  the left ideal  $\mathbb{C}S_d \cdot c_\lambda = V_\lambda$  is a simple  $S_d$ -representation and, conversely, any simple  $S_d$ -representation is isomorphic to  $V_\lambda$  for a unique partition  $\lambda$ .*

**PROOF.** (sketch) Observe that  $P_\lambda \cap Q_\lambda = \{e\}$  (any permutation preserving rows as well as columns preserves all boxes) and so any element of  $S_d$  can be written in at most one way as a product  $p \cdot q$  with  $p \in P_\lambda$  and  $q \in Q_\lambda$ . In particular, the Young symmetrizer can be written as  $c_\lambda = \sum \pm e_\sigma$  with  $\sigma = p \cdot q$  for unique  $p$  and  $q$  and the coefficient  $\pm 1 = \text{sgn}(q)$ . From this it follows that for all  $p \in P_\lambda$  and  $q \in Q_\lambda$  we have

$$p \cdot a_\lambda = a_\lambda \cdot p = a_\lambda \quad , \quad \text{sgn}(q)q \cdot b_\lambda = b_\lambda \cdot \text{sgn}(q)q = b_\lambda \quad , \quad p \cdot c_\lambda \cdot \text{sgn}(q)q = c_\lambda$$

Moreover, we claim that  $c_\lambda$  is the unique element in  $\mathbb{C}S_d$  (up to a scalar factor) satisfying the last property. This requires a few preparations.

Assume  $\sigma \notin P_\lambda \cdot Q_\lambda$  and consider the tableaux  $T' = \sigma T$ , that is, replacing the label  $i$  of each box in  $T$  by  $\sigma(i)$ . We claim that there are two distinct numbers which belong to the same row in  $T$  and to the same column in  $T'$ . If this were not the case, then all the distinct numbers in the first row of  $T$  appear in different columns of  $T'$ . But then we can find an element  $q'_1$  in the subgroup  $\sigma \cdot Q_\lambda \cdot \sigma^{-1}$  preserving the columns of  $T'$  to take all these elements to the first row of  $T'$ . But then, there is an element  $p_1 \in P_\lambda$  such that  $p_1 T$  and  $q'_1 T'$  have the same first row. We can proceed to the second row and so on and obtain elements  $p \in P_\lambda$  and  $q' \in \sigma \cdot Q_\lambda \cdot \sigma^{-1}$  such that the tableaux  $pT$  and  $q'T'$  are equal. Hence,  $pT = q'\sigma T$  entailing that  $p = q'\sigma$ . Further,  $q' = \sigma \cdot q \cdot \sigma^{-1}$  but then  $p = q'\sigma = \sigma q$  whence  $\sigma = p \cdot q^{-1} \in P_\lambda \cdot Q_\lambda$ , a contradiction. Therefore, to  $\sigma \notin P_\lambda \cdot Q_\lambda$  we can assign a *transposition*  $\tau = (ij)$  (replacing the two distinct numbers belonging to the same row in  $T$  and to the same column in  $T'$ ) for which  $p = \tau \in P_\lambda$  and  $q = \sigma^{-1} \cdot \tau \cdot \sigma \in Q_\lambda$ .

After these preliminaries, assume that  $c' = \sum a_\sigma e_\sigma$  is an element such that

$$p \cdot c' \cdot \text{sgn}(q)q = c' \quad \text{for all} \quad p \in P_\lambda, q \in Q_\lambda$$

We claim that  $a_\sigma = 0$  whenever  $\sigma \notin P_\lambda \cdot Q_\lambda$ . For take the transposition  $\tau$  found above and  $p = \tau$ ,  $q = \sigma^{-1} \cdot \tau \cdot \sigma$ , then  $p \cdot \sigma \cdot q = \tau \cdot \sigma \cdot \sigma^{-1} \cdot \tau \cdot \sigma = \sigma$ . However, the coefficient of  $\sigma$  in  $c'$  is  $a_\sigma$  and that of  $p \cdot c' \cdot q$  is  $-a_\sigma$  proving the claim. That is,

$$c' = \sum_{p,q} a_{pq} e_{p \cdot q}$$

but then by the property of  $c'$  we must have that  $a_{pq} = \text{sgn}(q)a_e$  whence  $c' = a_e c_\lambda$  finishing the proof of the claimed uniqueness of the element  $c_\lambda$ .

As a consequence we have for all elements  $x \in \mathbb{C}S_d$  that  $c_\lambda.x, c_\lambda = \alpha_x c_\lambda$  for some scalar  $\alpha_x \in \mathbb{C}$  and in particular that  $c_\lambda^2 = n_\lambda c_\lambda$ , for,

$$\begin{aligned} p.(c_\lambda.x.c_\lambda).\text{sgn}(q)q &= p.a_\lambda.b_\lambda.x.a_\lambda.b_\lambda.\text{sgn}(q)q \\ &= a_\lambda.b_\lambda.x.a_\lambda.b_\lambda = c_\lambda.x.c_\lambda \end{aligned}$$

and the statement follows from the uniqueness result for  $c_\lambda$ .

Define  $V_\lambda = \mathbb{C}S_d.c_\lambda$  then we have  $c_\lambda.V_\lambda \subset \mathbb{C}c_\lambda$ . We claim that  $V_\lambda$  is a simple  $S_d$ -representation. Let  $W \subset V_\lambda$  be a simple subrepresentation, then being a left ideal of  $\mathbb{C}S_d$  we can write  $W = \mathbb{C}S_d.x$  with  $x^2 = x$  (note that  $W$  is a direct summand). Assume that  $c_\lambda.W = 0$ , then  $W.W \subset \mathbb{C}S_d.c_\lambda.W = 0$  implying that  $x = 0$  whence  $W = 0$ , a contradiction. Hence,  $c_\lambda.W = \mathbb{C}c_\lambda \subset W$ , but then

$$V_\lambda = \mathbb{C}S_d.c_\lambda \subset W \quad \text{whence } V_\lambda = W$$

is simple. Remains to show that for different partitions, the corresponding simple representations cannot be isomorphic.

We put a *lexicographic* ordering on the partitions by the rule that

$$\lambda > \mu \quad \text{if the first nonvanishing } \lambda_i - \mu_i \text{ is positive}$$

We claim that if  $\lambda > \mu$  then  $a_\lambda.\mathbb{C}S_d.b_\mu = 0$ . It suffices to check that  $a_\lambda.\sigma.b_\mu = 0$  for  $\sigma \in S_d$ . As  $\sigma.b_\mu.\sigma^{-1}$  is the "b-element" constructed from the tableau  $b.T'$  where  $T'$  is the tableaux fixed for  $\mu$ , it is sufficient to check that  $a_\lambda.b_\mu = 0$ . As  $\lambda > \mu$  there are distinct numbers  $i$  and  $j$  belonging to the same row in  $T$  and to the same column in  $T'$ . If not, the distinct numbers in any fixed row of  $T$  must belong to different columns of  $T'$ , but this can only happen for all rows if  $\mu \geq \lambda$ . So consider  $\tau = (ij)$  which belongs to  $P_\lambda$  and to  $Q_\mu$ , whence  $a_\lambda.\tau = a_\lambda$  and  $\tau.b_\mu = -b_\mu$ . But then,

$$a_\lambda.b_\mu = a_\lambda.\tau.\tau.b_\mu = -a_\lambda.b_\mu$$

proving the claim.

If  $\lambda \neq \mu$  we claim that  $V_\lambda$  is not isomorphic to  $V_\mu$ . Assume that  $\lambda > \mu$  and  $\phi$  a  $\mathbb{C}S_d$ -isomorphism with  $\phi(V_\lambda) = V_\mu$ , then

$$\phi(c_\lambda V_\lambda) = c_\lambda \phi(V_\lambda) = c_\lambda V_\mu = c_\lambda \mathbb{C}S_d c_\mu = 0$$

Hence,  $c_\lambda V_\lambda = \mathbb{C}c_\lambda \neq 0$  lies in the kernel of an isomorphism which is clearly absurd.

Summarizing, we have constructed to distinct partitions of  $d$ ,  $\lambda$  and  $\mu$  nonisomorphic simple  $\mathbb{C}S_d$ -representations  $V_\lambda$  and  $V_\mu$ . As we know that there are as many isomorphism classes of simples as there are conjugacy classes in  $S_d$  (or partitions), the  $V_\lambda$  form a complete set of isomorphism classes of simple  $S_d$ -representations, finishing the proof of the theorem.  $\square$


### 1.6. Necklace relations.

In this section we will prove that all the relations holding among the elements of the necklace algebra  $\mathbb{N}_n^m$  are formal consequences of the Cayley-Hamilton equation. First, we will have to set up some notation to clarify what we mean by this.

For technical reasons it is sometimes convenient to have an infinite supply of noncommutative variables  $\{x_1, x_2, \dots, x_i, \dots\}$ . Two monomials of the same degree  $d$  in these variables

$$M = x_{i_1}x_{i_2}\dots x_{i_d} \quad \text{and} \quad M' = x_{j_1}x_{j_2}\dots x_{j_d}$$

are said to be *equivalent* if  $M'$  is obtained from  $M$  by a cyclic permutation, that is, there is a  $k$  such that  $i_1 = j_k$  and all  $i_a = j_b$  with  $b = k + a - 1 \pmod d$ . That is, if they determine the same necklace word



with each of the beads one of the noncommuting variables  $\boxed{i} = x_i$ . To each equivalence class we assign a formal variable that we denote by

$$t(x_{i_1} x_{i_2} \dots x_{i_d}).$$

The *formal necklace algebra*  $\mathbb{N}^\infty$  is then the polynomial algebra on all these (infinitely many) letters. Similarly, we define the *formal trace algebra*  $\mathbb{T}^\infty$  to be the algebra

$$\mathbb{T}^\infty = \mathbb{N}^\infty \otimes_{\mathbb{C}} \mathbb{C}\langle x_1, x_2, \dots, x_i, \dots \rangle$$

that is, the free associative algebra on the noncommuting variables  $x_i$  with coefficients in the polynomial algebra  $\mathbb{N}^\infty$ .

Crucial for our purposes is the existence of an  $\mathbb{N}^\infty$ -linear *formal trace map*

$$t : \mathbb{T}^\infty \longrightarrow \mathbb{N}^\infty$$

defined by the formula

$$t\left(\sum a_{i_1 \dots i_k} x_{i_1} \dots x_{i_k}\right) = \sum a_{i_1 \dots i_k} t(x_{i_1} \dots x_{i_k})$$

where  $a_{i_1 \dots i_k} \in \mathbb{N}^\infty$ .

In an analogous manner we will define infinite versions of the necklace and trace algebras. Let  $M_n^\infty$  be the space of all ordered sequences  $(A_1, A_2, \dots, A_i, \dots)$  with  $A_i \in M_n$  and all but finitely many of the  $A_i$  are the zero matrix. Again,  $GL_n$  acts on  $M_n^\infty$  by simultaneous conjugation and we denote the *infinite necklace algebra*  $\mathbb{N}_n^\infty$  to be the algebra of polynomial functions  $f$

$$M_n^\infty \xrightarrow{f} \mathbb{C}$$

which are constant along orbits. Clearly,  $\mathbb{N}_n^\infty$  is generated as  $\mathbb{C}$ -algebra by the invariants  $tr(M)$  where  $M$  runs over all monomials in the coordinate generic matrices  $X_k = (x_{ij}(k))_{i,j}$  belonging to the  $k$ -th factor of  $M_n^\infty$ . Similarly, the *infinite trace algebra*  $\mathbb{T}_n^\infty$  is the algebra of  $GL_n$ -equivariant polynomial maps

$$M_n^\infty \longrightarrow M_n.$$

Clearly,  $\mathbb{T}_n^\infty$  is the  $\mathbb{C}$ -algebra generated by  $\mathbb{N}_n^\infty$  and the generic matrices  $X_k$  for  $1 \leq k < \infty$ . Observe that  $\mathbb{T}_n^\infty$  is a subalgebra of the matrixring

$$\mathbb{T}_n^\infty \hookrightarrow M_n(\mathbb{C}[M_n^\infty])$$

and as such has a trace map  $tr$  defined on it and from our knowledge of the generators of  $\mathbb{N}_n^\infty$  we know that  $tr(\mathbb{T}_n^\infty) = \mathbb{N}_n^\infty$ .

Now, there are natural algebra epimorphisms

$$\mathbb{T}^\infty \xrightarrow{\tau} \mathbb{T}_n^\infty \quad \text{and} \quad \mathbb{N}^\infty \xrightarrow{\nu} \mathbb{N}_n^\infty$$

defined by  $\tau(t(x_{i_1} \dots x_{i_k})) = \nu(t(x_{i_1} \dots x_{i_k})) = tr(X_{i_1} \dots X_{i_k})$  and  $\tau(x_i) = X_i$ . That is,  $\nu$  and  $\tau$  are compatible with the trace maps

$$\begin{array}{ccc} \mathbb{T}^\infty & \xrightarrow{\tau} & \mathbb{T}_n^\infty \\ \downarrow t & & \downarrow tr \\ \mathbb{N}^\infty & \xrightarrow{\nu} & \mathbb{N}_n^\infty \end{array}$$

We are interested in describing the *necklace relations*, that is, the kernel of  $\nu$ . In the next section we will describe the *trace relations* which is the kernel of  $\tau$ . Note that we obtain the relations holding among the necklaces in  $\mathbb{N}_n^m$  by setting all  $x_i = 0$  with  $i > m$  and all  $t(x_{i_1} \dots x_{i_k}) = 0$  containing a variable with  $i_j > m$ .

In the description a map  $T : \mathbb{C}S_d \longrightarrow \mathbb{N}^\infty$  will be important. Let  $S_d$  be the symmetric group of permutations on  $\{1, \dots, d\}$  and let

$$\sigma = (i_1 i_2 \dots i_\alpha)(j_1 j_2 \dots j_\beta) \dots (z_1 z_2 \dots z_\zeta)$$

be a decomposition of  $\sigma \in S_d$  into cycles including those of length one. The map  $T$  assigns to  $\sigma$  a formal necklace  $T_\sigma(x_1, \dots, x_d)$  defined by

$$T_\sigma(x_1, \dots, x_d) = t(x_{i_1} x_{i_2} \dots x_{i_\alpha}) t(x_{j_1} x_{j_2} \dots x_{j_\beta}) \dots t(x_{z_1} x_{z_2} \dots x_{z_\zeta})$$

Let  $V = V_n$  be again the  $n$ -dimensional vectorspace of column vectors, then  $S_d$  acts naturally on  $V^{\otimes d}$  via

$$\sigma.(v_1 \otimes \dots \otimes v_d) = v_{\sigma(1)} \otimes \dots \otimes v_{\sigma(d)}$$

hence determines a linear map  $\lambda_\sigma \in \text{End}(V^{\otimes d})$ . Recall from section 3 that under the natural identifications

$$(M_n^{\otimes d})^* \simeq (V^{*\otimes d} \otimes V^{\otimes d})^* \simeq \text{End}(V^{\otimes d})$$

the map  $\lambda_\sigma$  defines the multilinear map


$$\mu_\sigma : \underbrace{M_n \otimes \dots \otimes M_n}_d \longrightarrow \mathbb{C}$$

defined by (using the cycle decomposition of  $\sigma$  as before)

$$\mu_\sigma(A_1 \otimes \dots \otimes A_d) = tr(A_{i_1} A_{i_2} \dots A_{i_\alpha}) tr(A_{j_1} A_{j_2} \dots A_{j_\beta}) \dots tr(A_{z_1} A_{z_2} \dots A_{z_\zeta}) \quad .$$

Therefore, a linear combination  $\sum a_\sigma T_\sigma(x_1, \dots, x_d)$  is a necklace relation (that is, belongs to  $\text{Ker } \nu$ ) if and only if the multilinear map  $\sum a_\sigma \mu_\sigma : M_n^{\otimes d} \longrightarrow \mathbb{C}$  is zero. This, in turn, is equivalent to the endomorphism  $\sum a_\sigma \lambda_\sigma \in \text{End}(V^{\otimes d})$ , induced by the action of the element  $\sum a_\sigma e_\sigma \in \mathbb{C}S_d$  on  $V^{\otimes d}$ , being zero. In order to answer the latter problem we have to understand the action of a Young symmetrizer  $c_\lambda \in \mathbb{C}S_d$  on  $V^{\otimes d}$ .

Let  $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$  be a partition of  $d$  and equip the corresponding Young diagram with the standard tableau (that is, order first the boxes in the first row from left to right, then the second row from left to right and so on).



The subgroup  $P_\lambda$  of  $S_d$  which preserves each row then becomes


$$P_\lambda = S_{\lambda_1} \times S_{\lambda_2} \times \dots \times S_{\lambda_k} \hookrightarrow S_d.$$

As  $a_\lambda = \sum_{p \in P_\lambda} e_p$  we see that the image of the action of  $a_\lambda$  on  $V^{\otimes d}$  is the subspace

$$Im(a_\lambda) = Sym^{\lambda_1} V \otimes Sym^{\lambda_2} V \otimes \dots \otimes Sym^{\lambda_k} V \hookrightarrow V^{\otimes d}.$$

Here,  $Sym^i V$  denotes the subspace of symmetric tensors in  $V^{\otimes i}$ .

Similarly, equip the Young diagram of  $\lambda$  with the tableau by ordering first the boxes in the first column from top to bottom, then those of the second column from top to bottom and so on.



Equivalently, give the Young diagram corresponding to the dual partition of  $\lambda$

$$\lambda^* = (\mu_1, \mu_2, \dots, \mu_l)$$

the standard tableau. Then, the subgroup  $Q_\lambda$  of  $S_d$  which preserves each row of  $\lambda$  (or equivalently, each column of  $\lambda^*$ ) is

$$Q_\lambda = S_{\mu_1} \times S_{\mu_2} \times \dots \times S_{\mu_l} \hookrightarrow S_d$$

As  $b_\lambda = \sum_{q \in Q_\lambda} sgn(q)e_q$  we see that the image of  $b_\lambda$  on  $V^{\otimes d}$  is the subspace

$$Im(b_\lambda) = \bigwedge^{\mu_1} V \otimes \bigwedge^{\mu_2} V \otimes \dots \otimes \bigwedge^{\mu_l} V \hookrightarrow V^{\otimes d}.$$

Here,  $\bigwedge^i V$  is the subspace of all anti-symmetric tensors in  $V^{\otimes i}$ . Note that  $\bigwedge^i V = 0$  whenever  $i$  is greater than the dimension  $dim V = n$ . That is, the image of the action of  $b_\lambda$  on  $V^{\otimes d}$  is zero whenever the dual partition  $\lambda^*$  contains a row of length  $\geq n + 1$ , or equivalently, whenever  $\lambda$  has  $\geq n + 1$  rows. Because the Young symmetrizer  $c_\lambda = a_\lambda.b_\lambda \in \mathbb{C} S_d$  we have proved the first result on necklace relations.


**THEOREM 1.18** (Second fundamental theorem of matrix invariants). *A formal necklace*

$$\sum_{\sigma \in S_d} a_\sigma T_\sigma(x_1, \dots, x_d)$$

is a necklace relation (for  $n \times n$  matrices) if and only if the element

$$\sum a_\sigma e_\sigma \in \mathbb{C}S_d$$

belongs to the ideal of  $\mathbb{C}S_d$  spanned by the Young symmetrizers  $c_\lambda$  relative to partitions  $\lambda = (\lambda_1, \dots, \lambda_k)$



with a least  $n + 1$  rows, that is,  $k \geq n + 1$ .

EXAMPLE 1.19. (Fundamental necklace and trace relation.) Consider the partition  $\lambda = (1, 1, \dots, 1)$  of  $n + 1$ , with corresponding Young tableau

1
2
⋮
$n+1$

Then,  $P_\lambda = \{e\}$ ,  $Q_\lambda = S_{n+1}$  and we have the Young symmetrizer

$$a_\lambda = 1 \quad b_\lambda = c_\lambda = \sum_{\sigma \in S_{n+1}} \text{sgn}(\sigma) e_\sigma.$$

The corresponding element is called the *fundamental necklace relation*

$$\text{fund}_n(x_1, \dots, x_{n+1}) = \sum_{\sigma \in S_{n+1}} \text{sgn}(\sigma) T_\sigma(x_1, \dots, x_{n+1}).$$

Clearly,  $\mathfrak{F}(x_1, \dots, x_{n+1})$  is multilinear of degree  $n+1$  in the variables  $\{x_1, \dots, x_{n+1}\}$ . Conversely, any multilinear necklace relation of degree  $n + 1$  must be a scalar multiple of  $\mathfrak{F}(x_1, \dots, x_{n+1})$ . This follows from the proposition as the ideal described there is for  $d = n + 1$  just the scalar multiples of  $\sum_{\sigma \in S_{n+1}} \text{sgn}(\sigma) e_\sigma$ .

Because  $\mathfrak{F}(x_1, \dots, x_{n+1})$  is multilinear in the variables  $x_i$  we can use the cyclic permutation property of the formal trace  $t$  to write it in the form

$$\mathfrak{F}(x_1, \dots, x_{n+1}) = t(\mathfrak{C}\mathfrak{H}(x_1, \dots, x_n) x_{n+1}) \quad \text{with} \quad \mathfrak{C}\mathfrak{H}(x_1, \dots, x_n) \in \mathbb{T}^\infty$$

Observe that  $\mathfrak{C}\mathfrak{H}(x_1, \dots, x_n)$  is multilinear in the variables  $x_i$ . Moreover, by the nondegeneracy of the trace map  $tr$  and the fact that  $\mathfrak{F}(x_1, \dots, x_{n+1})$  is a necklace relation, it follows that  $\mathfrak{C}\mathfrak{H}(x_1, \dots, x_n)$  is a trace relation. Again, any multilinear trace relation of degree  $n$  in the variables  $\{x_1, \dots, x_n\}$  is a scalar multiple of  $\mathfrak{C}\mathfrak{H}(x_1, \dots, x_n)$ . This follows from the corresponding uniqueness result for  $\mathfrak{F}(x_1, \dots, x_{n+1})$ .

We can give an explicit expression of this *fundamental trace relation*

$$\mathfrak{C}\mathfrak{H}(x_1, \dots, x_n) = \sum_{k=0}^n (-1)^k \sum_{i_1 \neq i_2 \neq \dots \neq i_k} x_{i_1} x_{i_2} \dots x_{i_k} \sum_{\sigma \in S_J} \text{sgn}(\sigma) T_\sigma(x_{j_1}, \dots, x_{j_{n-k}})$$



where  $J = \{1, \dots, n\} - \{i_1, \dots, i_k\}$ . In a moment we will see that  $\mathfrak{C}\mathfrak{H}(x_1, \dots, x_n)$  and hence also  $\mathfrak{F}(x_1, \dots, x_{n+1})$  is obtained by polarization of the Cayley-Hamilton identity for  $n \times n$  matrices.

We will explain what we mean by the Cayley-Hamilton polynomial for an element of  $\mathbb{T}^\infty$ . Recall that when  $X \in M_n(A)$  is a matrix with coefficients in a commutative  $\mathbb{C}$ -algebra  $A$  its *characteristic polynomial* is defined to be

$$\chi_X(t) = \det(t\mathbb{1}_n - X) \in A[t]$$

and by the *Cayley-Hamilton theorem* we have the basic relation that  $\chi_X(X) = 0$ . We have seen that the coefficients of the characteristic polynomial can be expressed as polynomial functions in the  $\text{tr}(X^i)$  for  $1 \leq i \leq n$ .

For example if  $n = 2$ , then the characteristic polynomial can be written as

$$\chi_X(t) = t^2 - \text{tr}(X)t + \frac{1}{2}(\text{tr}(X)^2 - \text{tr}(X^2)).$$

For general  $n$  the method for finding these polynomial functions is based on the formal recursive algorithm expressing elementary symmetric functions in term of *Newton functions*, usually expressed by the formulae

$$f(t) = \prod_{i=1}^n (t - \lambda_i),$$

$$\frac{f'(t)}{f(t)} = \frac{d \log f(t)}{dt} = \sum_{i=1}^n \frac{1}{t - \lambda_i} = \sum_{k=0}^{\infty} \frac{1}{t^{k+1}} \left( \sum_{i=1}^n \lambda_i^k \right)$$

Note, if  $\lambda_i$  are the eigenvalues of  $X \in M_n$ , then  $f(t) = \chi_X(t)$  and  $\sum_{i=1}^n \lambda_i^k = \text{tr}(X^k)$ . Therefore, one can use the formulae to express  $f(t)$  in terms of the elements  $\sum_{i=1}^n \lambda_i^k$ . To get the required expression for the characteristic polynomial of  $X$  one only has to substitute  $\sum_{i=1}^n \lambda_i^k$  with  $\text{tr}(X^k)$ .

This allows us to construct a *formal Cayley-Hamilton polynomial*  $\chi_x(x) \in \mathbb{T}^\infty$  of an element  $x \in \mathbb{T}^\infty$  by replacing in the above characteristic polynomial the term  $\text{tr}(X^k)$  with  $t(x^k)$  and  $t^l$  with  $x^l$ . If  $x$  is one of the variables  $x_i$  then  $\chi_x(x)$  is an element of  $\mathbb{T}^\infty$  homogeneous of degree  $n$ . Moreover, by the Cayley-Hamilton theorem it follows immediately that  $\chi_x(x)$  is a trace relation. Hence, if we fully polarize  $\chi_x(x)$  (say, using the variables  $\{x_1, \dots, x_n\}$ ) we obtain a multilinear trace relation of degree  $n$ . By the argument given in the example above we know that this element must be a scalar multiple of  $\mathfrak{C}\mathfrak{H}(x_1, \dots, x_n)$ . In fact, one can see that this scale factor must be  $(-1)^n$  as the leading term of the multilinearization is  $\sum_{\sigma \in S_n} x_{\sigma(1)} \dots x_{\sigma(n)}$  and compare this with the explicit form of  $\mathfrak{C}\mathfrak{H}(x_1, \dots, x_n)$ .

EXAMPLE 1.20. Consider the case  $n = 2$ . The formal Cayley-Hamilton polynomial of an element  $x \in \mathbb{T}^\infty$  is

$$\chi_x(x) = x^2 - t(x)x + \frac{1}{2}(t(x)^2 - t(x^2)) \quad .$$

Polarization with respect to the variables  $x_1$  and  $x_2$  gives the expression

$$x_1 x_2 + x_2 x_1 - t(x_1)x_2 - t(x_2)x_1 + t(x_1)t(x_2) - t(x_1 x_2)$$

which is  $\mathfrak{C}\mathfrak{H}(x_1, x_2)$ . Indeed, multiplying it on the right with  $x_3$  and applying the formal trace  $t$  to it we obtain

$$\begin{aligned} & t(x_1x_2x_3) + t(x_2x_1x_3) - t(x_1)t(x_2x_3) - t(x_2)t(x_1x_3) + t(x_1)t(x_2)t(x_3) - t(x_1x_2)t(x_3) \\ &= T_{(123)}(x_1, x_2, x_3) + T_{(213)}(x_1, x_2, x_3) - T_{(1)(23)}(x_1, x_2, x_3) - T_{(2)(13)}(x_1, x_2, x_3) \\ & \quad + T_{(1)(2)(3)}(x_1, x_2, x_3) - T_{(12)(3)}(x_1, x_2, x_3) \\ &= \sum_{\sigma \in S_3} T_{\sigma}(x_1, x_2, x_3) = \mathfrak{F}(x_1, x_2, x_3) \end{aligned}$$

as required.

**THEOREM 1.21.** *The necklace relations  $\text{Ker } \nu$  is the ideal of  $\mathbb{N}^{\infty}$  generated by all the elements*

$$\mathfrak{F}(m_1, \dots, m_{n+1})$$

where the  $m_i$  run over all monomials in the variables  $\{x_1, x_2, \dots, x_i, \dots\}$ .

**PROOF.** Take a homogeneous necklace relation  $f \in \text{Ker } \nu$  of degree  $d$  and polarize it to get a multilinear element  $f' \in \mathbb{N}^{\infty}$ . Clearly,  $f'$  is also a necklace relation and if we can show that  $f'$  belongs to the described ideal, then so does  $f$  as the process of restitution maps this ideal into itself.

Therefore, we may assume that  $f$  is multilinear of degree  $d$ . A priori  $f$  may depend on more than  $d$  variables  $x_k$ , but we can separate  $f$  as a sum of multilinear polynomials  $f_i$  each depending on precisely  $d$  variables such that for  $i \neq j$   $f_i$  and  $f_j$  do not depend on the same variables. Setting some of the variables equal to zero, we see that each of the  $f_i$  is again a necklace relation.

Thus, we may assume that  $f$  is a multilinear necklace identity of degree  $d$  depending on the variables  $\{x_1, \dots, x_d\}$ . But then we know from theorem 1.18 that we can write

$$f = \sum_{\tau \in S_d} a_{\tau} T_{\tau}(x_1, \dots, x_d)$$

where  $\sum a_{\tau} e_{\tau} \in \mathbb{C}S_d$  belongs to the ideal spanned by the Young symmetrizers of Young diagrams  $\lambda$  having at least  $n+1$  rows.

We claim that this ideal is generated by the Young symmetrizer of the partition  $(1, \dots, 1)$  of  $n+1$  under the natural embedding of  $S_{n+1}$  into  $S_d$ . Let  $\lambda$  be a Young diagram having  $k \geq n+1$  boxes and let  $c_{\lambda}$  be a Young symmetrizer with respect to a tableau where the boxes in the first column are labeled by the numbers  $I = \{i_1, \dots, i_k\}$  and let  $S_I$  be the obvious subgroup of  $S_d$ . As  $Q_{\lambda} = S_I \times Q'$  we see that  $b_{\lambda} = (\sum_{\sigma \in S_I} \text{sgn}(\sigma) e_{\sigma}) \cdot b'$  with  $b' \in \mathbb{C}Q'$ . Hence,  $c_{\lambda}$  belongs to the twosided ideal generated by  $c_I = \sum_{\sigma \in S_I} \text{sgn}(\sigma) e_{\sigma}$  but this is also the twosided ideal generated by  $c_k = \sum_{\sigma \in S_k} \text{sgn}(\sigma) e_{\sigma}$  as one verifies by conjugation with a partition sending  $I$  to  $\{1, \dots, k\}$ . Moreover, by induction one shows that the twosided ideal generated by  $c_k$  belongs to the twosided ideal generated by  $c_d = \sum_{\sigma \in S_d} \text{sgn}(\sigma) e_{\sigma}$ , finishing the proof of the claim.

From this claim, we can write

$$\sum_{\tau \in S_d} a_{\tau} e_{\tau} = \sum_{\tau_i, \tau_j \in S_d} a_{ij} e_{\tau_i} \cdot \left( \sum_{\sigma \in S_{n+1}} \text{sgn}(\sigma) e_{\sigma} \right) \cdot e_{\tau_j}$$

and therefore it suffices to analyze the form of the necklace identity associated to an element of the form

$$e_\tau \cdot \left( \sum_{\sigma \in S_{n+1}} \text{sgn}(\sigma) e_\sigma \right) \cdot e_{\tau'} \quad \text{with } \tau, \tau' \in S_d$$

Now, if a groupelement  $\sum_{\mu \in S_d} b_\mu e_\mu$  corresponds to the formal necklace polynomial  $\mathfrak{G}(x_1, \dots, x_d)$ , then the element  $e_\tau \cdot \left( \sum_{\mu \in S_d} b_\mu e_\mu \right) \cdot e_{\tau^{-1}}$  corresponds to the formal necklace polynomial  $\mathfrak{G}(x_{\tau(1)}, \dots, x_{\tau(d)})$ .

Therefore, we may replace the element  $e_\tau \cdot \left( \sum_{\sigma \in S_{n+1}} \text{sgn}(\sigma) e_\sigma \right) \cdot e_{\tau'}$  by the element

$$\left( \sum_{\sigma \in S_{n+1}} \text{sgn}(\sigma) e_\sigma \right) \cdot e_\eta \quad \text{with } \eta = \tau' \cdot \tau \in S_d$$

We claim that we can write  $\eta = \sigma' \cdot \theta$  with  $\sigma' \in S_{n+1}$  and  $\theta \in S_d$  such that each cycle of  $\theta$  contains at most one of the elements from  $\{1, 2, \dots, n+1\}$ . Indeed assume that  $\eta$  contains a cycle containing more than one element from  $\{1, \dots, n+1\}$ , say 1 and 2, that is

$$\eta = (1i_1i_2 \dots i_r 2j_1j_2 \dots j_s)(k_1 \dots k_\alpha) \dots (z_1 \dots z_\zeta)$$

then we can express the product  $(12) \cdot \eta$  in cycles as

$$(1i_1i_2 \dots i_r)(2j_1j_2 \dots j_s)(k_1 \dots k_\alpha) \dots (z_1 \dots z_\zeta)$$

Continuing in this manner we reduce the number of elements from  $\{1, \dots, n+1\}$  in every cycle to at most one.

But then as  $\sigma' \in S_{n+1}$  we have seen that  $\left( \sum \text{sgn}(\sigma) e_\sigma \right) \cdot e_{\sigma'} = \text{sgn}(\sigma') \left( \sum \text{sgn}(\sigma) e_\sigma \right)$  and consequently

$$\left( \sum_{\sigma \in S_{n+1}} \text{sgn}(\sigma) e_\sigma \right) \cdot e_\eta = \pm \left( \sum_{\sigma \in S_{n+1}} \text{sgn}(\sigma) e_\sigma \right) \cdot e_\theta$$

where each cycle of  $\theta$  contains at most one of  $\{1, \dots, n+1\}$ . Let us write

$$\theta = (1i_1 \dots i_\alpha)(2j_1 \dots j_\beta) \dots (n+1s_1 \dots s_\kappa)(t_1 \dots t_\lambda) \dots (z_1 \dots z_\zeta)$$

Now, let  $\sigma \in S_{n+1}$  then the cycle decomposition of  $\sigma \cdot \theta$  is obtained as follows : substitute in each cycle of  $\sigma$  the element 1 formally by the string  $1i_1 \dots i_\alpha$ , the element 2 by the string  $2j_1 \dots j_\beta$ , and so on until the element  $n+1$  by the string  $n+1s_1 \dots s_\kappa$  and finally adjoin the cycles of  $\theta$  in which no elements from  $\{1, \dots, n+1\}$  appear.

Finally, we can write out the formal necklace element corresponding to the element  $\left( \sum_{\sigma \in S_{n+1}} \text{sgn}(\sigma) e_\sigma \right) \cdot e_\theta$  as

$$\mathfrak{F}(x_1x_{i_1} \dots x_{i_\alpha}, x_2x_{j_1} \dots x_{j_\beta}, \dots, x_{n+1}x_{s_1} \dots x_{s_\kappa})t(x_{t_1} \dots x_{t_\lambda}) \dots t(x_{z_1} \dots x_{z_\zeta})$$

finishing the proof of the theorem.  $\square$

### 1.7. Trace relations.

We will again use the non-degeneracy of the trace map to deduce the trace relations, that is,  $\text{Ker } \tau$  from the description of the necklace relations.

**THEOREM 1.22.** *The trace relations  $\text{Ker } \tau$  is the twosided ideal of the formal trace algebra  $\mathbb{T}^\infty$  generated by all elements*

$$\mathfrak{F}(m_1, \dots, m_{n+1}) \quad \text{and} \quad \mathfrak{C}\mathfrak{H}(m_1, \dots, m_n)$$

where the  $m_i$  run over all monomials in the variables  $\{x_1, x_2, \dots, x_i, \dots\}$ .

PROOF. Consider a trace relation  $\mathfrak{H}(x_1, \dots, x_d) \in \text{Ker } \tau$ . Then, we have a necklace relation of the form

$$t(\mathfrak{H}(x_1, \dots, x_d)x_{d+1}) \in \text{Ker } \nu$$

By theorem 1.21 we know that this element must be of the form

$$\sum n_{i_1 \dots i_{n+1}} \mathfrak{F}(m_{i_1}, \dots, m_{i_{n+1}})$$

where the  $m_i$  are monomials, the  $n_{i_1 \dots i_{n+1}} \in \mathbb{N}^\infty$  and the expression must be linear in the variable  $x_{d+1}$ . That is,  $x_{d+1}$  appears linearly in each of the terms

$$n \mathfrak{F}(m_1, \dots, m_{n+1})$$

so appears linearly in  $n$  or in precisely one of the monomials  $m_i$ . If  $x_{d+1}$  appears linearly in  $n$  we can write

$$n = t(n' \cdot x_{d+1}) \quad \text{where} \quad n' \in \mathbb{T}^\infty.$$

If  $x_{d+1}$  appears linearly in one of the monomials  $m_i$  we may assume that it does so in  $m_{n+1}$ , permuting the monomials if necessary. That is, we may assume  $m_{n+1} = m'_{n+1} \cdot x_{d+1} \cdot m''_{n+1}$  with  $m, m'$  monomials. But then, we can write

$$\begin{aligned} n \mathfrak{F}(m_1, \dots, m_{n+1}) &= nt(\mathfrak{E}\mathfrak{H}(m_1, \dots, m_n) \cdot m'_{n+1} \cdot x_{d+1} \cdot m''_{n+1}) \\ &= t(n \cdot m''_{n+1} \cdot \mathfrak{E}\mathfrak{H}(m_1, \dots, m_n) \cdot m'_{n+1} \cdot x_{d+1}) \end{aligned}$$

using  $\mathbb{N}^\infty$ -linearity and the cyclic permutation property of the formal trace  $t$ . But then, separating the two cases, one can write the total expression

$$\begin{aligned} t(\mathfrak{H}(x_1, \dots, x_d)x_{d+1}) &= t\left[\sum_{\underline{i}} n'_{i_1 \dots i_{n+1}} \mathfrak{F}(m_{i_1}, \dots, m_{i_{n+1}}) \right. \\ &\quad \left. + \sum_{\underline{j}} n_{j_1 \dots j_{n+1}} \cdot m''_{j_{n+1}} \cdot \mathfrak{E}\mathfrak{H}(m_{j_1}, \dots, m_{j_n}) \cdot m'_{j_{n+1}} \right] x_{d+1} \end{aligned}$$

Finally, observe that two formal trace elements  $\mathfrak{H}(x_1, \dots, x_d)$  and  $\mathfrak{K}(x_1, \dots, x_d)$  are equal if the formal necklaces

$$t(\mathfrak{H}(x_1, \dots, x_d)x_{d+1}) = t(\mathfrak{K}(x_1, \dots, x_d)x_{d+1})$$

are equal, finishing the proof.  $\square$

We will give another description of the necklace relations  $\text{Ker } \tau$  which is better suited for the categorical interpretation of  $\mathbb{T}_n^\infty$  to be given in the next chapter. Consider formal trace elements  $m_1, m_2, \dots, m_i, \dots$  with  $m_j \in \mathbb{T}^\infty$ . The *formal substitution*

$$f \mapsto f(m_1, m_2, \dots, m_i, \dots)$$

is the uniquely determined algebra endomorphism of  $\mathbb{T}^\infty$  which maps the variable  $x_i$  to  $m_i$  and is compatible with the formal trace  $t$ . That is, the substitution sends a monomial  $x_{i_1} x_{i_2} \dots x_{i_k}$  to the element  $g_{i_1} g_{i_2} \dots g_{i_k}$  and an element  $t(x_{i_1} x_{i_2} \dots x_{i_k})$  to the element  $t(g_{i_1} g_{i_2} \dots g_{i_k})$ . A *substitution invariant ideal* of  $\mathbb{T}^\infty$  is a twosided ideal of  $\mathbb{T}^\infty$  that is closed under all possible substitutions as well as under the formal trace  $t$ . For any subset of elements  $E \subset \mathbb{T}^\infty$  there is a minimal substitution invariant ideal containing  $E$ . This is the ideal generated by all elements obtained

from  $E$  by making all possible substitutions and taking all their formal traces. We will refer to this ideal as the *substitution invariant ideal generated by  $E$* .

Recall the definition of the formal Cayley-Hamilton polynomial  $\chi_x(x)$  of an element  $x \in \mathbb{T}^\infty$  given in the previous section.

**THEOREM 1.23.** *The trace relations  $\text{Ker } \tau$  is the substitution invariant ideal of  $\mathbb{T}^\infty$  generated by the formal Cayley-Hamilton polynomials*

$$\chi_x(x) \quad \text{for all } x \in \mathbb{T}^\infty$$

**PROOF.** The result follows from theorem 1.22 and the definition of a substitution invariant ideal once we can show that the full polarization of  $\chi_x(x)$ , which we have seen is  $\mathfrak{C}\mathfrak{H}(x_1, \dots, x_n)$ , lies in the substitution invariant ideal generated by the  $\chi_x(x)$ .

This is true since we may replace the process of polarization with the process of multilinearization, whose first step is to replace, for instance

$$\chi_x(x) \quad \text{by} \quad \chi_{x+y}(x+y) - \chi_x(x) - \chi_y(y).$$

The final result of multilinearization is the same as of full polarization and the claim follows as multilinearizing a polynomial in a substitution invariant ideal, we remain in the same ideal.  $\square$

We will use our knowledge on the necklace and trace relations to improve the bound of  $2^n - 1$  in the *Nagata-Higman problem* to  $n^2$ . Recall that this problem asks for a number  $N(n)$  with the property that if  $R$  is an associative  $\mathbb{C}$ -algebra without unit such that  $r^n = 0$  for all  $r \in R$ , then we must have for all  $r_i \in R$  the identity

$$r_1 r_2 \dots r_{N(n)} = 0 \quad \text{in } R.$$

We start by reformulating the problem. Consider the positive part  $\mathbb{F}_+$  of the free  $\mathbb{C}$ -algebra generated by the variables  $\{x_1, x_2, \dots, x_i, \dots\}$

$$\mathbb{F}_+ = \mathbb{C}\langle x_1, x_2, \dots, x_i, \dots \rangle_+$$

which is an associative  $\mathbb{C}$ -algebra without unit. Let  $T(n)$  be the twosided ideal of  $\mathbb{F}_+$  generated by all  $n$ -powers  $f^n$  with  $f \in \mathbb{F}_+$ . Note that the ideal  $T(n)$  is invariant under all substitutions of  $\mathbb{F}_+$ . The Nagata-Higman problem then asks for a number  $N(n)$  such that the product

$$x_1 x_2 \dots x_{N(n)} \in T(n).$$

We will now give an alternative description of the quotient algebra  $\mathbb{F}_+/T(n)$ . Let  $\mathbb{N}_+$  be the positive part of the infinite necklace algebra  $\mathbb{N}_n^\infty$  and  $\mathbb{T}_+$  the positive part of the infinite trace algebra  $\mathbb{T}_n^\infty$ . Consider the quotient associative  $\mathbb{C}$ -algebra without unit

$$\overline{\mathbb{T}_+} = \mathbb{T}_+ / (\mathbb{N}_+ \mathbb{T}_n^\infty).$$

Observe the following facts about  $\overline{\mathbb{T}_+}$ : as a  $\mathbb{C}$ -algebra it is generated by the variables  $X_1, X_2, \dots$  as all the other algebra generators of the form  $t(x_{i_1} \dots x_{i_r})$  of  $\mathbb{T}^\infty$  are mapped to zero in  $\overline{\mathbb{T}_+}$ . Further, from the form of the Cayley-Hamilton polynomial it follows that every  $t \in \overline{\mathbb{T}_+}$  satisfies  $t^n = 0$ . That is, we have an algebra epimorphism

$$\mathbb{F}_+/T(n) \longrightarrow \overline{\mathbb{T}_+}$$

and we claim that it is also injective. To see this, observe that the quotient  $\mathbb{T}^\infty / \mathbb{N}_+^\infty \mathbb{T}^\infty$  is just the free  $\mathbb{C}$ -algebra on the variables  $\{x_1, x_2, \dots\}$ . To obtain  $\overline{\mathbb{T}_+}$  we have to factor out the ideal of trace relations. Now, a formal Cayley-Hamilton


polynomial  $\chi_x(x)$  is mapped to  $x^n$  in  $\mathbb{T}^\infty/\mathbb{N}_+^\infty\mathbb{T}^\infty$ . That is, to obtain  $\overline{\mathbb{T}_+}$  we factor out the substitution invariant ideal (observe that  $t$  is zero here) generated by the elements  $x^n$ , but this is just the definition of  $\mathbb{F}_+/T(n)$ .

Therefore, a reformulation of the Nagata-Higman problem is to find a number  $N = N(n)$  such that the product of the first  $N$  generic matrices

$$X_1 X_2 \dots X_N \in \mathbb{N}_+^\infty \mathbb{T}_n^\infty \quad \text{or, equivalently that} \quad \text{tr}(X_1 X_2 \dots X_N X_{N+1})$$


can be expressed as a linear combination of products of traces of lower degree. Using the description of the necklace relations given in theorem 1.18 we can reformulate this conditions in terms of the group algebra  $\mathbb{C}S_{N+1}$ . Let us introduce the following subspaces of the groupalgebra :

- $A$  will be the subspace spanned by all  $N + 1$  cycles in  $S_{N+1}$ ,
- $B$  will be the subspace spanned by all elements except  $N + 1$  cycles,
- $L(n)$  will be the ideal of  $\mathbb{C}S_{N+1}$  spanned by the Young symmetrizers associated to partitions



with  $\leq n$  rows, and

- $M(n)$  will be the ideal of  $\mathbb{C}S_{N+1}$  spanned by the Young symmetrizers associated to partitions



having more than  $n$  rows.

With these notations, we can reformulate the above condition as

$$(12 \dots NN + 1) \in B + M(n) \quad \text{and consequently} \quad \mathbb{C}S_{N+1} = B + M(n)$$

Define an inner product on the groupalgebra  $\mathbb{C}S_{N+1}$  such that the groupelements form an orthonormal basis, then  $A$  and  $B$  are orthogonal complements and also  $L(n)$  and  $M(n)$  are orthogonal complements. But then, taking orthogonal complements the condition can be rephrased as


$$(B + M(n))^\perp = A \cap L(n) = 0.$$

Finally, let us define an automorphism  $\tau$  on  $\mathbb{C}S_{N+1}$  induced by sending  $e_\sigma$  to  $\text{sgn}(\sigma)e_\sigma$ . Clearly,  $\tau$  is just multiplication by  $(-1)^N$  on  $A$  and therefore the above condition is equivalent to

$$A \cap L(n) \cap \tau L(n) = 0.$$


Moreover, for any Young tableau  $\lambda$  we have that  $\tau(a_\lambda) = b_{\lambda^*}$  and  $\tau(b_\lambda) = a_{\lambda^*}$ . Hence, the automorphism  $\tau$  sends the Young symmetrizer associated to a partition to the Young symmetrizer of the dual partition. This gives the following characterization

- $\tau L(n)$  is the ideal of  $\mathbb{C}S_{N+1}$  spanned by the Young symmetrizers associated to partitions



with  $\leq n$  columns.

Now, specialize to the case  $N = n^2$ . Clearly, any Young diagram having  $n^2 + 1$  boxes must have either more than  $n$  columns or more than  $n$  rows



and consequently we indeed have for  $N = n^2$  that

$$A \cap L(n) \cap \tau L(n) = 0$$

finishing the proof of the promised refinement of the Nagata-Higman bound

**THEOREM 1.24.** *Let  $R$  be an associative  $\mathbb{C}$ -algebra without unit element. Assume that  $r^n = 0$  for all  $r \in R$ . Then, for all  $r_i \in R$  we have*

$$r_1 r_2 \dots r_{n^2} = 0$$

**THEOREM 1.25.** *The necklace algebra  $\mathbb{N}_n^m$  is generated as a  $\mathbb{C}$ -algebra by all elements of the form*

$$\text{tr}(X_{i_1} X_{i_2} \dots X_{i_l})$$

with  $l \leq n^2 + 1$ . The trace algebra  $\mathbb{T}_n^m$  is spanned as a module over the necklace algebra  $\mathbb{N}_n^m$  by all monomials in the generic matrices

$$X_{i_1} X_{i_2} \dots X_{i_l}$$

of degree  $l \leq n^2$ .

### 1.8. Cayley-Hamilton algebras.

In this section we introduce the category  $\mathbf{alg} @_{\mathbf{n}}$  of Cayley-Hamilton algebras of degree  $n$ . A *trace map* on an (affine)  $\mathbb{C}$ -algebra  $A$  is a  $\mathbb{C}$ -linear map

$$\text{tr} : A \longrightarrow A$$

satisfying the following three properties for all  $a, b \in A$  :

- (1)  $\text{tr}(a)b = b\text{tr}(a)$ ,
- (2)  $\text{tr}(ab) = \text{tr}(ba)$  and
- (3)  $\text{tr}(\text{tr}(a)b) = \text{tr}(a)\text{tr}(b)$ .

Note that it follows from the first property that the image  $tr(A)$  of the trace map is contained in the *center* of  $A$ . Consider two algebras  $A$  and  $B$  equipped with a trace map which we will denote by  $tr_A$  respectively  $tr_B$ . A *trace morphism*  $\phi : A \longrightarrow B$  will be a  $\mathbb{C}$ -algebra morphism which is compatible with the trace maps, that is, the following diagram commutes

$$\begin{array}{ccc} A & \xrightarrow{\phi} & B \\ tr_A \downarrow & & \downarrow tr_B \\ A & \xrightarrow{\phi} & B \end{array}$$

This definition turns algebras with a trace map into a category, denoted by  $\mathbf{alg}^t$ . We will say that an algebra  $A$  with trace map  $tr$  is *trace generated* by a subset of elements  $I \subset A$  if the  $\mathbb{C}$ -algebra generated by  $B$  and  $tr(B)$  is equal to  $A$  where  $B$  is the  $\mathbb{C}$ -subalgebra generated by the elements of  $I$ . Note that  $A$  does not have to be generated as a  $\mathbb{C}$ -algebra by the elements from  $I$ .

Observe that for  $\mathbb{T}^\infty$  the formal trace  $t : \mathbb{T}^\infty \longrightarrow \mathbb{N}^\infty \hookrightarrow \mathbb{T}^\infty$  is a trace map. Property (1) follows because  $\mathbb{N}^\infty$  commutes with all elements of  $\mathbb{T}^\infty$ , property (2) is the cyclic permutation property for  $t$  and property (3) is the fact that  $t$  is a  $\mathbb{N}^\infty$ -linear map. The formal trace algebra  $\mathbb{T}^\infty$  is trace generated by the variables  $\{x_1, x_2, \dots, x_i, \dots\}$  but *not* as a  $\mathbb{C}$ -algebra.

Actually,  $\mathbb{T}^\infty$  is the *free algebra* in the generators  $\{x_1, x_2, \dots, x_i, \dots\}$  in the category of algebras with a trace map,  $\mathbf{alg}^t$ . That is, if  $A$  is an algebra with trace  $tr$  which is trace generated by  $\{a_1, a_2, \dots\}$ , then there is a trace preserving algebra epimorphism

$$\mathbb{T}^\infty \xrightarrow{\pi} A \quad .$$

For example, define  $\pi(x_i) = a_i$  and  $\pi(t(x_{i_1} \dots x_{i_l})) = tr(\pi(x_{i_1}) \dots \pi(x_{i_l}))$ . Also, the formal trace algebra  $\mathbb{T}^m$ , that is the subalgebra of  $\mathbb{T}^\infty$  trace generated by  $\{x_1, \dots, x_m\}$ , is the free algebra in the category of algebras with trace that are trace generated by at most  $m$  elements.

Given a trace map  $tr$  on  $A$ , we can define for any  $a \in A$  a *formal Cayley-Hamilton polynomial of degree  $n$* . Indeed, express

$$f(t) = \prod_{i=1}^n (t - \lambda_i)$$

as a polynomial in  $t$  with coefficients polynomial functions in the Newton functions  $\sum_{i=1}^n \lambda_i^k$ . Replacing the Newton function  $\sum \lambda_i^k$  by  $tr(a^k)$  we obtain the Cayley-Hamilton polynomial of degree  $n$

$$\chi_a^{(n)}(t) \in A[t] \quad .$$

DEFINITION 1.26. An (affine)  $\mathbb{C}$ -algebra  $A$  with trace map  $tr : A \longrightarrow A$  is said to be a *Cayley-Hamilton algebra of degree  $n$*  if the following two properties are satisfied :

- (1)  $tr(1) = n$ , and
- (2) For all  $a \in A$  we have  $\chi_a^{(n)}(a) = 0$  in  $A$ .

$\mathbf{alg} @_n$  is the category of Cayley-Hamilton algebras of degree  $n$  with trace preserving morphisms.



Observe that if  $R$  is a commutative  $\mathbb{C}$ -algebra, then  $M_n(R)$  is a Cayley-Hamilton algebra of degree  $n$ . The corresponding trace map is the composition of the usual trace with the inclusion of  $R \longrightarrow M_n(R)$  via scalar matrices. As a consequence, the infinite trace algebra  $\mathbb{T}_n^\infty$  has a trace map induced by the natural inclusion

$$\begin{array}{ccc} \mathbb{T}_n^\infty & \hookrightarrow & M_n(\mathbb{C}[M_n^\infty]) \\ \text{tr} \downarrow \cdots & & \downarrow \text{tr} \\ \mathbb{N}_n^\infty & \hookrightarrow & \mathbb{C}[M_n^\infty] \end{array}$$

which has image  $\text{tr}(\mathbb{T}_n^\infty)$  the infinite necklace algebra  $\mathbb{N}_n^\infty$ . Clearly, being a trace preserving inclusion,  $\mathbb{T}_n^\infty$  is a Cayley-Hamilton algebra of degree  $n$ . With this definition, we have the following categorical description of the trace algebra  $\mathbb{T}_n^\infty$ .

**THEOREM 1.27.** *The trace algebra  $\mathbb{T}_n^\infty$  is the free algebra in the generic matrix generators  $\{X_1, X_2, \dots, X_i, \dots\}$  in the category of Cayley-Hamilton algebras of degree  $n$ .*

*For any  $m$ , the trace algebra  $\mathbb{T}_n^m$  is the free algebra in the generic matrix generators  $\{X_1, \dots, X_m\}$  in the category  $\text{alg } @_{\mathbf{n}}$  of Cayley-Hamilton algebras of degree  $n$  which are trace generated by at most  $m$  elements.*

**PROOF.** Let  $F_n$  be the free algebra in the generators  $\{y_1, y_2, \dots\}$  in the category  $\text{alg } @_{\mathbf{n}}$ , then by freeness of  $\mathbb{T}^\infty$  there is a trace preserving algebra epimorphism

$$\mathbb{T}^\infty \xrightarrow{\pi} F_n \quad \text{with} \quad \pi(x_i) = y_i.$$

By the universal property of  $F_n$ , the ideal  $\text{Ker } \pi$  is the minimal ideal  $I$  of  $\mathbb{T}^\infty$  such that  $\mathbb{T}^\infty/I$  is Cayley-Hamilton of degree  $n$ .

We claim that  $\text{Ker } \pi$  is substitution invariant. Consider a substitution endomorphism  $\phi$  of  $\mathbb{T}^\infty$  and consider the diagram

$$\begin{array}{ccc} \mathbb{T}^\infty & \xrightarrow{\phi} & \mathbb{T}^\infty \\ \text{tr} \downarrow \cdots & \searrow \text{tr} & \downarrow \pi \\ \mathbb{T}^\infty/\text{Ker } \chi & \hookrightarrow & F_n \end{array}$$

then  $\text{Ker } \chi$  is an ideal closed under traces such that  $\mathbb{T}^\infty/\text{Ker } \chi$  is a Cayley-Hamilton algebra of degree  $n$  (being a subalgebra of  $F_n$ ). But then  $\text{Ker } \pi \subset \text{Ker } \chi$  (by minimality of  $\text{Ker } \pi$ ) and therefore  $\chi$  factors over  $F_n$ , that is, the substitution endomorphism  $\phi$  descends to an endomorphism  $\bar{\phi}: F_n \longrightarrow F_n$  meaning that  $\text{Ker } \pi$  is left invariant under  $\phi$ , proving the claim. Further, any formal Cayley-Hamilton polynomial  $\chi_x^{(n)}(x)$  of degree  $n$  of  $x \in \mathbb{T}^\infty$  maps to zero under  $\pi$ . By substitution invariance it follows that the ideal of trace relations  $\text{Ker } \tau \subset \text{Ker } \pi$ . We have seen that  $\mathbb{T}^\infty/\text{Ker } \tau = \mathbb{T}_n^\infty$  is the infinite trace algebra which is a Cayley-Hamilton algebra of degree  $n$ . Thus, by minimality of  $\text{Ker } \pi$  we must have  $\text{Ker } \tau = \text{Ker } \pi$  and hence  $F_n \simeq \mathbb{T}_n^\infty$ . The second assertion follows immediately.  $\square$

Let  $A$  be a Cayley-Hamilton algebra of degree  $n$  which is trace generated by the elements  $\{a_1, \dots, a_m\}$ . We have a trace preserving algebra epimorphism  $p_A$

defined by  $p(X_i) = a_i$

$$\begin{array}{ccc} \mathbb{T}_n^m & \xrightarrow{p_\alpha} & A \\ \text{tr} \downarrow & & \downarrow \text{tr}_A \\ \mathbb{T}_n^m & \xrightarrow{p_\alpha} & A \end{array}$$

and hence a *presentation*  $A \simeq \mathbb{T}_n^m / T_A$  where  $T_A \triangleleft \mathbb{T}_n^m$  is the *ideal of trace relations* holding among the generators  $a_i$ . We recall that  $\mathbb{T}_n^m$  is the ring of  $GL_n$ -equivariant polynomial maps  $M_n^m \xrightarrow{f} M_n$ , that is,

$$M_n(\mathbb{C}[M_n^m])^{GL_n} = \mathbb{T}_n^m$$

where the action of  $GL_n$  is the diagonal action on  $M_n(\mathbb{C}[M_n^m]) = M_n \otimes \mathbb{C}[M_n^m]$ .

Observe that if  $R$  is a commutative algebra, then any two-sided ideal  $I \triangleleft M_n(R)$  is of the form  $M_n(J)$  for an ideal  $J \triangleleft R$ . Indeed, the subsets  $J_{ij}$  of  $(i, j)$  entries of elements of  $I$  is an ideal of  $R$  as can be seen by multiplication with scalar matrices. Moreover, by multiplying on both sides with *permutation matrices* one verifies that  $J_{ij} = J_{kl}$  for all  $i, j, k, l$  proving the claim.

Applying this to the induced ideal  $M_n(\mathbb{C}[M_n^m]) \supseteq T_A \triangleleft M_n(\mathbb{C}[M_n^m])$  we find an ideal  $N_A \triangleleft \mathbb{C}[M_n^m]$  such that

$$M_n(\mathbb{C}[M_n^m]) \supseteq T_A \triangleleft M_n(\mathbb{C}[M_n^m]) = M_n(N_A)$$

Observe that both the induced ideal and  $N_A$  are stable under the respective  $GL_n$ -actions.

Assume that  $V$  and  $W$  are two (not necessarily finite dimensional)  $\mathbb{C}$ -vectorspaces with a locally finite  $GL_n$ -action (that is, every finite dimensional subspace is contained in a finite dimensional  $GL_n$ -stable subspace) and that  $V \xrightarrow{f} W$  is a linear map commuting with the  $GL_n$ -action. In section 2.5 we will see that we can decompose  $V$  and  $W$  uniquely in direct sums of simple representations and in their isotypical components (that is, collecting all factors isomorphic to a given simple  $GL_n$ -representation) and prove that  $V_{(0)} = V^{GL_n}$  respectively  $W_{(0)} = W^{GL_n}$  where  $(0)$  denotes the trivial  $GL_n$ -representation. We obtain a commutative diagram

$$\begin{array}{ccc} V & \xrightarrow{f} & W \\ \downarrow R & & \downarrow R \\ V^{GL_n} & \xrightarrow{f_0} & W^{GL_n} \end{array}$$

where  $R$  is the *Reynolds operator*, that is, the canonical projection to the isotypical component of the trivial representation. Clearly, the Reynolds operator commutes with the  $GL_n$ -action. Moreover, using complete decomposability we see that  $f_0$  is surjective (resp. injective) if  $f$  is surjective (resp. injective).

Because  $N_A$  is a  $GL_n$ -stable ideal of  $\mathbb{C}[M_n^m]$  we can apply the above in the situation

$$\begin{array}{ccc} M_n(\mathbb{C}[M_n^m]) & \xrightarrow{\pi} & M_n(\mathbb{C}[M_n^m]/N_A) \\ \downarrow R & & \downarrow R \\ \mathbb{T}_n^m & \xrightarrow{\pi_0} & M_n(\mathbb{C}[M_n^m]/N_A)^{GL_n} \end{array}$$

and the bottom map factorizes through  $A = \mathbb{T}_n^m/T_A$  giving a surjection

$$A \longrightarrow M_n(\mathbb{C}[M_n^m]/N_A)^{GL_n}.$$

In order to verify that this map is injective (and hence an isomorphism) it suffices to check that

$$M_n(\mathbb{C}[M_n^m]) \cap T_A = M_n(\mathbb{C}[M_n^m]) \cap \mathbb{T}_n^m = T_A.$$

Using the functoriality of the Reynolds operator with respect to multiplication in  $M_n(\mathbb{C}[M_n^\infty])$  with an element  $x \in \mathbb{T}_n^m$  or with respect to the trace map (both commuting with the  $GL_n$ -action) we deduce the following relations :

- For all  $x \in \mathbb{T}_n^m$  and all  $z \in M_n(\mathbb{C}[M_n^\infty])$  we have  $R(xz) = xR(z)$  and  $R(zx) = R(z)x$ .
- For all  $z \in M_n(\mathbb{C}[M_n^\infty])$  we have  $R(\text{tr}(z)) = \text{tr}(R(z))$ .

Assume that  $z = \sum_i t_i x_i n_i \in M_n(\mathbb{C}[M_n^m]) \cap T_A = M_n(\mathbb{C}[M_n^m]) \cap \mathbb{T}_n^m$  with  $m_i, n_i \in M_n(\mathbb{C}[M_n^m])$  and  $t_i \in T_A$ . Now, consider  $X_{m+1} \in \mathbb{T}_n^\infty$ . Using the cyclic property of traces we have

$$\text{tr}(zX_{m+1}) = \sum_i \text{tr}(m_i t_i n_i X_{m+1}) = \sum_i \text{tr}(n_i X_{m+1} m_i t_i)$$

and if we apply the Reynolds operator to it we obtain the equality

$$\text{tr}(zX_{m+1}) = \text{tr}\left(\sum_i R(n_i X_{m+1} m_i) t_i\right)$$

For any  $i$ , the term  $R(n_i X_{m+1} m_i)$  is invariant so belongs to  $\mathbb{T}_n^{m+1}$  and is linear in  $X_{m+1}$ . Knowing the generating elements of  $\mathbb{T}_n^{m+1}$  we can write

$$R(n_i X_{m+1} m_i) = \sum_j s_{ij} X_{m+1} t_{ij} + \sum_k \text{tr}(u_{ik} X_{m+1}) v_{ik}$$

with all of the elements  $s_{ij}, t_{ij}, u_{ik}$  and  $v_{ik}$  in  $\mathbb{T}_n^m$ . Substituting this information and again using the cyclic property of traces we obtain

$$\text{tr}(zX_{m+1}) = \text{tr}\left(\left(\sum_{i,j,k} s_{ij} t_{ij} t_i + \text{tr}(v_{ik} t_i)\right) X_{m+1}\right)$$

and by the nondegeneracy of the trace map we again deduce from this the equality

$$z = \sum_{i,j,k} s_{ij} t_{ij} t_i + \text{tr}(v_{ik} t_i)$$

Because  $t_i \in T_A$  and  $T_A$  is stable under taking traces we deduce from this that  $z \in T_A$  as required.

Because  $A = M_n(\mathbb{C}[M_n^m]/N_A)^{GL_n}$  we can apply functoriality of the Reynolds operator to the setting

$$\begin{array}{ccc} M_n(\mathbb{C}[M_n^m]/N_A) & \xrightleftharpoons{tr} & \mathbb{C}[M_n]/N_A \\ \downarrow R & & \downarrow R \\ A & \xrightleftharpoons{tr_A} & (\mathbb{C}[M_n]/N_A)^{GL_n} \end{array}$$

Concluding we also have the equality

$$tr_A(A) = (\mathbb{C}[M_n^m]/J_A)^{GL_n}.$$

Summarizing, we have proved the following invariant theoretic reconstruction result for Cayley-Hamilton algebras.

**THEOREM 1.28.** *Let  $A$  be a Cayley-Hamilton algebra of degree  $n$ , with trace map  $tr_A$ , which is trace generated by at most  $m$  elements. Then, there is a canonical ideal  $N_A \triangleleft \mathbb{C}[M_n^m]$  from which we can reconstruct the algebras  $A$  and  $tr_A(A)$  as invariant algebras*

$$A = M_n(\mathbb{C}[M_n^m]/N_A)^{GL_n} \quad \text{and} \quad tr_A(A) = (\mathbb{C}[M_n^m]/N_A)^{GL_n}$$

A direct consequence of the above proof is the following *universal property* of the embedding

$$A \xrightarrow{i_A} M_n(\mathbb{C}[M_n^m]/N_A).$$

Let  $R$  be a commutative  $\mathbb{C}$ -algebra, then  $M_n(R)$  with the usual trace is a Cayley-Hamilton algebra of degree  $n$ . If  $f : A \rightarrow M_n(R)$  is a trace preserving morphism, we claim that there exists a natural algebra morphism  $\bar{f} : \mathbb{C}[M_n^m]/N_A \rightarrow R$  such that the diagram

$$\begin{array}{ccc} A & \xrightarrow{f} & M_n(R) \\ \downarrow i_A & \nearrow M_n(\bar{f}) & \\ M_n(\mathbb{C}[M_n^m]/N_A) & & \end{array}$$

where  $M_n(\bar{f})$  is the algebra morphism defined entrywise. To see this, consider the composed trace preserving morphism  $\phi : \mathbb{T}_n^m \rightarrow A \xrightarrow{f} M_n(R)$ . Its image is fully determined by the images of the trace generators  $X_k$  of  $\mathbb{T}_n^m$  which are say  $m_k = (m_{ij}(k))_{i,j}$ . But then we have an algebra morphism  $\mathbb{C}[M_n^m] \xrightarrow{g} R$  defined by sending the variable  $x_{ij}(k)$  to  $m_{ij}(k)$ . Clearly,  $T_A \subset \text{Ker } \phi$  and after inducing to  $M_n(\mathbb{C}[M_n^m])$  it follows that  $N_A \subset \text{Ker } g$  proving that  $g$  factors through  $\mathbb{C}[M_n^m]/J_A \rightarrow R$ . This morphism has the required universal property.

### References.

The first fundamental theorem of matrix invariants, theorem 1.11, is due independently to G. B. Gurevich [26], C. Procesi [64] and K. S. Siberskii [75]. The second fundamental theorem of matrix invariants, theorem 1.18 is due independently to C. Procesi [64] and Y. P. Razmyslov [66]. Our treatment follows the paper [64] of C. Procesi, supplemented with material taken from the lecture notes

of H-P. Kraft [46] and E. Formanek [22]. The invariant theoretic reconstruction result, theorem 1.28, is due to C. Procesi [65].



## Reconstructing Algebras.

### 2.1. Representation spaces.

For a noncommutative affine algebra  $A$  with generating set  $\{a_1, \dots, a_m\}$ , there is an epimorphism

$$\mathbb{C}\langle x_1, \dots, x_m \rangle \xrightarrow{\phi} A$$

defined by  $\phi(x_i) = a_i$ . That is, a presentation of  $A$  as

$$A \simeq \mathbb{C}\langle x_1, \dots, x_m \rangle / I_A$$

where  $I_A$  is the *twosided ideal of relations* holding among the  $a_i$ . For example, if  $A = \mathbb{C}[x_1, \dots, x_m]$ , then  $I_A$  is the twosided ideal of  $\mathbb{C}\langle x_1, \dots, x_m \rangle$  generated by the elements  $x_i x_j - x_j x_i$  for all  $1 \leq i, j \leq m$ .

An  $n$ -dimensional representation of  $A$  is an algebra morphism

$$A \xrightarrow{\psi} M_n$$

from  $A$  to the algebra of  $n \times n$  matrices over  $\mathbb{C}$ . If  $A$  is generated by  $\{a_1, \dots, a_m\}$ , then  $\psi$  is fully determined by the point

$$(\psi(a_1), \psi(a_2), \dots, \psi(a_m)) \in M_n^m = \underbrace{M_n \oplus \dots \oplus M_n}_m.$$

We claim that  $\text{rep}_n(A)$ , the set of all  $n$ -dimensional representations of  $A$ , forms a Zariski closed subset of  $M_n^m$ . To begin, observe that

$$\text{rep}_n(\mathbb{C}\langle x_1, \dots, x_m \rangle) = M_n^m$$

as any  $m$ -tuple of  $n \times n$  matrices  $(A_1, \dots, A_m) \in M_n^m$  determines an algebra morphism  $\mathbb{C}\langle x_1, \dots, x_m \rangle \xrightarrow{\psi} M_n$  by taking  $\psi(x_i) = A_i$ .

Given a presentation  $A = \mathbb{C}\langle x_1, \dots, x_m \rangle / I_A$  an  $m$ -tuple  $(A_1, \dots, A_m) \in M_n^m$  determines an  $n$ -dimensional representation of  $A$  if (and only if) for every noncommutative polynomial  $r(x_1, \dots, x_m) \in I_A \triangleleft \mathbb{C}\langle x_1, \dots, x_m \rangle$  we have that

$$r(A_1, \dots, A_m) = \begin{bmatrix} 0 & \dots & 0 \\ \vdots & & \vdots \\ 0 & \dots & 0 \end{bmatrix} \in M_n.$$

Hence, consider the ideal  $I_A(n)$  of  $\mathbb{C}[M_n^m] = \mathbb{C}[x_{ij}(k) \mid 1 \leq i, j \leq n, 1 \leq k \leq m]$  generated by all the entries of the matrices in  $M_n(\mathbb{C}[M_n^m])$  of the form

$$r(X_1, \dots, X_m) \quad \text{for all } r(x_1, \dots, x_m) \in I_A.$$

We see that the *reduced representation variety*  $\text{rep}_n A$  is the set of simultaneous zeroes of the ideal  $I_A(n)$ , that is,

$$\text{rep}_n A = \mathbb{V}(I_A(n)) \hookrightarrow M_n^m$$

proving the claim, where  $\mathbb{V}$  denotes the *closed* set in the *Zariski topology* Zariski topology determined by an ideal, the complement of which we will denote with  $\mathbb{X}$ ). Observe that even when  $A$  is not finitely presented, the ideal  $I_A(n)$  is finitely generated as an ideal of the commutative polynomial algebra  $\mathbb{C}[M_n^m]$ .

Often, the ideal  $I_A(n)$  contains more information than the closed subset  $\text{rep}_n(A) = \mathbb{V}(I_A(n))$  which only determines the radical ideal of  $I_A(n)$ . This forces us to consider also the *representation variety* (or scheme)  $\underline{\text{rep}}_n A$ .

EXAMPLE 2.1. It may happen that  $\text{rep}_n A = \emptyset$ . For example, consider the *Weyl algebra*

$$A_1(\mathbb{C}) = \mathbb{C}\langle x, y \rangle / (xy - yx - 1)$$

If a couple of  $n \times n$ -matrices  $(A, B) \in \text{rep}_n A_1(\mathbb{C})$  then we must have

$$A.B - B.A = \mathbb{1}_n \in M_n$$

However, taking traces on both sides gives a contradiction as  $\text{tr}(AB) = \text{tr}(BA)$  and  $\text{tr}(\mathbb{1}_n) = n \neq 0$ .

In the foregoing chapter we studied the action of  $GL_n$  by simultaneous conjugation on  $M_n^m$ . We claim that  $\text{rep}_n A \hookrightarrow M_n^m$  is *stable* under this action, that is, if  $(A_1, \dots, A_m) \in \text{rep}_n A$ , then also  $(gA_1g^{-1}, \dots, gA_mg^{-1}) \in \text{rep}_n A$ . This is clear by composing the  $n$ -dimensional representation  $\psi$  of  $A$  determined by  $(A_1, \dots, A_m)$  with the algebra automorphism of  $M_n$  given by conjugation with  $g \in GL_n$ ,

$$\begin{array}{ccc} A & \xrightarrow{\psi} & M_n \\ & \searrow^{g \cdot \psi} & \downarrow^{g \cdot g^{-1}} \\ & & M_n \end{array}$$

That is,  $\text{rep}_n A$  is a  $GL_n$ -variety. We will give an interpretation of the orbits under this action.

Recall that a *left  $A$ -module  $M$*  is a vectorspace on which elements of  $A$  act on the left as linear operators satisfying the conditions

$$1.m = m \quad \text{and} \quad a.(b.m) = (ab).m$$

for all  $a, b \in A$  and all  $m \in M$ . An  $A$ -module *morphism*  $M \xrightarrow{f} N$  between two left  $A$ -modules is a linear map such that  $f(a.m) = a.f(m)$  for all  $a \in A$  and all  $m \in M$ . An  $A$ -module *automorphism* is an  $A$ -module morphism  $M \xrightarrow{f} N$  such that there is an  $A$ -module morphism  $N \xrightarrow{g} M$  such that  $f \circ g = \text{id}_M$  and  $g \circ f = \text{id}_N$ .

Assume the  $A$ -module  $M$  has complex dimension  $n$ , then after fixing a basis we can identify  $M$  with  $\mathbb{C}^n$  (column vectors). For any  $a \in A$  we can represent the linear action of  $a$  on  $M$  by an  $n \times n$  matrix  $\psi(a) \in M_n$ . The condition that  $a.(b.m) = (ab).m$  for all  $m \in M$  asserts that  $\psi(ab) = \psi(a)\psi(b)$  for all  $a, b \in A$ , that is,  $\psi$  is an algebra morphism  $A \xrightarrow{\psi} M_n$  and hence  $M$  determines an  $n$ -dimensional representation of  $A$ . Conversely, an  $n$ -dimensional representation  $A \xrightarrow{\psi} M_n$  determines an  $A$ -module structure on  $\mathbb{C}^n$  by the rule

$$a.v = \psi(a)v \quad \text{for all} \quad v \in \mathbb{C}^n.$$



Hence, there is a one-to-one correspondence between the  $n$ -dimensional representations of  $A$  and the  $A$ -module structures on  $\mathbb{C}^n$ . We call the reduced variety  $\text{rep}_n A$  the *reduced representation variety* of  $A$ . If two  $n$ -dimensional  $A$ -module structures  $M$  and  $N$  on  $\mathbb{C}^n$  are isomorphic (determined by a linear invertible map  $g \in GL_n$ ) then for all  $a \in A$  we have the commutative diagram

$$\begin{array}{ccc} M & \xrightarrow{g} & N \\ \downarrow a. & & \downarrow a. \\ M & \xrightarrow{g} & N \end{array}$$

Hence, if the action of  $a$  on  $M$  is represented by the matrix  $A$ , then the action of  $a$  on  $N$  is represented by the matrix  $g.A.g^{-1}$ . Therefore, two  $A$ -module structures on  $\mathbb{C}^n$  are isomorphic if and only if the points of  $\text{rep}_n A$  corresponding to them lie in the same  $GL_n$ -orbit. Concluding, studying  $n$ -dimensional  $A$ -modules up to isomorphism is the same as studying the  $GL_n$ -orbits in the reduced representation variety  $\text{rep}_n A$ .

If the defining ideal  $I_A(n)$  is a radical ideal, the above suffices. In general, the *scheme* structure of the representation variety  $\underline{\text{rep}}_n A$  will be important. By definition, the scheme  $\underline{\text{rep}}_n A$  is the functor assigning to any (affine) commutative  $\mathbb{C}$ -algebra  $R$ , the set

$$\underline{\text{rep}}_n A(R) = \text{Alg}_{\mathbb{C}}(\mathbb{C}[M_n^m]/I_A(n), R)$$

of  $\mathbb{C}$ -algebra morphisms  $\frac{\mathbb{C}[M_n^m]}{I_A(n)} \xrightarrow{\psi} R$ . Such a map  $\psi$  is determined by the image  $\psi(x_{ij}(k)) = r_{ij}(k) \in R$ . That is,  $\psi \in \underline{\text{rep}}_n A(R)$  determines an  $m$ -tuple of  $n \times n$  matrices with coefficients in  $R$

$$(r_1, \dots, r_m) \in \underbrace{M_n(R) \oplus \dots \oplus M_n(R)}_m \quad \text{where} \quad r_k = \begin{bmatrix} r_{11}(k) & \dots & r_{1n}(k) \\ \vdots & & \vdots \\ r_{n1}(k) & \dots & r_{nn}(k) \end{bmatrix}.$$

Clearly, for any  $r(x_1, \dots, x_m) \in I_A$  we must have that  $r(r_1, \dots, r_m)$  is the zero matrix in  $M_n(R)$ . That is,  $\psi$  determines uniquely an  $R$ -algebra morphism

$$\psi : R \otimes_{\mathbb{C}} A \longrightarrow M_n(R) \quad \text{by mapping} \quad x_k \mapsto r_k.$$

Alternatively, we can identify the set  $\underline{\text{rep}}_n A(R)$  with the set of left  $R \otimes A$ -module structures on the free  $R$ -module  $R^{\oplus n}$  of rank  $n$ .

## 2.2. Some algebraic geometry.

Throughout this book we assume that the reader has some familiarity with algebraic geometry, as contained in the first two chapters of the textbook [28]. In this section we restrict to the dimension formulas and the relation between Zariski and analytic closure, illustrating them with examples from representation varieties. We will work only with reduced varieties in this section.

A morphism  $X \xrightarrow{\phi} Y$  between two affine *irreducible varieties* (that is, the coordinate rings  $\mathbb{C}[X]$  and  $\mathbb{C}[Y]$  are domains) is said to be *dominant* if the image  $\phi(X)$  is Zariski dense in  $Y$ . On the level of the coordinate algebras dominance is

equivalent to  $\phi^* : \mathbb{C}[Y] \longrightarrow \mathbb{C}[X]$  being injective and hence inducing a field extension  $\phi^* : \mathbb{C}(Y) \hookrightarrow \mathbb{C}(X)$  between the function fields. Indeed, for  $f \in \mathbb{C}[Y]$  the function  $\phi^*(f)$  is by definition the composition

$$X \xrightarrow{\phi} Y \xrightarrow{f} \mathbb{C}$$

and therefore  $\phi^*(f) = 0$  iff  $f(\phi(X)) = 0$  iff  $f(\overline{\phi(X)}) = 0$ .

A morphism  $X \xrightarrow{\phi} Y$  between two affine varieties is said to be *finite* if under the algebra morphism  $\phi^*$  the coordinate algebra  $\mathbb{C}[X]$  is a finite  $\mathbb{C}[Y]$ -module. An important property of finite morphisms is that they are *closed*, that is the image of a closed subset is closed. Indeed, we can replace without loss of generality  $Y$  by the closed subset  $\overline{\phi(X)} = \mathbb{V}_Y(\text{Ker } \phi^*)$  and hence assume that  $\phi^*$  is an inclusion  $\mathbb{C}[Y] \hookrightarrow \mathbb{C}[X]$ . The claim then follows from the fact that in a finite extension there exists for any maximal ideal  $N \triangleleft \mathbb{C}[Y]$  a maximal ideal  $M \triangleleft \mathbb{C}[X]$  such that  $M \cap \mathbb{C}[Y] = N$ .

EXAMPLE 2.2. Let  $X$  be an irreducible affine variety of dimension  $d$ . By the *Noether normalization* result  $\mathbb{C}[X]$  is a finite module over a polynomial subalgebra  $\mathbb{C}[f_1, \dots, f_d]$ . But then, the finite inclusion  $\mathbb{C}[f_1, \dots, f_d] \hookrightarrow \mathbb{C}[X]$  determines a finite surjective morphism

$$X \xrightarrow{\phi} \mathbb{C}^d$$

An important source of finite morphisms is given by integral extensions. Recall that, if  $R \hookrightarrow S$  is an inclusion of domains we call  $S$  *integral* over  $R$  if every  $s \in S$  satisfies an equation

$$s^n = \sum_{i=0}^{n-1} r_i s^i \quad \text{with } r_i \in R.$$

A *normal* domain  $R$  has the property that any element of its field of fractions which is integral over  $R$  belongs already to  $R$ . If  $X \xrightarrow{\phi} Y$  is a dominant morphism between two irreducible affine varieties, then  $\phi$  is finite if and only if  $\mathbb{C}[X]$  is integral over  $\mathbb{C}[Y]$  for the embedding coming from  $\phi^*$ .

PROPOSITION 2.3. *Let  $X \xrightarrow{\phi} Y$  be a dominant morphism between irreducible affine varieties. Then, for any  $x \in X$  and any irreducible component  $C$  of the fiber  $\phi^{-1}(\phi(x))$  we have*

$$\dim C \geq \dim X - \dim Y.$$

Moreover, there is a nonempty open subset  $U$  of  $Y$  contained in the image  $\phi(X)$  such that for all  $u \in U$  we have

$$\dim \phi^{-1}(u) = \dim X - \dim Y.$$

PROOF. Let  $d = \dim X - \dim Y$  and apply the Noether normalization result to the affine  $\mathbb{C}(Y)$ -algebra  $\mathbb{C}(Y)\mathbb{C}[X]$ . Then, we can find a function  $g \in \mathbb{C}[Y]$  and algebraic independent functions  $f_1, \dots, f_d \in \mathbb{C}[X]_g$  ( $g$  clears away any denominators that occur after applying the normalization result) such that  $\mathbb{C}[X]_g$  is *integral* over

$\mathbb{C}[Y]_g[f_1, \dots, f_d]$ . That is, we have the commutative diagram

$$\begin{array}{ccc} \mathbb{X}_X(g) & \xrightarrow{\rho} & \mathbb{X}_Y(g) \times \mathbb{C}^d \\ \downarrow & & \downarrow pr_1 \\ X & \xrightarrow{\phi} & Y \longleftarrow \supset \mathbb{X}_Y(g) \end{array}$$

where we know that  $\rho$  is finite and surjective. But then we have that the open subset  $\mathbb{X}_Y(g)$  lies in the image of  $\phi$  and in  $\mathbb{X}_Y(g)$  all fibers of  $\phi$  have dimension  $d$ . For the first part of the statement we have to recall the statement of *Krull's Hauptideal result*: if  $X$  is an irreducible affine variety and  $g_1, \dots, g_r \in \mathbb{C}[X]$  with  $(g_1, \dots, g_r) \neq \mathbb{C}[X]$ , then any component  $C$  of  $\mathbb{V}_X(g_1, \dots, g_r)$  satisfies the inequality

$$\dim C \geq \dim X - r.$$

If  $\dim Y = r$  apply this result to the  $g_i$  determining the morphism

$$X \xrightarrow{\phi} Y \longrightarrow \mathbb{C}^r$$

where the latter morphism is the one from example 2.2.  $\square$

In fact, a stronger result holds. *Chevalley's theorem* asserts the following.

**THEOREM 2.4.** *Let  $X \xrightarrow{\phi} Y$  be a morphism between affine varieties, the function*

$$X \longrightarrow \mathbb{N} \quad \text{defined by} \quad x \mapsto \dim_x \phi^{-1}(\phi(x))$$

*is upper-semicontinuous. That is, for all  $n \in \mathbb{N}$ , the set*

$$\{x \in X \mid \dim_x \phi^{-1}(\phi(x)) \leq n\}$$

*is Zariski open in  $X$ .*

**PROOF.** Let  $Z(\phi, n)$  be the set  $\{x \in X \mid \dim_x \phi^{-1}(\phi(x)) \geq n\}$ . We will prove that  $Z(\phi, n)$  is closed by induction on the dimension of  $X$ . We first make some reductions. We may assume that  $X$  is irreducible. For, let  $X = \cup_i X_i$  be the decomposition of  $X$  into irreducible components, then  $Z(\phi, n) = \cup Z(\phi \mid X_i, n)$ . Next, we may assume that  $Y = \overline{\phi(X)}$  whence  $Y$  is also irreducible and  $\phi$  is a dominant map. Now, we are in the setting of proposition 2.3. Therefore, if  $n \leq \dim X - \dim Y$  we have  $Z(\phi, n) = X$  by that proposition, so it is closed. If  $n > \dim X - \dim Y$  consider the open set  $U$  in  $Y$  of proposition 2.3. Then,  $Z(\phi, n) = Z(\phi \mid (X - \phi^{-1}(U)), n)$ . the dimension of the closed subvariety  $X - \phi^{-1}(U)$  is strictly smaller than  $\dim X$  hence by induction we may assume that  $Z(\phi \mid (X - \phi^{-1}(U)), n)$  is closed in  $X - \phi^{-1}(U)$  whence closed in  $X$ .  $\square$

An immediate consequence of the foregoing proposition is that for any morphism  $X \xrightarrow{\phi} Y$  between affine varieties, the image  $\phi(X)$  contains an open dense subset of  $\overline{\phi(X)}$  (reduce to irreducible components and apply the proposition).

**EXAMPLE 2.5.** Let  $A$  be an affine  $\mathbb{C}$ -algebra and  $M \in \text{rep}_n A$ . We claim that the orbit

$$\mathcal{O}(M) = GL_n.M \quad \text{is Zariski open in its closure} \quad \overline{\mathcal{O}(M)}.$$

Consider the 'orbit-map'  $GL_n \xrightarrow{\phi} \text{rep}_n A$  defined by  $g \mapsto g.M$ . then, by the above remark  $\overline{\mathcal{O}(M)} = \overline{\phi(GL_n)}$  contains a Zariski open subset  $U$  of  $\overline{\mathcal{O}(M)}$  contained in the image of  $\phi$  which is  $\mathcal{O}(M)$ . But then,

$$\mathcal{O}(M) = GL_n.M = \cup_{g \in GL_n} g.U$$

is also open in  $\overline{\mathcal{O}(M)}$ . Next, we claim that  $\overline{\mathcal{O}(M)}$  contains a closed orbit. Indeed, assume  $\mathcal{O}(M)$  is not closed, then the complement  $C_M = \overline{\mathcal{O}(M)} - \mathcal{O}(M)$  is a proper Zariski closed subset whence  $\dim C < \dim \overline{\mathcal{O}(M)}$ . But,  $C$  is the union of  $GL_n$ -orbits  $\mathcal{O}(M_i)$  with  $\dim \mathcal{O}(M_i) < \dim \overline{\mathcal{O}(M)}$ . Repeating the argument with the  $M_i$  and induction on the dimension we will obtain a closed orbit in  $\overline{\mathcal{O}(M)}$ .

Next, we want to relate the Zariski closure with the  $\mathbb{C}$ -closure (that is, the usual complex or analytic topology). Whereas they are usually not equal (for example, the unit circle in  $\mathbb{C}^1$ ), we will show that they coincide for the important class of *constructible* subsets. A subset  $Z$  of an affine variety  $X$  is said to be *locally closed* if  $Z$  is open in its Zariski closure  $\overline{Z}$ . A subset  $Z$  is said to be *constructible* if  $Z$  is the union of finitely many locally closed subsets. Clearly, finite unions, finite intersections and complements of constructible subsets are again constructible. The importance of constructible sets for algebraic geometry is clear from the following result.

**PROPOSITION 2.6.** *Let  $X \xrightarrow{\phi} Y$  be a morphism between affine varieties. If  $Z$  is a constructible subset of  $X$ , then  $\phi(Z)$  is a constructible subset of  $Y$ .*

**PROOF.** Because every open subset of  $X$  is a finite union of special open sets which are themselves affine varieties, it suffices to show that  $\phi(X)$  is constructible. We will use induction on  $\dim \overline{\phi(X)}$ . There exists an open subset  $U \subset \overline{\phi(X)}$  which is contained in  $\phi(X)$ . Consider the closed complement  $W = \overline{\phi(X)} - U$  and its inverse image  $X' = \phi^{-1}(W)$ . Then,  $X'$  is an affine variety and by induction we may assume that  $\phi(X')$  is constructible. But then,  $\phi(X) = U \cup \phi(X')$  is also constructible.  $\square$

**EXAMPLE 2.7.** Let  $A$  be an affine  $\mathbb{C}$ -algebra. The subset  $\text{ind}_n A \hookrightarrow \text{rep}_n A$  of the *indecomposable*  $n$ -dimensional  $A$ -modules is constructible. Indeed, define for any pair  $k, l$  such that  $k + l = n$  the morphism

$$GL_n \times \text{rep}_k A \times \text{rep}_l A \longrightarrow \text{rep}_n A$$

by sending a triple  $(g, M, N)$  to  $g.(M \oplus N)$ . By the foregoing result the image of this map is constructible. The decomposable  $n$ -dimensional  $A$ -modules belong to one of these finitely many sets whence are constructible, but then so is its complement which in  $\text{ind}_n A$ .

Apart from being closed, finite morphisms often satisfy the *going-down property*. That is, consider a finite and surjective morphism

$$X \xrightarrow{\phi} Y$$

where  $X$  is irreducible and  $Y$  is normal (that is,  $\mathbb{C}[Y]$  is a normal domain). Let  $Y' \hookrightarrow Y$  an irreducible Zariski closed subvariety and  $x \in X$  with image  $\phi(x) = y' \in Y'$ . Then, the going-down property asserts the existence of an irreducible Zariski closed subvariety  $X' \hookrightarrow X$  such that  $x \in X'$  and  $\phi(X') = Y'$ . In particular, the morphism  $X' \xrightarrow{\phi} Y'$  is again finite and surjective and in particular  $\dim X' = \dim Y'$ .

LEMMA 2.8. *Let  $x \in X$  an irreducible affine variety and  $U$  a Zariski open subset. Then, there is an irreducible curve  $C \hookrightarrow X$  through  $x$  and intersecting  $U$ .*

PROOF. If  $d = \dim X$  consider the finite surjective morphism  $X \xrightarrow{\phi} \mathbb{C}^d$  of example 2.2. Let  $y \in \mathbb{C}^d - \phi(X - U)$  and consider the line  $L$  through  $y$  and  $\phi(x)$ . By the going-down property there is an irreducible curve  $C \hookrightarrow X$  containing  $x$  such that  $\phi(C) = L$  and by construction  $C \cap U \neq \emptyset$ .  $\square$

PROPOSITION 2.9. *Let  $X \xrightarrow{\phi} Y$  be a dominant morphism between irreducible affine varieties any  $y \in Y$ . Then, there is an irreducible curve  $C \hookrightarrow X$  such that  $y \in \overline{\phi(C)}$ .*

PROOF. Consider an open dense subset  $U \hookrightarrow Y$  contained in the image  $\phi(X)$ . By the lemma there is a curve  $C' \hookrightarrow Y$  containing  $y$  and such that  $C' \cap U \neq \emptyset$ . Then, again applying the lemma to an irreducible component of  $\phi^{-1}(C')$  not contained in a fiber, we obtain an irreducible curve  $C \hookrightarrow X$  with  $\overline{\phi(C)} = \overline{C'}$ .  $\square$

Any affine variety  $X \hookrightarrow \mathbb{C}^k$  can also be equipped with the induced  $\mathbb{C}$ -topology (or *analytic topology*) from  $\mathbb{C}^k$  which is much finer than the *Zariski topology*. Usually there is no relation between the closure  $\overline{Z}^{\mathbb{C}}$  of a subset  $Z \hookrightarrow X$  in the  $\mathbb{C}$ -topology and the Zariski closure  $\overline{Z}$ .

LEMMA 2.10. *Let  $U \subset \mathbb{C}^k$  containing a subset  $V$  which is Zariski open and dense in  $\overline{U}$ . Then,*

$$\overline{U}^{\mathbb{C}} = \overline{U}$$

PROOF. By reducing to irreducible components, we may assume that  $\overline{U}$  is irreducible. Assume first that  $\dim \overline{U} = 1$ , that is,  $\overline{U}$  is an irreducible curve in  $\mathbb{C}^k$ . Let  $U_s$  be the subset of points where  $\overline{U}$  is a complex manifold, then  $\overline{U} - U_s$  is finite and by the *implicit function theorem* in analysis every  $u \in U_s$  has a  $\mathbb{C}$ -open neighborhood which is  $\mathbb{C}$ -homeomorphic to the complex line  $\mathbb{C}^1$ , whence the result holds in this case.

If  $\overline{U}$  is general and  $x \in \overline{U}$  we can take by the lemma above an irreducible curve  $C \hookrightarrow \overline{U}$  containing  $x$  and such that  $C \cap V \neq \emptyset$ . Then,  $C \cap V$  is Zariski open and dense in  $C$  and by the curve argument above  $x \in \overline{(C \cap V)}^{\mathbb{C}} \subset \overline{U}^{\mathbb{C}}$ . We can do this for any  $x \in \overline{U}$  finishing the proof.  $\square$

Consider the embedding of an affine variety  $X \hookrightarrow \mathbb{C}^k$ , proposition 2.6 and the fact that any constructible set  $Z$  contains a subset  $U$  which is open and dense in  $\overline{Z}$  we deduce from the lemma at once the next result.

PROPOSITION 2.11. *If  $Z$  is a constructible subset of an affine variety  $X$ , then*

$$\overline{Z}^{\mathbb{C}} = \overline{Z}$$

EXAMPLE 2.12. Let  $A$  be an affine  $\mathbb{C}$ -algebra and  $M \in \text{rep}_n A$ . We have proved in example 2.5 that the orbit  $\mathcal{O}(M) = GL_n \cdot M$  is Zariski open in its closure  $\overline{\mathcal{O}(M)}$ . Therefore, the orbit  $\mathcal{O}(M)$  is a constructible subset of  $\text{rep}_n A$ . By the proposition above, the Zariski closure  $\overline{\mathcal{O}(M)}$  of the orbit coincides with the closure of  $\mathcal{O}(M)$  in the  $\mathbb{C}$ -topology.

### 2.3. The Hilbert criterium.

A *one parameter subgroup* of a linear algebraic group  $G$  is a morphism

$$\lambda : \mathbb{C}^* \longrightarrow G$$

of affine algebraic groups. That is,  $\lambda$  is both a groupmorphism and a morphism of affine varieties. The set of all one parameter subgroup of  $G$  will be denoted by  $Y(G)$ .

If  $G$  is commutative algebraic group, then  $Y(G)$  is an Abelian group with additive notation

$$\lambda_1 + \lambda_2 : \mathbb{C}^* \longrightarrow G \quad \text{with } (\lambda_1 + \lambda_2)(t) = \lambda_1(t) \cdot \lambda_2(t)$$

Recall that an  $n$ -dimensional torus is an affine algebraic group isomorphic to

$$\underbrace{\mathbb{C}^* \times \dots \times \mathbb{C}^*}_n = T_n$$

the closed subgroup of invertible diagonal matrices in  $GL_n$ .

LEMMA 2.13.  $Y(T_n) \simeq \mathbb{Z}^n$ . The correspondence is given by assigning to  $(r_1, \dots, r_n) \in \mathbb{Z}^n$  the one-parameter subgroup

$$\lambda : \mathbb{C}^* \longrightarrow T_n \quad \text{given by } t \mapsto (t^{r_1}, \dots, t^{r_n})$$

PROOF. For any two affine algebraic groups  $G$  and  $H$  there is a canonical bijection  $Y(G \times H) = Y(G) \times Y(H)$  so it suffices to verify that  $Y(\mathbb{C}^*) \simeq \mathbb{Z}$  with any  $\lambda : \mathbb{C}^* \longrightarrow \mathbb{C}^*$  given by  $t \mapsto t^r$  for some  $r \in \mathbb{Z}$ . This is obvious as  $\lambda$  induces the algebra morphism

$$\mathbb{C}[\mathbb{C}^*] = \mathbb{C}[x, x^{-1}] \xrightarrow{\lambda^*} \mathbb{C}[x, x^{-1}] = \mathbb{C}[\mathbb{C}^*]$$

which is fully determined by the image of  $x$  which must be an invertible element. Now, any invertible element in  $\mathbb{C}[x, x^{-1}]$  is homogeneous of the form  $cx^r$  for some  $r \in \mathbb{Z}$  and  $c \in \mathbb{C}^*$ . The corresponding morphism maps  $t$  to  $ct^r$  which is only a groupmorphism if it maps the identity element 1 to 1 so  $c = 1$ , finishing the proof.  $\square$

PROPOSITION 2.14. Any one-parameter subgroup  $\lambda : \mathbb{C}^* \longrightarrow GL_n$  is of the form

$$t \mapsto g^{-1} \cdot \begin{bmatrix} t^{r_1} & & 0 \\ & \ddots & \\ 0 & & t^{r_n} \end{bmatrix} \cdot g$$

for some  $g \in GL_n$  and some  $n$ -tuple  $(r_1, \dots, r_n) \in \mathbb{Z}^n$ .

PROOF. Let  $H$  be the image under  $\lambda$  of the subgroup  $\mu_\infty$  of roots of unity in  $\mathbb{C}^*$ . We claim that there is a basechange matrix  $g \in GL_n$  such that

$$g.H.g^{-1} \hookrightarrow \begin{bmatrix} \mathbb{C}^* & & 0 \\ & \ddots & \\ 0 & & \mathbb{C}^* \end{bmatrix}$$

Assume  $h \in H$  not a scalar matrix, then  $h$  has a proper eigenspace decomposition  $V \oplus W = \mathbb{C}^n$ . We use that  $h^l = \mathbb{1}_n$  and hence all its Jordan blocks must have size

one as for any  $\lambda \neq 0$  we have

$$\begin{bmatrix} \lambda & 1 & & 0 \\ & \ddots & \ddots & \\ & & \ddots & 1 \\ & & & \lambda \end{bmatrix}^l = \begin{bmatrix} \lambda^l & l\lambda^{l-1} & & * \\ & \ddots & \ddots & \\ & & \ddots & l\lambda^{l-1} \\ & & & \lambda^l \end{bmatrix} \neq \mathbb{1}_n$$

Because  $H$  is commutative, both  $V$  and  $W$  are stable under  $H$ . By induction on  $n$  we may assume that the images of  $H$  in  $GL(V)$  and  $GL(W)$  are diagonalizable, but then the same holds in  $GL_n$ .

As  $\mu_\infty$  is infinite, it is Zariski dense in  $\mathbb{C}^*$  and because the diagonal matrices are Zariski closed in  $GL_n$  we have

$$g \cdot \lambda(\mathbb{C}^*) \cdot g^{-1} = g \cdot \overline{H} \cdot g^{-1} \hookrightarrow T_n$$

and the result follows from the lemma above  $\square$

Let  $V$  be a general  $GL_n$ -representation considered as an affine space with  $GL_n$ -action, let  $X$  be a  $GL_n$ -stable closed subvariety and consider a point  $x \in X$ . A one-parameter subgroup  $\mathbb{C}^* \xrightarrow{\lambda} GL_n$  determines a morphism

$$\mathbb{C}^* \xrightarrow{\lambda_x} X \quad \text{defined by} \quad t \mapsto \lambda(t).x$$

Observe that the image  $\lambda_x(\mathbb{C}^*)$  lies in the orbit  $GL_n.x$  of  $x$ . Assume there is a continuous extension of this map to the whole of  $\mathbb{C}$ . We claim that this extension must then be a morphism. If not, the induced algebra morphism

$$\mathbb{C}[X] \xrightarrow{\lambda_x^*} \mathbb{C}[t, t^{-1}]$$

does not have its image in  $\mathbb{C}[t]$ , so for some  $f \in \mathbb{C}[Z]$  we have that

$$\lambda_x^*(f) = \frac{a_0 + a_1 t + \dots + a_s t^s}{t^s} \quad \text{with} \quad a_0 \neq 0 \quad \text{and} \quad s > 0$$

But then  $\lambda_x^*(f)(t) \longrightarrow \pm \infty$  when  $t$  goes to zero, that is,  $\lambda_x^*$  cannot have a continuous extension, a contradiction.

So, if a continuous extension exists there is morphism  $\lambda_x : \mathbb{C} \longrightarrow X$ . Then,  $\lambda_x(0) = y$  and we denote this by

$$\lim_{t \rightarrow 0} \lambda(t).x = y$$

Clearly, the point  $y \in X$  must belong to the orbitclosure  $\overline{GL_n.x}$  in the Zariski topology (or in the  $\mathbb{C}$ -topology as orbits are constructible). Conversely, one might ask whether if  $y \in \overline{GL_n.x}$  we can always approach  $y$  via a one-parameter subgroup. The *Hilbert criterium* gives situations when this is indeed possible.

The only ideals of the formal power series  $\mathbb{C}[[t]]$  are principal and generated by  $t^r$  for some  $r \in \mathbb{N}_+$ . With  $\mathbb{C}((t))$  we will denote the field of fractions of the domain  $\mathbb{C}((t))$ .

LEMMA 2.15. *Let  $V$  be a  $GL_n$ -representation,  $v \in V$  and a point  $w \in V$  lying in the orbitclosure  $\overline{GL_n.v}$ . Then, there exists a matrix  $g$  with coefficients in the field  $\mathbb{C}((t))$  and  $\det(g) \neq 0$  such that*

$$(g.v)_{t=0} \quad \text{is well defined and is equal to} \quad w$$

PROOF. Note that  $g.v$  is a vector with coordinates in the field  $\mathbb{C}((t))$ . If all coordinates belong to  $\mathbb{C}[[t]]$  we can set  $t = 0$  in this vector and obtain a vector in  $V$ . It is this vector that we denote with  $(g.v)_{t=0}$ .

Consider the orbit map  $\mu : GL_n \longrightarrow V$  defined by  $g' \mapsto g'.v$ . As  $w \in \overline{GL_n.v}$  we have seen that there is an irreducible curve  $C \hookrightarrow GL_n$  such that  $w \in \overline{\mu(C)}$ . We obtain a diagram of  $\mathbb{C}$ -algebras

$$\begin{array}{ccccc} \mathbb{C}[GL_n] & \longrightarrow & \mathbb{C}[C] & \hookrightarrow & \mathbb{C}(C) \\ \uparrow \mu^* & & \uparrow \mu^* & & \uparrow \\ \mathbb{C}[V] & \longrightarrow & \mathbb{C}[\overline{\mu(C)}] & \hookrightarrow & \mathbb{C}[C'] \end{array}$$

Here,  $\mathbb{C}[C]$  is defined to be the integral closure of  $\mathbb{C}[\overline{\mu(C)}]$  in the functionfield  $\mathbb{C}(C)$  of  $C$ . Two things are important to note here :  $C' \longrightarrow \overline{\mu(C)}$  is finite, so surjective and take  $c \in C'$  be a point lying over  $w \in \overline{\mu(C)}$ . Further,  $C'$  having an integrally closed coordinate ring is a complex manifold. Hence, by the implicit function theorem polynomial functions on  $C$  can be expressed in a neighborhood of  $c$  as power series in one variable, giving an embedding  $\mathbb{C}[C'] \hookrightarrow \mathbb{C}[[t]]$  with  $(t) \cap \mathbb{C}[C'] = M_c$ . This inclusion extends to one on the level of their fields of fractions. That is, we have a diagram of  $\mathbb{C}$ -algebra morphisms

$$\begin{array}{ccccccc} \mathbb{C}[GL_n] & \longrightarrow & \mathbb{C}(C) & = & \mathbb{C}(C') & \hookrightarrow & \mathbb{C}((t)) \\ \uparrow \mu^* & & \uparrow & & \uparrow & & \uparrow \\ \mathbb{C}[V] & \longrightarrow & \mathbb{C}[\overline{\mu(C)}] & \hookrightarrow & \mathbb{C}[C'] & \hookrightarrow & \mathbb{C}[[t]] \end{array}$$

The upper composition defines an invertible matrix  $g(t)$  with coefficients in  $\mathbb{C}((t))$ , its  $(i, j)$ -entry being the image of the coordinate function  $x_{ij} \in \mathbb{C}[GL_n]$ . Moreover, the inverse image of the maximal ideal  $(t) \triangleleft \mathbb{C}[[t]]$  under the lower composition gives the maximal ideal  $M_w \triangleleft \mathbb{C}[V]$ . This proves the claim.  $\square$

LEMMA 2.16. *Let  $g$  be an  $n \times n$  matrix with coefficients in  $\mathbb{C}((t))$  and  $\det g \neq 0$ . Then there exist  $u_1, u_2 \in GL_n(\mathbb{C}[[t]])$  such that*

$$g = u_1 \cdot \begin{bmatrix} t^{r_1} & & & 0 \\ & \ddots & & \\ 0 & & & t^{r_n} \end{bmatrix} \cdot u_2$$

with  $r_i \in \mathbb{Z}$  and  $r_1 \leq r_2 \leq \dots \leq r_n$ .

PROOF. By multiplying  $g$  with a suitable power of  $t$  we may assume that  $g = (g_{ij}(t))_{i,j} \in M_n(\mathbb{C}[[t]])$ . If  $f(t) = \sum_{i=0}^{\infty} f_i t^i \in \mathbb{C}[[t]]$  define  $v(f(t))$  to be the minimal  $i$  such that  $a_i \neq 0$ . Let  $(i_0, j_0)$  be an entry where  $v(g_{ij}(t))$  attains a minimum, say  $r_1$ . That is, for all  $(i, j)$  we have  $g_{ij}(t) = t^{r_1} t^r f(t)$  with  $r \geq 0$  and  $f(t)$  an invertible element of  $\mathbb{C}[[t]]$ .

By suitable row and column interchanges we can take the entry  $(i_0, j_0)$  to the  $(1, 1)$ -position. Then, multiplying with a unit we can replace it by  $t^{r_1}$  and by elementary row and column operations all the remaining entries in the first row and



column can be made zero. That is, we have invertible matrices  $a_1, a_2 \in GL_n(\mathbb{C}[[t]])$  such that

$$g = a_1 \cdot \begin{bmatrix} t^{r_1} & \underline{0}^r \\ \underline{0} & \boxed{g_1} \end{bmatrix} \cdot a_2$$

Repeating the same idea on the submatrix  $g_1$  and continuing gives the result.  $\square$

We can now state and prove the *Hilbert criterium* which allows us to study orbit-closures by one parameter subgroups.

**THEOREM 2.17.** *Let  $V$  be a  $GL_n$ -representation and  $X \hookrightarrow V$  a closed  $GL_n$ -stable subvariety. Let  $\mathcal{O}(x) = GL_n \cdot x$  be the orbit of a point  $x \in X$ . Let  $Y \hookrightarrow \overline{\mathcal{O}(x)}$  be a closed  $GL_n$ -stable subset. Then, there exists a one-parameter subgroup  $\lambda : \mathbb{C}^* \rightarrow GL_n$  such that*

$$\lim_{t \rightarrow 0} \lambda(t) \cdot x \in Y$$

**PROOF.** It suffices to prove the result for  $X = V$ . By lemma 2.15 there is an invertible matrix  $g \in M_n(\mathbb{C}((t)))$  such that

$$(g \cdot x)_{t=0} = y \in Y$$

By lemma 2.16 we can find  $u_1, u_2 \in GL_n(\mathbb{C}[[t]])$  such that

$$g = u_1 \cdot \lambda'(t) \cdot u_2 \quad \text{with} \quad \lambda'(t) = \begin{bmatrix} t^{r_1} & & 0 \\ & \ddots & \\ 0 & & t^{r_n} \end{bmatrix}$$

a one-parameter subgroup. There exist  $x_i \in V$  such that  $u_2 \cdot x = \sum_{i=0}^{\infty} z_i t^i$  in particular  $u_2(0) \cdot x = x_0$ . But then,

$$\begin{aligned} (\lambda'(t) \cdot u_2 \cdot x)_{t=0} &= \sum_{i=0}^{\infty} (\lambda'(t) \cdot x_i t^i)_{t=0} \\ &= (\lambda'(t) \cdot x_0)_{t=0} + (\lambda'(t) \cdot x_1 t)_{t=0} + \dots \end{aligned}$$

But one easily verifies (using a basis of eigenvectors of  $\lambda'(t)$ ) that

$$\lim_{s \rightarrow 0} \lambda'^{-1}(s) \cdot (\lambda'(t) x_i t^i)_{t=0} = \begin{cases} (\lambda'(t) \cdot x_0)_{t=0} & \text{if } i = 0, \\ 0 & \text{if } i \neq 0 \end{cases}$$

As  $(\lambda'(t) \cdot u_2 \cdot x)_{t=0} \in Y$  and  $Y$  is a closed  $GL_n$ -stable subset, we also have that

$$\lim_{s \rightarrow 0} \lambda'^{-1}(s) \cdot (\lambda'(t) \cdot u_2 \cdot x)_{t=0} \in Y \quad \text{that is,} \quad (\lambda'(t) \cdot x_0)_{t=0} \in Y$$

But then, we have for the one-parameter subgroup  $\lambda(t) = u_2(0)^{-1} \cdot \lambda'(t) \cdot u_2(0)$  that

$$\lim_{t \rightarrow 0} \lambda(t) \cdot x \in Y$$

finishing the proof.  $\square$

An important special case occurs when  $x \in V$  belongs to the *nullcone*, that is, when the orbit closure  $\overline{\mathcal{O}(x)}$  contains the fixed point  $0 \in V$ . The original Hilbert criterium asserts the following.

**PROPOSITION 2.18.** *Let  $V$  be a  $GL_n$ -representation and  $x \in V$  in the nullcone. Then, there is a one-parameter subgroup  $\mathbb{C}^* \xrightarrow{\lambda} GL_n$  such that*

$$\lim_{t \rightarrow 0} \lambda(t) \cdot x = 0$$

In the statement of theorem 2.17 it is important that  $Y$  is closed. In particular, it does *not* follow that any orbit  $\mathcal{O}(y) \hookrightarrow \overline{\mathcal{O}(x)}$  can be reached via one-parameter subgroups, see example 2.21 below.

## 2.4. Semisimple modules

In this section we will characterize the closed  $GL_n$ -orbits in the representation variety  $\text{rep}_n A$  for an affine  $\mathbb{C}$ -algebra  $A$ . We have seen that any point  $\psi \in \text{rep}_n A$ , that is any  $n$ -dimensional representation  $A \xrightarrow{\psi} M_n$ , determines an  $n$ -dimensional  $A$ -module which we will denote with  $M_\psi$ .

A *finite filtration*  $F$  on an  $n$ -dimensional module  $M$  is a sequence of  $A$ -submodules

$$F : 0 = M_{t+1} \subset M_t \subset \dots \subset M_1 \subset M_0 = M.$$

The *associated graded*  $A$ -module is the  $n$ -dimensional module

$$\text{gr}_F M = \bigoplus_{i=0}^t M_i/M_{i+1}.$$

We have the following ringtheoretical interpretation of the action of one-parameter subgroups of  $GL_n$  on the representation variety  $\text{rep}_n A$ .

LEMMA 2.19. *Let  $\psi, \rho \in \text{rep}_n A$ . Equivalent are,*

- (1) *There is a one-parameter subgroup  $\mathbb{C}^* \xrightarrow{\lambda} GL_n$  such that*

$$\lim_{t \rightarrow 0} \lambda(t) \cdot \psi = \rho$$

- (2) *There is a finite filtration  $F$  on the  $A$ -module  $M_\psi$  such that*

$$\text{gr}_F M_\psi \simeq M_\rho$$

*as  $A$ -modules.*

PROOF. (1)  $\Rightarrow$  (2) : If  $V$  is any  $GL_n$ -representation and  $\mathbb{C}^* \xrightarrow{\lambda} GL_n$  a one-parameter subgroup, we have an induced *weight space decomposition* of  $V$

$$V = \bigoplus_i V_{\lambda,i} \quad \text{where} \quad V_{\lambda,i} = \{v \in V \mid \lambda(t) \cdot v = t^i v, \forall t \in \mathbb{C}^*\}.$$

In particular, we apply this to the underlying vectorspace of  $M_\psi$  which is  $V = \mathbb{C}^n$  (column vectors) on which  $GL_n$  acts by left multiplication. We define

$$M_j = \bigoplus_{i>j} V_{\lambda,i}$$

and claim that this defines a finite filtration on  $M_\psi$  with associated graded  $A$ -module  $M_\rho$ . For any  $a \in A$  (it suffices to vary  $a$  over the generators of  $A$ ) we can consider the linear maps

$$\phi_{ij}(a) : V_{\lambda,i} \hookrightarrow V = M_\psi \xrightarrow{a} M_\psi = V \twoheadrightarrow V_{\lambda,j}$$

(that is, we express the action of  $a$  in a blockmatrix  $\Phi_a$  with respect to the decomposition of  $V$ ). Then, the action of  $a$  on the module corresponding to  $\lambda(t) \cdot \psi$  is given by the matrix  $\Phi'_a = \lambda(t) \cdot \Phi_a \cdot \lambda(t)^{-1}$  with corresponding blocks

$$\begin{array}{ccc} V_{\lambda,i} & \xrightarrow{\phi_{ij}(a)} & V_{\lambda,j} \\ \uparrow \lambda(t)^{-1} & & \downarrow \lambda(t) \\ V_{\lambda,i} & \xrightarrow{\phi'_{ij}(a)} & V_{\lambda,j} \end{array}$$

that is  $\phi'_{ij}(a) = t^{j-i}\phi_{ij}(a)$ . Therefore, if  $\lim_{t \rightarrow 0} \lambda(t) \cdot \psi$  exists we must have that

$$\phi_{ij}(a) = 0 \quad \text{for all } j < i.$$

But then, the action by  $a$  sends any  $M_k = \bigoplus_{i>k} V_{\lambda,i}$  to itself, that is, the  $M_k$  are  $A$ -submodules of  $M_\psi$ . Moreover, for  $j > i$  we have

$$\lim_{t \rightarrow 0} \phi'_{ij}(a) = \lim_{t \rightarrow 0} t^{j-i} \phi_{ij}(a) = 0$$

Consequently, the action of  $a$  on  $\rho$  is given by the diagonal blockmatrix with blocks  $\phi_{ii}(a)$ , but this is precisely the action of  $a$  on  $V_i = M_{i-1}/M_i$ , that is,  $\rho$  corresponds to the associated graded module.

(2)  $\Rightarrow$  (1) : Given a finite filtration on  $M_\psi$

$$F \quad : \quad 0 = M_{t+1} \subset \dots \subset M_1 \subset M_0 = M_\psi$$

we have to find a one-parameter subgroup  $\mathbb{C}^* \xrightarrow{\lambda} GL_n$  which induces the filtration  $F$  as in the first part of the proof. Clearly, there exist subspaces  $V_i$  for  $0 \leq i \leq t$  such that

$$V = \bigoplus_{i=0}^t V_i \quad \text{and} \quad M_j = \bigoplus_{i=j}^t V_i.$$

Then we take  $\lambda$  to be defined by  $\lambda(t) = t^i Id_{V_i}$  for all  $i$  and it verifies the claims.  $\square$

EXAMPLE 2.20. Let  $M_\psi$  be the 2-dimensional  $\mathbb{C}[x]$ -module determined by the Jordan block and consider the canonical basevectors

$$\begin{bmatrix} \lambda & 1 \\ 0 & \lambda \end{bmatrix} \quad e_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad e_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

Then,  $\mathbb{C}e_1$  is a  $\mathbb{C}[x]$ -submodule of  $M_\psi$  and we have a filtration

$$0 = M_2 \subset \mathbb{C}e_1 = M_1 \subset \mathbb{C}e_1 \oplus \mathbb{C}e_2 = M_0 = M_\psi$$

Using the conventions of the second part of the above proof we then have

$$V_1 = \mathbb{C}e_1 \quad V_2 = \mathbb{C}e_2 \quad \text{hence} \quad \lambda(t) = \begin{bmatrix} t & 0 \\ 0 & 1 \end{bmatrix}$$

Indeed, we then obtain that


$$\begin{bmatrix} t & 0 \\ 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} \lambda & 1 \\ 0 & \lambda \end{bmatrix} \cdot \begin{bmatrix} t^{-1} & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} \lambda & t \\ 0 & \lambda \end{bmatrix}$$

and the limit  $t \rightarrow 0$  exists and is the associated graded module  $gr_F M_\psi = M_\rho$  determined by the diagonal matrix.

Consider two modules  $M_\psi, M_\psi \in rep_n A$ . Assume that  $\mathcal{O}(M_\rho) \hookrightarrow \overline{\mathcal{O}(M_\psi)}$  and that we can reach the orbit of  $M_\rho$  via a one-parameter subgroup. Then, lemma 2.19 asserts that  $M_\rho$  must be *decomposable* as it is the associated graded of a nontrivial filtration on  $M_\psi$ . This gives us a criterium to construct examples showing that the closedness assumption in the formulation of Hilbert's criterium is essential.

EXAMPLE 2.21. (Nullcone of  $M_3^2 = M_3 \oplus M_3$ )

In chapter 6 we will describe a method to determine the nullcones of  $m$ -tuples of  $n \times n$  matrices. The special case of  $2 \times 3 \times 3$  matrices has been worked out by H.P. Kraft in [44, p.202]. The orbits are depicted in figure 1 In this picture, each node corresponds to a torus. The right hand number is the dimension of the torus and the left hand number is the dimension of the orbit represented by a point in the

FIGURE 1. Kraft's diamond describing the nullcone of  $M_3^2$ .

torus. The solid or dashed lines indicate orbit closures. For example, the dashed line corresponds to the following two points in  $M_3^2 = M_3 \oplus M_3$

$$\psi = \left( \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \right) \quad \rho = \left( \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \right)$$

We claim that  $M_\rho$  is an indecomposable 3-dimensional module of  $\mathbb{C}\langle x, y \rangle$ . Indeed, the only subspace of the column vectors  $\mathbb{C}^3$  left invariant under both  $x$  and  $y$  is equal to

$$\begin{bmatrix} \mathbb{C} \\ 0 \\ 0 \end{bmatrix}$$

hence  $M_\rho$  cannot have a direct sum decomposition of two or more modules. Next, we claim that  $\mathcal{O}(M_\rho) \xrightarrow{\text{closure}} \overline{\mathcal{O}(M_\psi)}$ . Indeed, simultaneous conjugating  $\psi$  with the invertible matrix

$$\begin{bmatrix} 1 & \epsilon^{-1} - 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \epsilon^{-1} \end{bmatrix} \quad \text{we obtain the couple} \quad \left( \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & \epsilon \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \right)$$

and letting  $\epsilon \rightarrow 0$  we see that the limiting point is  $\rho$ .

The *Jordan-Hölder theorem*, see for example [63, 2.6] asserts that any finite dimensional  $A$ -module  $M$  has a *composition series*, that is,  $M$  has a finite filtration

$$F \quad 0 = M_{t+1} \subset M_t \subset \dots \subset M_1 \subset M_0 = M$$

such that the successive quotients  $S_i = M_i/M_{i+1}$  are all *simple*  $A$ -modules for  $0 \leq i \leq t$ . Moreover, these *composition factors*  $S$  and their *multiplicities* are independent of the chosen composition series, that is, the set  $\{S_0, \dots, S_t\}$  is the same for every composition series. In particular, the associated graded module for a composition series is determined only up to isomorphism and is the semisimple  $n$ -dimensional module

$$gr M = \bigoplus_{i=0}^t S_i$$

THEOREM 2.22. *Let  $A$  be an affine  $\mathbb{C}$ -algebra and  $M \in \text{rep}_n A$ .*

- (1) *The orbit  $\mathcal{O}(M)$  is closed in  $\text{rep}_n A$  if and only if  $M$  is an  $n$ -dimensional semisimple  $A$ -module.*
- (2) *The orbitclosure  $\overline{\mathcal{O}(M)}$  contains exactly one closed orbit, corresponding to the direct sum of the composition factors of  $M$ .*
- (3) *The points of the quotient variety of  $\text{rep}_n A$  under  $GL_n$  parameterize the isomorphism classes of  $n$ -dimensional semisimple  $A$ -modules. We will denote the quotient variety by  $\text{iss}_n A$ .*

PROOF. (1) : Assume that the orbit  $\mathcal{O}(M)$  is Zariski closed. Let  $gr M$  be the associated graded module for a composition series of  $M$ . From lemma 2.19 we know that  $\mathcal{O}(gr M)$  is contained in  $\overline{\mathcal{O}(M)} = \mathcal{O}(M)$ . But then  $gr M \simeq M$  whence  $M$  is semisimple.

Conversely, assume  $M$  is semisimple. We know that the orbitclosure  $\overline{\mathcal{O}(M)}$  contains a closed orbit, say  $\mathcal{O}(N)$ . By the Hilbert criterium we have a one-parameter subgroup  $\mathbb{C}^* \xrightarrow{\lambda} GL_n$  such that


$$\lim_{t \rightarrow 0} \lambda(t).M = N' \simeq N.$$

By lemma 2.19 this means that there is a finite filtration  $F$  on  $M$  with associated graded module  $gr_F M \simeq N$ . For the semisimple module  $M$  the only possible finite filtrations are such that each of the submodules is a direct sum of simple components, so  $gr_F M \simeq M$ , whence  $M \simeq N$  and hence the orbit  $\mathcal{O}(M)$  is closed.

(2) : Remains only to prove uniqueness of the closed orbit in  $\overline{\mathcal{O}(M)}$ . This either follows from the Jordan-Hölder theorem or, alternatively, from the separation property of the quotient map to be proved in the next section.

(3) : We will prove in the next section that the points of a quotient variety parameterize the closed orbits.  $\square$

EXAMPLE 2.23. Recall the description of the orbits in  $M_2^2 = M_2 \oplus M_2$  given in the previous chapter.



and each fiber contains a unique closed orbit. The one over a point in  $H - S$  corresponding to the matrix couple

$$\left( \begin{bmatrix} a_1 & 0 \\ 0 & a_2 \end{bmatrix}, \begin{bmatrix} b_1 & 0 \\ 0 & b_2 \end{bmatrix} \right)$$

which is indeed a semi-simple module of  $\mathbb{C}\langle x, y \rangle$  (the direct sum of the two 1-dimensional simple representations determined by  $x \mapsto a_i$  and  $y \mapsto b_i$ ). In case  $a_1 = a_2$  and  $b_1 = b_2$  these two simples coincide and the semi-simple module having this factor with multiplicity two is the unique closed orbit in the fiber of a point in  $S$ .

EXAMPLE 2.24. Assume  $A$  is a finite dimensional  $\mathbb{C}$ -algebra. Then, there are only a finite number, say  $k$ , of nonisomorphic  $n$ -dimensional semisimple  $A$ -modules. Hence  $\text{iss}_n A$  is a finite number of  $k$  points, whence  $\text{rep}_n A$  is the disjoint union of  $k$  connected components, each consisting of all  $n$ -dimensional  $A$ -modules with the same composition factors. Connectivity follows from the fact that the orbit of the sum of the composition factors lies in the closure of each orbit.

EXAMPLE 2.25. Let  $A$  be an affine commutative algebra with presentation  $A = \mathbb{C}[x_1, \dots, x_m]/I_A$  and let  $X$  be the affine variety  $\mathbb{V}(I_A)$ . Observe that any simple  $A$ -module is one-dimensional hence corresponds to a point in  $X$ . (Indeed, for any algebra  $A$  a simple  $k$ -dimensional module determines an epimorphism  $A \twoheadrightarrow M_k$  and  $M_k$  is only commutative if  $k = 1$ ). Applying the Jordan-Hölder theorem we see that

$$\text{iss}_n A \simeq X^{(n)} = \underbrace{X \times \dots \times X}_n / S_n$$

the  $n$ -th symmetric product of  $X$ .

## 2.5. Some invariant theory.

The results in this section hold for any reductive algebraic group. Because we work with  $GL_n$  (or later with products  $GL(\alpha) = GL_{a_1} \times \dots \times GL_{a_k}$ ) we include a proof in this case. Our first aim is to prove that  $GL_n$  is a *reductive group*, that is, all  $GL_n$ -representations are completely reducible. Consider the *unitary group*

$$U_n = \{A \in GL_n \mid A.A^* = \mathbb{1}_n\}$$

where  $A^*$  is the *Hermitian transpose* of  $A$ . Clearly,  $U_n$  is a compact Lie group. Any compact Lie group has a so called *Haar measure* which allows one to integrate continuous complex valued functions over the group in an invariant way. That is, there is a linear function assigning to every continuous function  $f : U_n \rightarrow \mathbb{C}$  its integral

$$f \mapsto \int_{U_n} f(g) dg \in \mathbb{C}$$

which is normalized such that  $\int_{U_n} dg = 1$  and is left and right invariant, which means that for all  $u \in U_n$  we have the equalities

$$\int_{U_n} f(gu) dg = \int_{U_n} f(g) dg = \int_{U_n} f(ug) dg.$$

This integral replaces the classical idea in representation theory of averaging functions over a finite group.

PROPOSITION 2.26. *Every  $U_n$ -representation is completely reducible.*

PROOF. Take a finite dimensional complex vectorspace  $V$  with a  $U_n$ -action and assume that  $W$  is a subspace of  $V$  left invariant under this action. Extending a basis of  $W$  to  $V$  we get a linear map  $V \xrightarrow{\phi} W$  which is the identity on  $W$ . For any  $v \in V$  we have a continuous map

$$U_n \longrightarrow W \quad g \mapsto g.\phi(g^{-1}.v)$$

(use that  $W$  is left invariant) and hence we can integrate it over  $U_n$  (integrate the coordinate functions). Hence we can define a map  $\phi_0 : V \longrightarrow W$  by

$$\phi_0(v) = \int_{U_n} g.\phi(g^{-1}.v)dg.$$

Clearly,  $\phi_0$  is linear and is the identity on  $W$ . Moreover,

$$\begin{aligned} \phi_0(u.v) &= \int_{U_n} g.\phi(g^{-1}u.v)dg = u. \int_{U_n} u^{-1}g.\phi(g^{-1}u.v)dg \\ &\stackrel{*}{=} u. \int_{U_n} g\phi(g^{-1}.v)dg = u.\phi_0(v) \end{aligned}$$

where the starred equality uses the invariance of the Haar measure. Hence,  $V = W \oplus \text{Ker } \phi_0$  is a decomposition as  $U_n$ -representations. Continuing whenever one of the components has a nontrivial subrepresentation we arrive at a decomposition of  $V$  into simple  $U_n$ -representations.  $\square$

We claim that for any  $n$ ,  $U_n$  is Zariski dense in  $GL_n$ . Let  $D_n$  be the group of all diagonal matrices in  $GL_n$ . The *Cartan decomposition* for  $GL_n$  asserts that

$$GL_n = U_n.D_n.U_n$$

For, take  $g \in GL_n$  then  $g.g^*$  is an Hermitian matrix and hence diagonalizable by unitary matrices. So, there is a  $u \in U_n$  such that

$$u^{-1}.g.g^*.u = \begin{bmatrix} \alpha_1 & & \\ & \ddots & \\ & & \alpha_n \end{bmatrix} = \underbrace{s^{-1}.g.s}_p . \underbrace{s^{-1}.g^*.s}_{p^*}$$

Then, each  $\alpha_i > 0 \in \mathbb{R}$  as  $\alpha_i = \sum_{j=1}^n \|p_{ij}\|^2$ . Let  $\beta_i = \sqrt{\alpha_i}$  and let  $d$  the diagonal matrix  $\text{diag}(\beta_1, \dots, \beta_n)$ . Clearly,

$$g = u.d.(d^{-1}.u^{-1}.g) \quad \text{and we claim} \quad v = d^{-1}.u^{-1}.g \in U_n.$$

Indeed, we have

$$\begin{aligned} v.v^* &= (d^{-1}.u^{-1}.g).(g^*.u.d^{-1}) = d^{-1}.(u^{-1}.g.g^*.u).d^{-1} \\ &= d^{-1}.d^2.d^{-1} = \mathbb{1}_n \end{aligned}$$

proving the Cartan decomposition. Now,  $D_n = \mathbb{C}^* \times \dots \times \mathbb{C}^*$  and  $D_n \cap U_n = U_1 \times \dots \times U_1$  and because  $U_1 = \mu$  is Zariski dense (being infinite) in  $D_1 = \mathbb{C}^*$ , we have that  $D_n$  is contained in the Zariski closure of  $U_n$ . By the Cartan decomposition we then have that the Zariski closure of  $U_n$  is  $GL_n$ .

**THEOREM 2.27.**  *$GL_n$  is a reductive group. That is, all  $GL_n$ -representations are completely reducible.*

**PROOF.** Let  $V$  be a  $GL_n$ -representation having a subrepresentation  $W$ . In particular,  $V$  and  $W$  are  $U_n$ -representations, so by the foregoing proposition we have a decomposition  $V = W \oplus W'$  as  $U_n$ -representations. Consider the subgroup

$$N = N_{GL_n}(W') = \{g \in GL_n \mid g.W' \subset W'\}$$

then  $N$  is a Zariski closed subgroup of  $GL_n$  containing  $U_n$ . As the Zariski closure of  $U_n$  is  $GL_n$  we have  $N = GL_n$  and hence that  $W'$  is a representation of  $GL_n$ . Continuing gives a decomposition of  $V$  in simple  $GL_n$ -representations.  $\square$

Let  $S = S_{GL_n}$  be the set of isomorphism classes of simple  $GL_n$ -representations. If  $W$  is a simple  $GL_n$ -representation belonging to the isomorphism class  $s \in S$ , we say that  $W$  is of *type*  $s$  and denote this by  $W \in s$ . Let  $X$  be a complex vectorspace (not necessarily finite dimensional) with a linear action of  $GL_n$ . We say that the action is *locally finite* on  $X$  if, for any finite dimensional subspace  $Y$  of  $X$ , there exists a finite dimensional subspace  $Y \subset Y' \subset X$  which is a  $GL_n$ -representation. The *isotypical component* of  $X$  of type  $s \in S$  is defined to be the subspace

$$X_{(s)} = \sum \{W \mid W \subset X, W \in s\}.$$

If  $V$  is a  $GL_n$ -representation, we have seen that  $V$  is completely reducible. Then,  $V = \oplus V_{(s)}$  and every isotypical component  $V_{(s)} \simeq W^{\oplus e_s}$  for  $W \in s$  and some number  $e_s$ . Clearly,  $e_s \neq 0$  for only finitely many classes  $s \in S$ . We call the decomposition  $V = \oplus_{s \in S} V_{(s)}$  the *isotypical decomposition* of  $V$  and we say that the simple representation  $W \in s$  occurs *with multiplicity*  $e_s$  in  $V$ .

If  $V'$  is another  $GL_n$ -representation and if  $V \xrightarrow{\phi} V'$  is a morphism of  $GL_n$ -representations (that is, a linear map commuting with the action), then for any  $s \in S$  we have that  $\phi(V_{(s)}) \subset V'_{(s)}$ . If the action of  $GL_n$  on  $X$  is locally finite, we can reduce to finite dimensional  $GL_n$ -subrepresentation and obtain a decomposition

$$X = \oplus_{s \in S} X_{(s)},$$

which is again called the *isotypical decomposition* of  $X$ .

Let  $V$  be a  $GL_n$ -representation of some dimension  $m$ . Then, we can view  $V$  as an affine space  $\mathbb{C}^m$  and we have an induced action of  $GL_n$  on the polynomial functions  $f \in \mathbb{C}[V]$  by the rule

$$\begin{array}{ccc} V & \xrightarrow{f} & \mathbb{C} \\ \downarrow g & \searrow g.f & \\ V & & \end{array}$$

that is  $(g.f)(v) = f(g^{-1}.v)$  for all  $g \in GL_n$  and all  $v \in V$ . If  $\mathbb{C}[V] = \mathbb{C}[x_1, \dots, x_m]$  is graded by giving all the  $x_i$  degree one, then each of the homogeneous components of  $\mathbb{C}[V]$  is a finite dimensional  $GL_n$ -representation. Hence, the action of  $GL_n$  on  $\mathbb{C}[V]$  is locally finite. Indeed, let  $\{y_1, \dots, y_l\}$  be a basis of a finite dimensional subspace  $Y \subset \mathbb{C}[V]$  and let  $d$  be the maximum of the  $\deg(y_i)$ . Then  $Y' = \oplus_{i=0}^d \mathbb{C}[V]_i$  is a  $GL_n$ -representation containing  $Y$ .

Therefore, we have an isotypical decomposition  $\mathbb{C}[V] = \oplus_{s \in S} \mathbb{C}[V]_{(s)}$ . In particular, if  $0 \in S$  denotes the isomorphism class of the trivial  $GL_n$ -representation ( $\mathbb{C}_{triv} = \mathbb{C}x$  with  $g.x = x$  for every  $g \in GL_n$ ) then we have

$$\mathbb{C}[V]_{(0)} = \{f \in \mathbb{C}[V] \mid g.f = f, \forall g \in GL_n\} = \mathbb{C}[V]^{GL_n}$$

the ring of *polynomial invariants*, that is, of polynomial functions which are constant along orbits in  $V$ .

LEMMA 2.28. *Let  $V$  be a  $GL_n$ -representation.*

(1) *Let  $I \triangleleft \mathbb{C}[V]$  be a  $GL_n$ -stable ideal, that is,  $g.I \subset I$  for all  $g \in GL_n$ , then*

$$(\mathbb{C}[V]/I)^{GL_n} \simeq \mathbb{C}[V]^{GL_n} / (I \cap \mathbb{C}[V]^{GL_n}).$$



(2) Let  $J \triangleleft \mathbb{C}[V]^{GL_n}$  be an ideal, then we have a lying-over property

$$J = J\mathbb{C}[V] \cap \mathbb{C}[V]^{GL_n}.$$

Hence,  $\mathbb{C}[V]^{GL_n}$  is Noetherian, that is, every increasing chain of ideals stabilizes.

(3) Let  $I_j$  be a family of  $GL_n$ -stable ideals of  $\mathbb{C}[V]$ , then

$$\left(\sum_j I_j\right) \cap \mathbb{C}[V]^{GL_n} = \sum_j (I_j \cap \mathbb{C}[V]^{GL_n}).$$

PROOF. (1) : As  $I$  has the induced  $GL_n$ -action which is locally finite we have the isotypical decomposition  $I = \bigoplus I_{(s)}$  and clearly  $I_{(s)} = \mathbb{C}[V]_{(s)} \cap I$ . But then also, taking quotients we have

$$\bigoplus_s (\mathbb{C}[V]/I)_{(s)} = \mathbb{C}[V]/I = \bigoplus_s \mathbb{C}[V]_{(s)}/I_{(s)}.$$

Therefore,  $(\mathbb{C}[V]/I)_{(s)} = \mathbb{C}[V]_{(s)}/I_{(s)}$  and taking the special case  $s = 0$  is the statement.

(2) : For any  $f \in \mathbb{C}[V]^{GL_n}$  left-multiplication by  $f$  in  $\mathbb{C}[V]$  commutes with the  $GL_n$ -action, whence  $f \cdot \mathbb{C}[V]_{(s)} \subset \mathbb{C}[V]_{(s)}$ . That is,  $\mathbb{C}[V]_{(s)}$  is a  $\mathbb{C}[V]^{GL_n}$ -module. But then, as  $J \subset \mathbb{C}[V]^{GL_n}$  we have

$$\bigoplus_s (J\mathbb{C}[V])_{(s)} = J\mathbb{C}[V] = \bigoplus_s J\mathbb{C}[V]_{(s)}.$$

Therefore,  $(J\mathbb{C}[V])_{(s)} = J\mathbb{C}[V]_{(s)}$  and again taking the special value  $s = 0$  we obtain  $J\mathbb{C}[V] \cap \mathbb{C}[V]^{GL_n} = (J\mathbb{C}[V])_{(0)} = J$ . The Noetherian statement follows from the fact that  $\mathbb{C}[V]$  is Noetherian (the Hilbert basis theorem).

(3) : For any  $j$  we have the decomposition  $I_j = \bigoplus_s (I_j)_{(s)}$ . But then, we have

$$\bigoplus_s \left(\sum_j I_j\right)_{(s)} = \sum_j I_j = \sum_j \bigoplus_s (I_j)_{(s)} = \bigoplus_s \sum_j (I_j)_{(s)}.$$

Therefore,  $(\sum_j I_j)_{(s)} = \sum_j (I_j)_{(s)}$  and taking  $s = 0$  gives the required statement.  $\square$

**THEOREM 2.29.** *Let  $V$  be a  $GL_n$ -representation. Then, the ring of polynomial invariants  $\mathbb{C}[V]^{GL_n}$  is an affine  $\mathbb{C}$ -algebra.*

PROOF. Because the action of  $GL_n$  on  $\mathbb{C}[V]$  preserves the gradation, the ring of invariants is also graded

$$\mathbb{C}[V]^{GL_n} = R = \mathbb{C} \oplus R_1 \oplus R_2 \oplus \dots$$

From lemma 2.28(2) we know that  $\mathbb{C}[V]^{GL_n}$  is Noetherian and hence the ideal  $R_+ = R_1 \oplus R_2 \oplus \dots$  is finitely generated  $R_+ = Rf_1 + \dots + Rf_l$  by homogeneous elements  $f_1, \dots, f_l$ . We claim that as a  $\mathbb{C}$ -algebra  $\mathbb{C}[V]^{GL_n}$  is generated by the  $f_i$ . Indeed, we have  $R_+ = \sum_{i=1}^l \mathbb{C}f_i + R_+^2$  and then also

$$R_+^2 = \sum_{i,j=1}^l \mathbb{C}f_i f_j + R_+^3$$

and iterating this procedure we obtain for all powers  $m$  that

$$R_+^m = \sum_{\sum m_i = m} \mathbb{C}f_1^{m_1} \dots f_l^{m_l} + R_+^{m+1}.$$

Now, consider the subalgebra  $\mathbb{C}[f_1, \dots, f_l]$  of  $R = \mathbb{C}[V]^{GL_n}$ , then we obtain for any power  $d > 0$  that

$$\mathbb{C}[V]^{GL_n} = \mathbb{C}[f_1, \dots, f_l] + R_+^d.$$

For any  $i$  we then have for the homogeneous components of degree  $i$

$$R_i = \mathbb{C}[f_1, \dots, f_l]_i + (R_+^d)_i.$$

Now, if  $d > i$  we have that  $(R_+^d)_i = 0$  and hence that  $R_i = \mathbb{C}[f_1, \dots, f_l]_i$ . As this holds for all  $i$  we proved the claim.  $\square$

Choose generating invariants  $f_1, \dots, f_l$  of  $\mathbb{C}[V]^{GL_n}$ , consider the morphism

$$V \xrightarrow{\phi} \mathbb{C}^l \quad \text{defined by} \quad v \mapsto (f_1(v), \dots, f_l(v))$$

and define  $W$  to be the Zariski closure  $\overline{\phi(V)}$  in  $\mathbb{C}^l$ . Then, we have a diagram

$$\begin{array}{ccc} V & \xrightarrow{\phi} & \mathbb{C}^l \\ & \searrow \cong & \uparrow \\ & & W \end{array}$$

and an isomorphism  $\mathbb{C}[W] \xrightarrow{\pi^*} \mathbb{C}[V]^{GL_n}$ . More general, let  $X$  be a closed  $GL_n$ -stable subvariety of  $V$ , then  $X = \mathbb{V}_V(I)$  for some  $GL_n$ -stable ideal  $I$  of  $\mathbb{C}[V]$ . From lemma 2.28(1) we obtain

$$\mathbb{C}[X]^{GL_n} = (\mathbb{C}[V]/I)^{GL_n} = \mathbb{C}[V]^{GL_n} / (I \cap \mathbb{C}[V]^{GL_n})$$

whence  $\mathbb{C}[X]^{GL_n}$  is also an affine algebra (and generated by the images of the  $f_i$ ). Define  $Y$  to be the Zariski closure of  $\phi(X)$  in  $\mathbb{C}^l$ , then we have a diagram

$$\begin{array}{ccc} X & \xrightarrow{\phi} & \mathbb{C}^l \\ & \searrow \cong & \uparrow \\ & & Y \end{array}$$

and an isomorphism  $\mathbb{C}[Y] \xrightarrow{\pi} \mathbb{C}[X]^{GL_n}$ . We call the morphism  $X \xrightarrow{\pi} Y$  an *algebraic quotient* of  $X$  under  $GL_n$ . We will now prove some important properties of this quotient.

**PROPOSITION 2.30 (universal property).** *If  $X \xrightarrow{\mu} Z$  is a morphism which is constant along  $GL_n$ -orbits in  $X$ , then there exists a unique factoring morphism  $\bar{\mu}$*

$$\begin{array}{ccc} X & \xrightarrow{\pi} & Y \\ & \searrow \mu & \uparrow \bar{\mu} \\ & & Z \end{array}$$

PROOF. As  $\mu$  is constant along  $GL_n$ -orbits in  $X$ , we have an inclusion  $\mu^*(\mathbb{C}[Z]) \subset \mathbb{C}[X]^{GL_n}$ . We have the commutative diagram

$$\begin{array}{ccc}
 & \mathbb{C}[X] & \\
 & \uparrow & \\
 & \mathbb{C}[X]^{GL_n} & \\
 \nearrow \mu^* & & \nwarrow \mu_* \\
 \mathbb{C}[Z] & \xrightarrow{\quad \bar{\mu}^* \quad} & \mathbb{C}[Y]
 \end{array}$$

from which the existence and uniqueness of  $\bar{\mu}$  follows.  $\square$

As a consequence, an algebraic quotient is uniquely determined up to isomorphism (that is, we might have started from other generating invariants and still obtain the same quotient variety up to isomorphism).

PROPOSITION 2.31 (onto property). *The algebraic quotient  $X \xrightarrow{\pi} Y$  is surjective. Moreover, if  $Z \hookrightarrow X$  is a closed  $GL_n$ -stable subset, then  $\pi(Z)$  is closed in  $Y$  and the morphism*

$$\pi_X|_Z : Z \longrightarrow \pi(Z)$$

*is an algebraic quotient, that is,  $\mathbb{C}[\pi(Z)] \simeq \mathbb{C}[Z]^{GL_n}$ .*

PROOF. Let  $y \in Y$  with maximal ideal  $M_y \triangleleft \mathbb{C}[Y]$ . By lemma 2.28(2) we have  $M_y \mathbb{C}[X] \neq \mathbb{C}[X]$  and hence there is a maximal ideal  $M_x$  of  $\mathbb{C}[X]$  containing  $M_y \mathbb{C}[X]$ , but then  $\pi(x) = y$ . Let  $Z = \mathbb{V}_X(I)$  for a  $G$ -stable ideal  $I$  of  $\mathbb{C}[X]$ , then  $\overline{\pi(Z)} = \mathbb{V}_Y(I \cap \mathbb{C}[Y])$ . That is,  $\mathbb{C}[\overline{\pi(Z)}] = \mathbb{C}[Y]/(I \cap \mathbb{C}[Y])$ . However, we have from lemma 2.28(1) that

$$\mathbb{C}[Y]/(\mathbb{C}[Y] \cap I) \simeq (\mathbb{C}[X]/I)^{GL_n} = \mathbb{C}[Z]^{GL_n}$$

and hence  $\mathbb{C}[\overline{\pi(Z)}] = \mathbb{C}[Z]^{GL_n}$ . Finally, surjectivity of  $\pi|_Z$  is proved as above.  $\square$

An immediate consequence is that the Zariski topology on  $Y$  is the quotient topology of that on  $X$ . For, take  $U \subset Y$  with  $\pi^{-1}(U)$  Zariski open in  $X$ . Then,  $X - \pi^{-1}(U)$  is a  $GL_n$ -stable closed subset of  $X$ . Then,  $\pi(X - \pi^{-1}(U)) = Y - U$  is Zariski closed in  $Y$ .

PROPOSITION 2.32 (separation property). *The quotient  $X \xrightarrow{\pi} Y$  separates disjoint closed  $GL_n$ -stable subvarieties of  $X$ .*

PROOF. Let  $Z_j$  be closed  $GL_n$ -stable subvarieties of  $X$  with defining ideals  $Z_j = \mathbb{V}_X(I_j)$ . Then,  $\cap_j Z_j = \mathbb{V}_X(\sum_j I_j)$ . Applying lemma 2.28(3) we obtain

$$\begin{aligned}
 \overline{\pi(\cap_j Z_j)} &= \mathbb{V}_Y((\sum_j I_j) \cap \mathbb{C}[Y]) = \mathbb{V}_Y(\sum_j (I_j \cap \mathbb{C}[Y])) \\
 &= \cap_j \mathbb{V}_Y(I_j \cap \mathbb{C}[Y]) = \cap_j \overline{\pi(Z_j)}
 \end{aligned}$$

The onto property implies that  $\overline{\pi(Z_j)} = \pi(Z_j)$  from which the statement follows.  $\square$

It follows from the universal property that the quotient variety  $Y$  determined by the ring of polynomial invariants  $\mathbb{C}[Y]^{GL_n}$  is the best algebraic approximation to the orbit space problem. From the separation property a stronger fact follows.

**PROPOSITION 2.33.** *The algebraic quotient  $X \xrightarrow{\pi} Y$  is the best continuous approximation to the orbit space. That is, points of  $Y$  parameterize the closed  $GL_n$ -orbits in  $X$ . In fact, every fiber  $\pi^{-1}(y)$  contains exactly one closed orbit  $C$  and we have*

$$\pi^{-1}(y) = \{x \in X \mid C \subset \overline{GL_n \cdot x}\}$$

**PROOF.** The fiber  $F = \pi^{-1}(y)$  is a  $GL_n$ -stable closed subvariety of  $X$ . Take any orbit  $GL_n \cdot x \subset F$  then either it is closed or contains in its closure an orbit of strictly smaller dimension. Induction on the dimension then shows that  $\overline{GL_n \cdot x}$  contains a closed orbit  $C$ . On the other hand, assume that  $F$  contains two closed orbits, then they have to be disjoint contradicting the separation property.  $\square$

## 2.6. Geometric reconstruction

In this section we will give a geometric interpretation of the reconstruction result of theorem 1.28. Let  $A$  be a Cayley-Hamilton algebra of degree  $n$ , with trace map  $tr_A$ , which is generated by at most  $m$  elements  $a_1, \dots, a_m$ . We will give a functorial interpretation to the affine scheme determined by the canonical ideal  $N_A \triangleleft \mathbb{C}[M_n^m]$  in the formulation of theorem 1.28. First, let us identify the reduced affine variety  $\mathbb{V}(N_A)$ . A point  $m = (m_1, \dots, m_m) \in \mathbb{V}(N_A)$  determines an algebra map  $f_m : \mathbb{C}[M_n^m]/N_A \longrightarrow \mathbb{C}$  and hence an algebra map  $\phi_m$

$$\begin{array}{ccc} A & \xrightarrow{\phi_m} & M_n(\mathbb{C}) \\ \downarrow & \nearrow M_n(f_m) & \\ M_n(\mathbb{C}[M_n^m]/N_A) & & \end{array}$$

which is trace preserving. Conversely, from the universal property it follows that any trace preserving algebra morphism  $A \longrightarrow M_n(\mathbb{C})$  is of this form by considering the images of the trace generators  $a_1, \dots, a_m$  of  $A$ . Alternatively, the points of  $\mathbb{V}(N_A)$  parameterize  $n$ -dimensional *trace preserving* representations of  $A$ . That is,  $n$ -dimensional representations for which the morphism  $A \longrightarrow M_n(\mathbb{C})$  describing the action is trace preserving. For this reason we will denote the variety  $\mathbb{V}(N_A)$  by  $rep_n^{tr} A$  and call it the *trace preserving reduced representation variety* of  $A$ .

Assume that  $A$  is generated as a  $\mathbb{C}$ -algebra by  $a_1, \dots, a_m$  (observe that this is no restriction as trace affine algebras are affine) then clearly  $I_A(n) \subset N_A$ . That is,

**LEMMA 2.34.** *For  $A$  a Cayley-Hamilton algebra of degree  $n$  generated by  $\{a_1, \dots, a_m\}$ , the reduced trace preserving representation variety*

$$rep_n^{tr} A \hookrightarrow rep_n A$$

*is a closed subvariety of the reduced representation variety.*

It is easy to determine the additional defining equations. For, write any trace monomial out in the generators

$$tr_A(a_{i_1} \dots a_{i_k}) = \sum \alpha_{j_1 \dots j_l} a_{j_1} \dots a_{j_l}$$

then for a point  $m = (m_1, \dots, m_m) \in \text{rep}_n A$  to belong to  $\text{rep}_n^{\text{tr}} A$ , it must satisfy all the relations of the form

$$\text{tr}(m_{i_1} \dots m_{i_k}) = \sum \alpha_{j_1 \dots j_l} m_{j_1} \dots m_{j_l}$$

with  $\text{tr}$  the usual trace on  $M_n(\mathbb{C})$ . These relations define the closed subvariety  $\text{rep}_n^{\text{tr}}(A)$ . Usually, this is a proper subvariety.

EXAMPLE 2.35. Let  $A$  be a finite dimensional semi-simple algebra  $A = M_{d_1}(\mathbb{C}) \oplus \dots \oplus M_{d_k}(\mathbb{C})$ , then  $A$  has precisely  $k$  distinct simple modules  $\{M_1, \dots, M_k\}$  of dimensions  $\{d_1, \dots, d_k\}$ . Here,  $M_i$  can be viewed as column vectors of size  $d_i$  on which the component  $M_{d_i}(\mathbb{C})$  acts by left multiplication and the other factors act as zero. Because  $A$  is semi-simple every  $n$ -dimensional  $A$ -representation  $M$  is isomorphic to

$$M = M_1^{\oplus e_1} \oplus \dots \oplus M_k^{\oplus e_k}$$

for certain multiplicities  $e_i$  satisfying the numerical condition

$$n = e_1 d_1 + \dots + e_k d_k$$

That is,  $\text{rep}_n A$  is the disjoint union of a finite number of (closed) orbits each determined by an integral vector  $(e_1, \dots, e_k)$  satisfying the condition called the *dimension vector* of  $M$ .

$$\text{rep}_n A \simeq \bigsqcup_{(e_1, \dots, e_k)} GL_n / (GL_{e_1} \times \dots \times GL_{e_k})$$

Let  $f_i \geq 1$  be natural numbers such that  $n = f_1 d_1 + \dots + f_k d_k$  and consider the embedding of  $A$  into  $M_n(\mathbb{C})$  defined by

$$(a_1, \dots, a_k) \in A \longrightarrow \left[ \begin{array}{ccc} \underbrace{\begin{bmatrix} a_1 & & 0 \\ & \ddots & \\ 0 & & a_1 \end{bmatrix}}_{f_1} & & \\ & \ddots & \\ & & \underbrace{\begin{bmatrix} a_k & & 0 \\ & \ddots & \\ 0 & & a_k \end{bmatrix}}_{f_k} \end{array} \right] \in M_n(\mathbb{C})$$

Via this embedding,  $A$  becomes a Cayley-Hamilton algebra of degree  $n$  when equipped with the induced trace  $\text{tr}$  from  $M_n(\mathbb{C})$ .

Let  $M$  be the  $n$ -dimensional  $A$ -representation with dimension vector  $(e_1, \dots, e_k)$  and choose a basis compatible with this decomposition. Let  $E_i$  be the idempotent of  $A$  corresponding to the identity matrix  $I_{d_i}$  of the  $i$ -th factor. Then, the trace of the matrix defining the action of  $E_i$  on  $M$  is clearly  $e_i d_i \cdot I_n$ . On the other hand,  $\text{tr}(E_i) = f_i d_i \cdot I_n$ , hence the only trace preserving  $n$ -dimensional  $A$ -representation is that of dimension vector  $(f_1, \dots, f_k)$ . Therefore,  $\text{rep}_n^{\text{tr}} A$  consists of the single closed orbit determined by the integral vector  $(f_1, \dots, f_k)$ .

$$\text{rep}_n^{\text{tr}} A \simeq GL_n / (GL_{f_1} \times \dots \times GL_{f_k})$$

Consider the scheme structure of the trace preserving representation variety  $\underline{rep}_n^{tr} A$ . The corresponding functor assigns to a commutative affine  $\mathbb{C}$ -algebra  $R$

$$\underline{rep}_n^{tr}(R) = \text{Alg}_{\mathbb{C}}(\mathbb{C}[M_n^m]/N_A, R).$$

An algebra morphism  $\psi : \mathbb{C}[M_n^m]/N_A \longrightarrow R$  determines uniquely an  $m$ -tuple of  $n \times n$  matrices with coefficients in  $R$  by

$$r_k = \begin{bmatrix} \psi(x_{11}(k)) & \cdots & \psi(x_{1n}(k)) \\ \vdots & & \vdots \\ \psi(x_{n1}(k)) & \cdots & \psi(x_{nn}(k)) \end{bmatrix}$$

Composing with the canonical embedding

$$\begin{array}{ccc} A & \xrightarrow{\phi} & M_n(R) \\ \downarrow & \nearrow M_n(\psi) & \\ M_n(\mathbb{C}[M_n^m]/N_A) & & \end{array}$$

determines the trace preserving algebra morphism  $\phi : A \longrightarrow M_n(R)$  where the trace map on  $M_n(R)$  is the usual trace. By the universal property any trace preserving map  $A \longrightarrow M_n(R)$  is also of this form.

LEMMA 2.36. *Let  $A$  be a Cayley-Hamilton algebra of degree  $n$  which is generated by  $\{a_1, \dots, a_m\}$ . The trace preserving representation variety  $\underline{rep}_n^{tr} A$  represents the functor*

$$\underline{rep}_n^{tr} A(R) = \{A \xrightarrow{\phi} M_n(R) \mid \phi \text{ is trace preserving}\}$$

Moreover,  $\underline{rep}_n^{tr} A$  is a closed subscheme of  $\underline{rep}_n A$ .

Recall that there is an action of  $GL_n$  on  $\mathbb{C}[M_n^m]$  and from the definition of the ideals  $I_A(n)$  and  $N_A$  it is clear that they are stable under the  $GL_n$ -action. That is, there is an action by automorphisms on the quotient algebras  $\mathbb{C}[M_n^m]/I_A(n)$  and  $\mathbb{C}[M_n^m]/N_A$ . But then, their algebras of invariants are equal to

$$\begin{cases} \mathbb{C}[\underline{rep}_n A]^{GL_n} &= (\mathbb{C}[M_n^m]/I_A(n))^{GL_n} = \frac{\mathbb{N}_n^m}{(I_A(n) \cap \mathbb{N}_n^m)} \\ \mathbb{C}[\underline{rep}_n^{tr} A]^{GL_n} &= (\mathbb{C}[M_n^m]/N_A)^{GL_n} = \frac{\mathbb{N}_n^m}{(N_A \cap \mathbb{N}_n^m)} \end{cases}$$

That is, these rings of invariants define closed subschemes of the affine (reduced) variety associated to the necklace algebra  $\mathbb{N}_n^m$ . We will call these schemes the *quotient schemes* for the action of  $GL_n$  and denote them respectively by

$$\underline{iss}_n A = \underline{rep}_n A/GL_n \quad \text{and} \quad \underline{iss}_n^{tr} A = \underline{rep}_n^{tr} A/GL_n.$$

We have seen that the geometric points of the reduced variety  $\underline{iss}_n A$  of the affine quotient scheme  $\underline{iss}_n A$  parameterize the isomorphism classes of  $n$ -dimensional semisimple  $A$ -representations. Similarly, the geometric points of the reduced variety  $\underline{iss}_n^{tr} A$  of the quotient scheme  $\underline{iss}_n^{tr} A$  parameterize isomorphism classes of *trace preserving*  $n$ -dimensional semisimple  $A$ -representations.

PROPOSITION 2.37. *Let  $A$  be a Cayley-Hamilton algebra of degree  $n$  with trace map  $tr_A$ . Then, we have that*

$$tr_A(A) = \mathbb{C}[\underline{iss}_n^{tr} A],$$

the coordinate ring of the quotient scheme  $\text{iss}_n^{\text{tr}} A$ . In particular, maximal ideals of  $\text{tr}_A(A)$  parameterize the isomorphism classes of trace preserving  $n$ -dimensional semi-simple  $A$ -representations.

By definition, a  $GL_n$ -equivariant map between the affine  $GL_n$ -schemes

$$\underline{\text{rep}}_n^{\text{tr}} A \xrightarrow{f} M_n = \underline{M}_n$$

means that for any commutative affine  $\mathbb{C}$ -algebra  $R$  the corresponding map

$$\underline{\text{rep}}_n^{\text{tr}} A(R) \xrightarrow{f(R)} M_n(R)$$

commutes with the action of  $GL_n(R)$ . Alternatively, the ring of all morphisms  $\underline{\text{rep}}_n^{\text{tr}} A \longrightarrow M_n$  is the matrixalgebra  $M_n(\mathbb{C}[M_n^m]/N_A)$  and those that commute with the  $GL_n$  action are precisely the invariants. That is, we have the following description of  $A$ .

**THEOREM 2.38.** *Let  $A$  be a Cayley-Hamilton algebra of degree  $n$  with trace map  $\text{tr}_A$ . Then, we can recover  $A$  as the ring of  $GL_n$ -equivariant maps*

$$A = \{f : \underline{\text{rep}}_n^{\text{tr}} A \longrightarrow M_n \text{ equivariant} \}$$

of affine  $GL_n$ -schemes.

Summarizing the results of this and the previous section we have

**THEOREM 2.39.** *The functor*

$$\mathbf{alg} @_{\mathbf{n}} \xrightarrow{\text{rep}^t} GL_n - \mathbf{aff}$$

which assigns to a Cayley-Hamilton algebra  $A$  of degree  $n$  the  $GL_n$ -affine scheme  $\underline{\text{rep}}_n^{\text{tr}} A$  of trace preserving  $n$ -dimensional representations has a left inverse. This left inverse functor

$$GL_n - \mathbf{aff} \xrightarrow{\text{wit}} \mathbf{alg} @_{\mathbf{n}}$$

assigns to a  $GL_n$ -affine scheme  $\underline{X}$  its witness algebra  $M_n(\mathbb{C}[\underline{X}])^{GL_n}$  which is a Cayley-Hamilton algebra of degree  $n$ .

Note however that this functor is *not* an equivalence of categories. For, there are many affine  $GL_n$ -schemes having the same witness algebra as we will see in the next section.

We will give an application of the algebraic reconstruction result, theorem 1.28, to finite dimensional algebras. First, we define a *norm map* for algebras in  $\mathbf{alg} @_{\mathbf{n}}$ .

Let  $A$  be a Cayley-Hamilton algebra of degree  $n$  with trace map  $\text{tr}$ , then we can define a *norm map* on  $A$  by defining

$$N(a) = \sigma_n(a) \quad \text{for all } a \in A.$$

Recall that the elementary symmetric function  $\sigma_n$  is a polynomial function  $f(t_1, t_2, \dots, t_n)$  in the Newton functions  $t_i = \sum_{j=1}^n x_j^i$ . Then,  $\sigma(a) = f(\text{tr}(a), \text{tr}(a^2), \dots, \text{tr}(a^n))$ . Because, we have a trace preserving embedding  $A \hookrightarrow M_n(\mathbb{C}[\underline{\text{rep}}_n^{\text{tr}} A])$  and the norm map  $N$  coincides with the determinant in this matrix-algebra, we have that

$$N(1) = 1 \quad \text{and} \quad \forall a, b \in A \quad N(ab) = N(a)N(b).$$

Furthermore, the norm map extends to a polynomial map on  $A[t]$  and we have that  $\chi_a^{(n)}(t) = N(t - a)$ , in particular we can obtain the trace by polarization of the norm map. Consider a finite dimensional semi-simple  $\mathbb{C}$ -algebra

$$A = M_{d_1}(\mathbb{C}) \oplus \dots \oplus M_{d_k}(\mathbb{C}),$$

then all the Cayley-Hamilton structures of degree  $n$  on  $A$  with trace values in  $\mathbb{C}$  are given by the following result.

LEMMA 2.40. *Let  $A$  be a semi-simple algebra as above and  $tr$  a trace map on  $A$  making it into a Cayley-Hamilton algebra of degree  $n$  with  $tr(A) = \mathbb{C}$ . Then, there exist a dimension vector  $\alpha = (m_1, \dots, m_k) \in \mathbb{N}_+^k$  such that  $n = \sum_{i=1}^k m_i d_i$  and for any  $a = (A_1, \dots, A_k) \in A$  with  $A_i \in M_{d_i}(\mathbb{C})$  we have that*

$$tr(a) = m_1 Tr(A_1) + \dots + m_k Tr(A_k)$$

where  $Tr$  are the usual trace maps on matrices.

PROOF. The norm-map  $N$  on  $A$  defined by the trace map  $tr$  induces a group morphism on the invertible elements of  $A$

$$N : A^* = GL_{d_1}(\mathbb{C}) \times \dots \times GL_{d_k}(\mathbb{C}) \longrightarrow \mathbb{C}^*$$

that is, a *character*. Now, any character is of the following form, let  $A_i \in GL_{d_i}(\mathbb{C})$ , then for  $a = (A_1, \dots, A_k)$  we must have

$$N(a) = \det(A_1)^{m_1} \det(A_2)^{m_2} \dots \det(A_k)^{m_k}$$

for certain integers  $m_i \in \mathbb{Z}$ . Since  $N$  extends to a polynomial map on the whole of  $A$  we must have that all  $m_i \geq 0$ . By polarization it then follows that

$$tr(a) = m_1 Tr(A_1) + \dots + m_k Tr(A_k)$$

and it remains to show that no  $m_i = 0$ . Indeed, if  $m_i = 0$  then  $tr$  would be the zero map on  $M_{d_i}(\mathbb{C})$ , but then we would have for any  $a = (0, \dots, 0, A, 0, \dots, 0)$  with  $A \in M_{d_i}(\mathbb{C})$  that

$$\chi_a^{(n)}(t) = t^n$$

whence  $\chi_a^{(n)}(a) \neq 0$  whenever  $A$  is not nilpotent. This contradiction finishes the proof.  $\square$

We can extend this to all finite dimensional  $\mathbb{C}$ -algebras. Let  $A$  be a finite dimensional algebra with radical  $J$  and assume there is a trace map  $tr$  on  $A$  making  $A$  into a Cayley-Hamilton algebra of degree  $n$  and such that  $tr(A) = \mathbb{C}$ . We claim that the norm map  $N : A \longrightarrow \mathbb{C}$  is zero on  $J$ . Indeed, any  $j \in J$  satisfies  $j^l = 0$  for some  $l$  whence  $N(j)^l = 0$ . But then, polarization gives that  $tr(j) = 0$  and we have that the semisimple algebra

$$A^{ss} = A/J = M_{d_1}(\mathbb{C}) \oplus \dots \oplus M_{d_k}(\mathbb{C})$$

is a semisimple Cayley-hamilton algebra of degree  $n$  on which we can apply the foregoing lemma. Finally, note that  $A \simeq A^{ss} \oplus J$  as  $\mathbb{C}$ -vectorspaces. This concludes the proof of

PROPOSITION 2.41. *Let  $A$  be a finite dimensional  $\mathbb{C}$ -algebra with radical  $J$  and semisimple part*

$$A^{ss} = A/J = M_{d_1}(\mathbb{C}) \oplus \dots \oplus M_{d_k}(\mathbb{C}).$$



If  $tr : A \longrightarrow \mathbb{C} \hookrightarrow A$  is a trace map such that  $A$  is a Cayley-Hamilton algebra of degree  $n$ , there exists a dimension vector  $\alpha = (m_1, \dots, m_k) \in \mathbb{N}_+^k$  such that for all  $a = (A_1, \dots, A_k, j)$  with  $A_i \in M_{d_i}(\mathbb{C})$  and  $j \in J$  we have

$$tr(a) = m_1 Tr(A_1) + \dots + m_k Tr(A_k)$$

with  $Tr$  the usual traces on  $M_{d_i}(\mathbb{C})$  and  $\sum_i m_i d_i = n$ .

Fix a trace map  $tr$  on  $A$  determined by a dimension vector  $\alpha = (m_1, \dots, m_k) \in \mathbb{N}^k$ . Then, the trace preserving variety  $rep_n^{tr} A$  is the scheme of  $A$ -modules of dimension vector  $\alpha$ , that is, those  $A$ -modules  $M$  such that

$$M^{ss} = S_1^{\oplus m_1} \oplus \dots \oplus S_k^{\oplus m_k}$$

where  $S_i$  is the simple  $A$ -module of dimension  $d_i$  determined by the  $i$ -th factor in  $A^{ss}$ . An immediate consequence of the reconstruction theorem 2.38 is

**PROPOSITION 2.42.** *Let  $A$  be a finite dimensional algebra with trace map  $tr : A \longrightarrow \mathbb{C}$  determined by a dimension vector  $\alpha = (m_1, \dots, m_k)$  as before with all  $m_i > 0$ . Then,  $A$  can be recovered from the  $GL_n$ -structure of the affine scheme  $rep_n^{tr} A$  of all  $A$ -modules of dimension vector  $\alpha$ .*

Still, there can be other trace maps on  $A$  making  $A$  into a Cayley-Hamilton algebra of degree  $n$ . For example let  $C$  be a finite dimensional commutative  $\mathbb{C}$ -algebra with radical  $N$ , then  $A = M_n(C)$  is finite dimensional with radical  $J = M_n(N)$  and the usual trace map  $tr : M_n(C) \longrightarrow C$  makes  $A$  into a Cayley-Hamilton algebra of degree  $n$  such that  $tr(J) = N \neq 0$ . Still, if  $A$  is semi-simple, the center  $Z(A) = \mathbb{C} \oplus \dots \oplus \mathbb{C}$  (as many terms as there are matrix components in  $A$ ) and any subring of  $Z(A)$  is of the form  $\mathbb{C} \oplus \dots \oplus \mathbb{C}$ . In particular,  $tr(A)$  has this form and composing the trace map with projection on the  $j$ -th component we have a trace map  $tr_j$  on which we can apply lemma 2.40.

### 2.7. The Gerstenhaber-Hesselink theorem.


In this section we will give examples of distinct  $GL_n$ -affine schemes having the same witness algebra, proving that the left inverse of theorem 2.39 is *not* an equivalence of categories. We will study the orbits in  $rep_n \mathbb{C}[x]$  or, equivalent, conjugacy classes of  $n \times n$  matrices.

It is sometimes convenient to relax our definition of partitions to include zeroes at its tail. That is, a partition  $p$  of  $n$  is an integral  $n$ -tuple  $(a_1, a_2, \dots, a_n)$  with  $a_1 \geq a_2 \geq \dots \geq a_n \geq 0$  with  $\sum_{i=1}^n a_i = n$ . As before, we represent a partition by a Young diagram by omitting rows corresponding to zeroes.

If  $q = (b_1, \dots, b_n)$  is another partition of  $n$  we say that  $p$  *dominates*  $q$  and write

$$p > q \quad \text{if and only if} \quad \sum_{i=1}^r a_i \geq \sum_{i=1}^r b_i \quad \text{for all } 1 \leq r \leq n.$$


For example, the partitions of 4 are ordered as indicated below




Note however that the dominance relation is *not* a total ordering whenever  $n \geq 6$ . For example, the following two partition of 6



are not comparable. The *dominance order* is induced by the *Young move* of throwing a row-ending box down the diagram. Indeed, let  $p$  and  $q$  be partitions of  $n$  such that  $p > q$  and assume there is no partition  $r$  such that  $p > r$  and  $r > q$ . Let  $i$  be the minimal number such that  $a_i > b_i$ . Then by the assumption  $a_i = b_i + 1$ . Let  $j > i$  be minimal such that  $a_j \neq b_j$ , then we have  $b_j = a_j + 1$  because  $p$  dominates  $q$ . But then, the remaining rows of  $p$  and  $q$  must be equal. That is, a Young move can be depicted as



For example, the Young moves between the partitions of 4 given above are as indicated



A *Young  $p$ -tableau* is the Young diagram of  $p$  with the boxes labeled by integers from  $\{1, 2, \dots, s\}$  for some  $s$  such that each label appears at least ones. A Young  $p$ -tableau is said to be *of type  $q$*  for some partition  $q = (b_1, \dots, b_n)$  of  $n$  if the following conditions are met :

- the labels are non-decreasing along rows,
- the labels are strictly increasing along columns, and
- the label  $i$  appears exactly  $b_i$  times.

For example, if  $p = (3, 2, 1, 1)$  and  $q = (2, 2, 2, 1)$  then the  $p$ -tableau below

1	1	3
2	2	
3		
4		

is of type  $q$  (observe that  $p > q$  and even  $p \rightarrow q$ ). In general, let  $p = (a_1, \dots, a_n)$  and  $q = (b_1, \dots, b_n)$  be partitions of  $n$  and assume that  $p \rightarrow q$ . Then, there is a Young  $p$ -tableau of type  $q$ . For, fill the Young diagram of  $q$  by putting 1's in the first row, 2's in the second and so on. Then, upgrade the fallen box together with



Now label the boxes of this tableau by the base vectors  $\{v_1, \dots, v_n\}$  such that the boxes labeled  $i$  in the Young  $q^*$ -tableau of type  $p^*$  are filled with the base vectors from  $V_i - V_{i-1}$ . Call this tableau  $T$ . In the example, we can take

$$T = \begin{array}{|c|c|c|c|} \hline e_1 & e_4 & e_6 & e_7 \\ \hline e_2 & e_5 & e_3 & \\ \hline \end{array}$$

Define a linear operator  $F$  on  $\mathbb{C}^n$  by the rule that  $F(v_i) = v_j$  if  $v_j$  is the label of the box in  $T$  just above the box labeled  $v_i$ . In case  $v_i$  is a label of a box in the first row of  $T$  we take  $F(v_i) = 0$ . Obviously,  $F$  is a nilpotent  $n \times n$  matrix and by construction we have that

$$rk F^l = n - (b_1^* + \dots + b_l^*)$$

That is,  $F$  is nilpotent of type  $q = (b_1, \dots, b_n)$ . Moreover,  $F$  satisfies  $F(V_i) \subset V_{i-1}$  for all  $i$  by the way we have labeled the tableau  $T$  and defined  $F$ .

In the example above, we have  $F(e_2) = e_1$ ,  $F(e_5) = e_4$ ,  $F(e_3) = e_6$  and all other  $F(e_i) = 0$ . That is,  $F$  is the matrix

$$\begin{bmatrix} 0 & 1 & & & & & & \\ 0 & 0 & & & & & & \\ & & 0 & & & & 0 & \\ & & & 0 & 1 & & & \\ & & & 0 & 0 & & & \\ & & 1 & & & & 0 & \\ & & & & & & & 0 \end{bmatrix}$$

which is seen to be of type  $(2, 2, 2, 1)$  after performing a few Jordan moves.

Returning to the general case, consider for all  $\epsilon \in \mathbb{C}$  the  $n \times n$  matrix :

$$F_\epsilon = (1 - \epsilon)F + \epsilon A.$$

We claim that for all but finitely many values of  $\epsilon$  we have  $F_\epsilon \in \mathcal{O}(A)$ . Indeed, we have seen that  $F(V_i) \subset V_{i-1}$  where  $V_i$  is defined as the subspace such that  $A^i(V_i) = 0$ . Hence,  $F(V_1) = 0$  and therefore

$$F_\epsilon(V_1) = (1 - \epsilon)F + \epsilon A(V_1) = 0.$$

Assume by induction that  $F_\epsilon^i(V_i) = 0$  holds for all  $i < l$ , then we have that

$$\begin{aligned} F_\epsilon^l(V_l) &= F_\epsilon^{l-1}((1 - \epsilon)F + \epsilon A)(V_l) \\ &\subset F_\epsilon^{l-1}(V_{l-1}) = 0 \end{aligned}$$

because  $A(V_l) \subset V_{l-1}$  and  $F(V_l) \subset V_{l-1}$ . But then we have for all  $l$  that

$$rk F_\epsilon^l \leq \dim V_l = n - (a_1^* + \dots + a_l^*) = rk A^l \stackrel{def}{=} r_l.$$

Then for at least one  $r_l \times r_l$  submatrix of  $F_\epsilon^l$  its determinant considered it as a polynomial of degree  $r_l$  in  $\epsilon$  is not identically zero (as it is nonzero for  $\epsilon = 1$ ). But then this determinant is non-zero for all but finitely many  $\epsilon$ . Hence,  $rk F_\epsilon^l = rk A^l$  for all  $l$  for all but finitely many  $\epsilon$ . As these numbers determine the dual partition  $p^*$  of the type of  $A$ ,  $F_\epsilon$  is a nilpotent  $n \times n$  matrix of type  $p$  for all but finitely many values of  $\epsilon$ , proving the claim. But then,  $F_0 = F$  which we have proved to

be a nilpotent matrix of type  $q$  belongs to the closure of the orbit  $\mathcal{O}(A)$ . That is, we have proved the difficult part of the *Gerstenhaber-Hesselink theorem*.

**THEOREM 2.43.** *Let  $A$  be a nilpotent  $n \times n$  matrix of type  $p = (a_1, \dots, a_n)$  and  $B$  nilpotent of type  $q = (b_1, \dots, b_n)$ . Then,  $B$  belongs to the closure of the orbit  $\mathcal{O}(A)$ , that is,*

$$B \in \overline{\mathcal{O}(A)} \quad \text{if and only if} \quad p > q$$

*in the domination order on partitions of  $n$ .*

To prove the theorem we only have to observe that if  $B$  is contained in the closure of  $A$ , then  $B^l$  is contained in the closure of  $A^l$  and hence  $rk A^l \geq rk B^l$  (because  $rk A^l < k$  is equivalent to vanishing of all determinants of  $k \times k$  minors which is a closed condition). But then,

$$n - \sum_{i=1}^l a_i^* \geq n - \sum_{i=1}^l b_i^*$$

for all  $l$ , that is,  $q^* > p^*$  and hence  $p > q$ . The other implication was proved above if we remember that the domination order was induced by the Young moves and clearly we have that if  $B \in \overline{\mathcal{O}(C)}$  and  $C \in \overline{\mathcal{O}(A)}$  then also  $B \in \overline{\mathcal{O}(A)}$ .


**EXAMPLE 2.44.** Nilpotent matrices for small  $n$ .

We will apply theorem 2.43 to describe the orbit-closures of nilpotent matrices of  $8 \times 8$  matrices. The following table lists all partitions (and their dual in the other column)

The partitions of 8.


a	(8)	v	(1,1,1,1,1,1,1,1)
b	(7,1)	u	(2,1,1,1,1,1,1)
c	(6,2)	t	(2,2,1,1,1,1)
d	(6,1,1)	s	(3,1,1,1,1,1)
e	(5,3)	r	(2,2,2,1,1)
f	(5,2,1)	q	(3,2,1,1,1)
g	(5,1,1,1)	p	(4,1,1,1,1)
h	(4,4)	o	(2,2,2,2)
i	(4,3,1)	n	(3,2,2,1)
j	(4,2,2)	m	(3,3,1,1)
k	(3,3,2)	k	(3,3,2)
l	(4,2,1,1)	l	(4,2,1,1)

The domination order between these partitions can be depicted as follows where all the Young moves are from left to right




Of course, from this graph we can read off the dominance order graphs for partitions of  $n \leq 8$ . The trick is to identify a partition of  $n$  with that of 8 by throwing in a tail of ones and to look at the relative position of both partitions in the above

picture. Using these conventions we get the following graph for partitions of 7



and for partitions of 6 the dominance order is depicted as follows



The dominance order on partitions of  $n \leq 5$  is a total ordering.

The Gerstenhaber-Hesselink theorem can be applied to describe the module varieties of the algebras  $\frac{\mathbb{C}[x]}{(x^r)}$ .

EXAMPLE 2.45. The representation variety  $rep_n \frac{\mathbb{C}[x]}{(x^r)}$ .

Any algebra morphism  $\mathbb{C}[x] \rightarrow M_n$  is determined by the image of  $x$ , whence  $rep_n(\mathbb{C}[x]) = M_n$ . We have seen that conjugacy classes in  $M_n$  are classified by the Jordan normalform. Let  $A$  is conjugated to a matrix in normalform

$$\begin{bmatrix} J_1 & & & \\ & J_2 & & \\ & & \ddots & \\ & & & J_s \end{bmatrix}$$

where  $J_i$  is a Jordan block of size  $d_i$ , hence  $n = d_1 + d_2 + \dots + d_s$ . Then, the  $n$ -dimensional  $\mathbb{C}[x]$ -module  $M$  determined by  $A$  can be decomposed uniquely as

$$M = M_1 \oplus M_2 \oplus \dots \oplus M_s$$

where  $M_i$  is a  $\mathbb{C}[x]$ -module of dimension  $d_i$  which is *indecomposable*, that is, cannot be decomposed as a direct sum of proper submodules.

Now, consider the quotient algebra  $R = \mathbb{C}[x]/(x^r)$ , then the ideal  $I_R(n)$  of  $\mathbb{C}[x_{11}, x_{12}, \dots, x_{nn}]$  is generated by the  $n^2$  entries of the matrix

$$\begin{bmatrix} x_{11} & \dots & x_{1n} \\ \vdots & & \vdots \\ x_{n1} & \dots & x_{nn} \end{bmatrix}^r.$$

For example if  $r = m = 2$ , then the ideal is generated by the entries of the matrix

$$\begin{bmatrix} x_1 & x_2 \\ x_3 & x_4 \end{bmatrix}^2 = \begin{bmatrix} x_1^2 + x_2x_3 & x_2(x_1 + x_4) \\ x_3(x_1 + x_4) & x_4^2 + x_2x_3 \end{bmatrix}$$

That is, the ideal with generators

$$I_R = (x_1^2 + x_2x_3, x_2(x_1 + x_4), x_3(x_1 + x_4), (x_1 - x_4)(x_1 + x_4))$$

The variety  $\mathbb{V}(I_R) \hookrightarrow M_2$  consists of all matrices  $A$  such that  $A^2 = 0$ . Conjugating  $A$  to an upper triangular form we see that the eigenvalues of  $A$  must be zero,

hence

$$\text{rep}_2 \mathbb{C}[x]/(x^2) = \mathcal{O}\left(\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}\right) \cup \mathcal{O}\left(\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}\right)$$

and we have seen that this variety is a cone with top the zero matrix and defining equations

$$\mathbb{V}(x_1 + x_4, x_1^2 + x_2x_3)$$

and we see that  $I_R$  is properly contained in this ideal. Still, we have that

$$\text{rad}(I_R) = (x_1 + x_4, x_1^2 + x_3x_4)$$

for an easy computation shows that  $\overline{x_1 + x_4}^3 = 0 \in \mathbb{C}[x_1, x_2, x_3, x_4]/I_R$ . Therefore, even in the easiest of examples, the representation variety does not have to be reduced.

For the general case, observe that when  $J$  is a Jordan block of size  $d$  with eigenvalue zero an easy calculation shows that

$$J^{d-1} = \begin{bmatrix} 0 & \dots & 0 & d-1 \\ & \ddots & & 0 \\ & & \ddots & \vdots \\ & & & 0 \end{bmatrix} \quad \text{and} \quad J^d = \begin{bmatrix} 0 & \dots & \dots & 0 \\ \vdots & & & \vdots \\ \vdots & & & \vdots \\ 0 & \dots & \dots & 0 \end{bmatrix}$$

Therefore, we see that the representation variety  $\text{rep}_n \mathbb{C}[x]/(x^r)$  is the union of all conjugacy classes of matrices having 0 as only eigenvalue and all of which Jordan blocks have size  $\leq r$ . Expressed in module theoretic terms, any  $n$ -dimensional  $R = \mathbb{C}[x]/(x^r)$ -module  $M$  is isomorphic to a direct sum of indecomposables

$$M = I_1^{\oplus e_1} \oplus I_2^{\oplus e_2} \oplus \dots \oplus I_r^{\oplus e_r}$$

where  $I_j$  is the unique indecomposable  $j$ -dimensional  $R$ -module (corresponding to the Jordan block of size  $j$ ). Of course, the multiplicities  $e_i$  of the factors must satisfy the equation

$$e_1 + 2e_2 + 3e_3 + \dots + re_r = n$$

In  $M$  we can consider the subspaces for all  $1 \leq i \leq r-1$

$$M_i = \{m \in M \mid x^i.m = 0\}$$

the dimension of which can be computed knowing the powers of Jordan blocks (observe that the dimension of  $M_i$  is equal to  $n - \text{rank}(A^i)$ )

$$t_i = \dim_{\mathbb{C}} M_i = e_1 + 2e_2 + \dots + (i-1)e_i + i(e_i + e_{i+1} + \dots + e_r)$$

Observe that giving  $n$  and the  $r-1$ -tuple  $(t_1, t_2, \dots, t_{r-1})$  is the same as giving the multiplicities  $e_i$  because

$$\begin{cases} 2t_1 & = t_2 + e_1 \\ 2t_2 & = t_3 + t_1 + e_2 \\ 2t_3 & = t_4 + t_2 + e_3 \\ & \vdots \\ 2t_{n-2} & = t_{n-1} + t_{n-3} + e_{n-2} \\ 2t_{n-1} & = n + t_{n-2} + e_{n-1} \\ n & = t_{n-1} + e_n \end{cases}$$

Let  $n$ -dimensional  $\mathbb{C}[x]/(x^r)$ -modules  $M$  and  $M'$  (or associated matrices  $A$  and  $A'$ ) be determined by the  $r - 1$ -tuples  $(t_1, \dots, t_{r-1})$  respectively  $(t'_1, \dots, t'_{r-1})$  then we have that

$$\mathcal{O}(A') \hookrightarrow \overline{\mathcal{O}(A)} \quad \text{if and only if} \quad t_1 \leq t'_1, t_2 \leq t'_2, \dots, t_{r-1} \leq t'_{r-1}$$

Therefore, we have an inverse order isomorphism between the orbits in  $\text{rep}_n(\mathbb{C}[x]/(x^r))$  and the  $r - 1$ -tuples of natural numbers  $(t_1, \dots, t_{r-1})$  satisfying the following linear inequalities (which follow from the above system)

$$2t_1 \geq t_2, 2t_2 \geq t_3 + t_1, 2t_3 \geq t_4 + t_2, \dots, 2t_{n-1} \geq n + t_{n-2}, n \geq t_{n-2}.$$

Let us apply this general result in a few easy cases. First, consider  $r = 2$ , then the orbits in  $\text{rep}_n \mathbb{C}[x]/(x^2)$  are parameterized by a natural number  $t_1$  satisfying the inequalities  $n \geq t_1$  and  $2t_1 \geq n$ , the multiplicities are given by  $e_1 = 2t_1 - n$  and  $e_2 = n - t_1$ . Moreover, the orbit of the module  $M(t'_1)$  lies in the closure of the orbit of  $M(t_1)$  whenever  $t_1 \leq t'_1$ .


That is, if  $n = 2k + \delta$  with  $\delta = 0$  or  $1$ , then  $\text{rep}_n \mathbb{C}[x]/(x^2)$  is the union of  $k + 1$  orbits and the orbitclosures form a linear order as follows (from big to small)

$$I_1^\delta \oplus I_2^{\oplus k} \text{ --- } I_1^{\oplus \delta+2} \oplus I_2^{\oplus k-1} \text{ --- } \dots \text{ --- } I_1^{\oplus n}$$

If  $r = 3$ , orbits in  $\text{rep}_n \mathbb{C}[x]/(x^3)$  are determined by couples of natural numbers  $(t_1, t_2)$  satisfying the following three linear inequalities


$$\begin{cases} 2t_1 & \geq t_2 \\ 2t_2 & \geq n + t_1 \\ n & \geq t_2 \end{cases}$$

For example, for  $n = 8$  we obtain the following situation





Therefore,  $rep_8 \mathbb{C}[x]/(x^3)$  consists of 10 orbits with orbitclosure diagram as below (the nodes represent the multiplicities  $[e_1 e_2 e_3]$ ).



Here we used the equalities  $e_1 = 2t_1 - t_2$ ,  $e_2 = 2t_2 - n - t_1$  and  $e_3 = n - t_2$ . For general  $n$  and  $r$  this result shows that  $rep_n \mathbb{C}[x]/(x^r)$  is the closure of the orbit of the module with decomposition

$$M_{gen} = I_r^{\oplus e} \oplus I_s \quad \text{if} \quad n = er + s$$

We are now in a position to give the promised examples of affine  $GL_n$ -schemes having the same witness algebra.

EXAMPLE 2.46. Consider the action of  $GL_n$  on  $M_n$  by conjugation and take a nilpotent matrix  $A$ . All eigenvalues of  $A$  are zero, so the conjugacy class of  $A$  is fully determined by the sizes of its Jordan blocks. These sizes determine a partition  $\lambda(A) = (\lambda_1, \lambda_2, \dots, \lambda_k)$  of  $n$  with  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_k$ . Moreover, we have given an algorithm to determine whether an orbit  $\mathcal{O}(B)$  of another nilpotent matrix  $B$  is contained in the orbit closure  $\overline{\mathcal{O}(A)}$ , the criterium being that


$$\mathcal{O}(B) \subset \overline{\mathcal{O}(A)} \iff \lambda(B)^* \geq \lambda(A)^*.$$

where  $\lambda^*$  denotes the dual partition. We see that the witness algebra of  $\overline{\mathcal{O}(A)}$  is equal to

$$M_n(\mathbb{C}[\overline{\mathcal{O}(A)}])^{GL_n} = \mathbb{C}[X]/(X^k)$$


where  $k$  is the number of columns of the Young diagram  $\lambda(A)$ .

Hence, the orbit closures of nilpotent matrices such that their associated Young diagrams have equal number of columns have the same witness algebras. For example, if  $n = 4$  then the closures of the orbits corresponding to



have the same witness algebra, although the closure of the second is a proper closed subscheme of the closure of the first.

Recall the orbitclosure diagram of conjugacy classes of nilpotent  $8 \times 8$  matrices given by the Gerstenhaber-Hesslink theorem. In the picture below, the closures of orbits corresponding to connected nodes of the same colour have the same witness algebra.



### 2.8. The real moment map.

In this section we will give another interpretation of the algebraic quotient variety  $\underline{iss}_n^{tr} A$  with methods coming from symplectic geometry. We have an involution

$$GL_n \xrightarrow{i} GL_n \quad \text{defined by} \quad g \longrightarrow (g^*)^{-1}$$

where  $A^*$  is the *adjoint matrix* of  $g$ , that is, the conjugate transpose

$$M = \begin{bmatrix} m_{11} & \dots & m_{1n} \\ \vdots & & \vdots \\ m_{n1} & \dots & m_{nn} \end{bmatrix} \quad M^* = \begin{bmatrix} \overline{m_{11}} & \dots & \overline{m_{n1}} \\ \vdots & & \vdots \\ \overline{m_{1n}} & \dots & \overline{m_{nn}} \end{bmatrix}$$

The *real points* of this involution, that is

$$(GL_n)^i = \{g \in GL_n \mid g = (g^*)^{-1}\} = U_n = \{u \in GL_n \mid uu^* = \mathbb{1}_n\}$$

is the *unitary group*. On the level of Lie algebras, the involution  $i$  gives rise to the linear map

$$M_n \xrightarrow{di} M_n \quad \text{defined by} \quad M \longrightarrow -M^*$$

corresponding to the fact that the Lie algebra of the unitary group, that is, the kernel of  $di$ , is the space of *skew-Hermitian matrices*

$$\text{Lie } U_n = \{M \in M_n \mid M = -M^*\} = i\text{Herm}_n$$

Consider the standard *Hermitian inproduct* on  $M_n$  defined by

$$(A, B) = \text{tr}(A^*B) \quad \text{which satisfies} \quad \begin{cases} (cA, B) = \overline{c}(A, B) \\ (A, cB) = c(A, B) \\ (B, A) = \overline{(A, B)} \end{cases}$$

As a subgroup of  $GL_n$ ,  $U_n$  acts on  $M_n$  by conjugation and because  $(uAu^*, uBu^*) = \text{tr}(uA^*u^*uBu^*) = \text{tr}(A^*B)$ , the inproduct is invariant under the  $U_n$ -action. The action of  $U_n$  on  $M_n$  induces an action of  $\text{Lie } U_n$  on  $M_n$  given for all  $h \in \text{Lie } U_n$  and  $M \in M_n$

$$h.M = hM + Mh^* = hM - Mh$$

Using this action, we define the *real moment map*  $\mu$  for the action of  $U_n$  on  $M_n$  as the map from  $M_n$  to the linear dual of the Lie algebra

$$M_n \xrightarrow{\mu} (i\text{Lie } U_n)^* \quad M \longrightarrow (h \mapsto i(h.M, M))$$

We will identify the inverse image of the zero map  $\underline{0} : Lie U_n \longrightarrow 0$  under  $\mu$ . Because

$$\begin{aligned} (h.M, M) &= tr((h.M - M.h)^*M) \\ &= tr(M^*h^*M - h^*M^*M) \\ &= tr(h^*(MM^* - M^*M)) \end{aligned}$$

and using the nondegeneracy of the *Killing form* on *Lie U<sub>n</sub>* we have the identification

$$\mu^{-1}(\underline{0}) = \{M \in M_n \mid MM^* = M^*M\} = Nor_n$$

the space of *normal matrices*. Alternatively, we can define the real moment map to be determined by

$$M_n \xrightarrow{\mu_{\mathbb{R}}} Lie U_n \quad M \longrightarrow i(MM^* - M^*M) = i[M, M^*]$$

Recall that a matrix  $M \in M_n(\mathbb{C})$  is said to be *normal* if it commutes with its adjoint. For example, diagonal matrices are normal as are unitary matrices. Further, it is clear that if  $M$  is normal and  $u$  unitary, then the conjugated matrix  $uMu^{-1} = uMu^*$  is again a normal matrix, that is we have an action of the compact Lie group  $U_n$  on the subset  $Nor_n \hookrightarrow M_n(\mathbb{C})$  of normal matrices. We recall the proof of the following classical result

**THEOREM 2.47.** *Every  $U_n$  orbit in  $Nor_n$  contains a diagonal matrix. This gives a natural one-to-one correspondence*

$$\mu^{-1}(\underline{0})/U_n = Nor_n/U_n \longleftrightarrow M_n/GL_n$$

*between the  $U_n$ -orbits in  $Nor_n$  and the closed  $GL_n$ -orbits in  $M_n$ .*

**PROOF.** Equip  $\mathbb{C}^n$  with the standard Hermitian form, that is,

$$\langle v, w \rangle = \bar{v}^T \cdot w = \bar{v}_1 w_1 + \dots + \bar{v}_n w_n$$

Take a non-zero eigenvector  $v$  of  $M \in Nor_n$  and normalize it such that  $\langle v, v \rangle = 1$ . Extend  $v = v_1$  to an orthonormal basis  $\{v_1, \dots, v_n\}$  of  $\mathbb{C}^n$  and let  $u$  be the basechange matrix from the standard basis. With respect to the new basis, the linear map determined by  $M$  and  $M^*$  are represented by the normal matrices

$$M_1 = uMu^* = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ 0 & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & & \vdots \\ 0 & a_{n2} & \dots & a_{nn} \end{bmatrix} \quad M_1^* = uM^*u^* = \begin{bmatrix} \bar{a}_{11} & 0 & \dots & 0 \\ \bar{a}_{12} & \bar{a}_{22} & \dots & \bar{a}_{n2} \\ \vdots & \vdots & & \vdots \\ \bar{a}_{1n} & \bar{a}_{2n} & \dots & \bar{a}_{nn} \end{bmatrix}$$

Because  $M$  is normal, so is  $M_1$ . The left hand corner of  $M_1^*M_1$  is  $a_{11}\bar{a}_{11}$  whereas that of  $M_1M_1^*$  is  $a_{11}\bar{a}_{11} + a_{12}\bar{a}_{12} + \dots + a_{1n}\bar{a}_{1n}$ , whence

$$a_{12}\bar{a}_{12} + \dots + a_{1n}\bar{a}_{1n} = 0$$

but as all  $a_{1i}\bar{a}_{1i} = \|a_{1i}\|^2 \geq 0$ , this implies that all  $a_{1i} = 0$ , whence

$$M_1 = \begin{bmatrix} a_{11} & 0 & \dots & 0 \\ 0 & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & & \vdots \\ 0 & a_{n2} & \dots & a_{nn} \end{bmatrix}$$

and induction finishes the claim. Because permutation matrices are unitary we see that the diagonal entries are determined up to permutation, so every  $U_n$ -orbit

determines a unique conjugacy class of semi-simple matrices, that is, a closed  $GL_n$ -orbit in  $M_n$ .  $\square$

We will generalize this classical result to  $m$ -tuples of  $n \times n$  matrices,  $M_n^m$ , and then by restriction to trace preserving representation varieties. Take  $A = (A_1, \dots, A_m)$  and  $B = (B_1, \dots, B_m)$  in  $M_n^m$  and define an Hermitian inproduct on  $M_n^m$  by

$$(A, B) = \text{tr}(A_1^* B_1 + \dots + A_m^* B_m)$$

which is again invariant under the action of  $U_n$  by simultaneous conjugation on  $M_n^m$ . The induced action of  $Lie U_n$  on  $M_n^m$  is given by

$$h.A = (hA_1 - A_1 h, \dots, hA_m - A_m h)$$

This allows us to define the *real moment map*  $\mu$  for the action of  $U_n$  on  $M_n^m$  to be the assignment

$$M_n^m \xrightarrow{\mu} (iLie U_n)^* \quad A \longrightarrow (h \mapsto i(h.A, A))$$

and again using the nondegeneracy of the Killing form on  $Lie U_n$  we have the identification

$$\mu^{-1}(0) = \{A \in M_n^m \mid \sum_{i=1}^m (A_i A_i^* - A_i^* A_i) = 0\}$$

Again, the real moment map is determined by

$$M_n^m \xrightarrow{\mu_{\mathbb{R}}} Lie U_n \quad A = (A_1, \dots, A_m) \mapsto i[A, A^*] = i \sum_{j=1}^m [A_j, A_j^*]$$

We will show that there is a natural one-to-one correspondence between  $U_n$ -orbits in the set  $\mu^{-1}(0)$  and closed  $GL_n$ -orbits in  $M_n^m$ . We first consider the properties of the real valued function  $p_A$  defined as the norm on the orbit of any  $A \in M_n^m$

$$GL_n \xrightarrow{p_A} \mathbb{R}_+ \quad g \longrightarrow \|g.A\|^2$$

Because the Hermitian inproduct is invariant under  $U_n$  we have  $p_A(ug) = p_A(g)$  for any  $u \in U_n$ . If  $Stab(A)$  denotes the stabilizer subgroup of  $A \in GL_n$ , then for any  $s \in Stab(A)$  we also have  $p_A(gs) = p_A(g)$  hence  $p_A$  is constant along  $U_n g Stab(A)$ -cosets. We aim to prove that the critical points of  $p_A$  are minima and that the minimum is attained if and only if  $\mathcal{O}(A)$  is a closed  $GL_n$ -orbit.

Consider the restriction of  $p_A$  to the maximal torus  $T_n \hookrightarrow GL_n$  of invertible diagonal matrices. Then,  $T_n \cap U_n = K = U_1 \times \dots \times U_1$  is the subgroup

$$K = \left\{ \begin{bmatrix} k_1 & & 0 \\ & \ddots & \\ 0 & & k_n \end{bmatrix} \quad \text{where } |k_i| = 1 \right\}$$

The action by conjugation of  $T_n$  on  $M_n^m$  decomposes this space into *weight spaces*

$$M_n^m = M_n^m(0) \oplus \bigoplus_{i,j=1}^n M_n^m(\pi_i - \pi_j)$$

where  $M_n^m(\pi_i - \pi_j) = \{A \in M_n^m \mid \text{diag}(t_1, \dots, t_n).A = t_i t_j^{-1} A\}$ . It follows from the definition of the Hermitian inproduct on  $M_n^m$  that the different weightspaces

are orthogonal to each other. We decompose  $A \in M_n^m$  into eigenvectors for the  $T_n$ -action as

$$A = A(0) + \sum_{i,j=1}^n A(i,j) \quad \text{with} \quad \begin{cases} A(0) & \in M_n^m(0) \\ A(i,j) & \in M_n^m(\pi_i - \pi_j) \end{cases}$$

With this convention we have for  $t = \text{diag}(t_1, \dots, t_n) \in T_n$  that

$$\begin{aligned} p_A(t) &= \|A(0) + \sum_{i,j=1}^n t_i t_j^{-1} A(i,j)\|^2 \\ &= \|A(0)\|^2 + \sum_{i,j=1}^n t_i^2 t_j^{-2} \|A(i,j)\|^2 \end{aligned}$$

where the last equality follows from the orthogonality of the different weight spaces. Further, remark that the stabilizer subgroup  $\text{Stab}_T(A)$  of  $A$  in  $T$  can be identified with

$$\text{Stab}_T(A) = \{t = \text{diag}(t_1, \dots, t_n) \mid t_i = t_j \text{ if } A(i,j) \neq 0\}.$$

As before,  $p_A$  induces a function on double cosets  $K \backslash T_n / \text{Stab}_T(A)$ , in particular  $p_M$  determines a real valued function on  $K \backslash T_n \simeq \mathbb{R}^n$  (the isomorphism is given by the map  $\text{diag}(t_1, \dots, t_n) \xrightarrow{\log} (\log |t_1|, \dots, \log |t_n|)$ ). That is,

$$\begin{array}{ccc} T_n & \xrightarrow{\log} & K \backslash T_n \simeq \mathbb{R}^n \\ \downarrow p_A & \swarrow \hat{p}' & \downarrow \\ \mathbb{R}_+ & \xleftarrow{p_A''} & K \backslash T_n / \text{Stab}_T(A) \end{array}$$

where the function  $p_A''$  is the *special function*

$$p_A''(r_1, \dots, r_n) = e^{2 \log \|A(0)\|} + \sum_{i,j:A(i,j) \neq 0}^n e^{2 \log \|A(i,j)\| + 2x_i - 2x_j}$$

and where  $K \backslash T_n / \text{Stab}_T(A)$  is the quotient space of  $\mathbb{R}^n$  by the subspace  $V_A$  which is the image of  $\text{Stab}_T(A)$  under  $\log$

$$V_A = \sum_{i:\exists A(i,j) \neq 0} \mathbb{R}e_i + \sum_{i,j:A(i,j) \neq 0} \mathbb{R}(e_i - e_j)$$

where  $e_i$  are the standard basis vectors of  $\mathbb{R}^n$ . Let  $\{i_1, \dots, i_k\}$  be the minimal elements of the non-empty equivalence classes induced by the relation  $i \sim j$  iff  $A(i,j) \neq 0$ , then

$$\begin{cases} K \backslash T_n / \text{Stab}_T(A) \simeq \sum_{j=1}^k \mathbb{R}e_{i_j} \\ p_A''(y_1, \dots, y_k) = c_0 + \sum_{j=1}^k (\sum_{l(j)} c_{l(j)} e^{a_{l(j)} y_j}) \end{cases}$$

for certain positive real numbers  $c_0, c_{l(j)}$  and real numbers  $a_{l(j)}$ . But then, elementary calculus shows that the  $k \times k$  matrix

$$\begin{bmatrix} \frac{\partial^2 p_A''}{\partial y_1 \partial y_1}(m) & \dots & \frac{\partial^2 p_A''}{\partial y_1 \partial y_k}(m) \\ \vdots & & \vdots \\ \frac{\partial^2 p_A''}{\partial y_k \partial y_1}(m) & \dots & \frac{\partial^2 p_A''}{\partial y_k \partial y_k}(m) \end{bmatrix}$$

is a positive definite diagonal matrix in every point  $m \in \mathbb{R}^k$ . That is,  $p_A$  is a strictly convex *Morse function* and if it has a critical point  $m_0$  (that is, if all  $\frac{\partial p_A}{\partial y_i}(m_0) = 0$ ), it must be a unique minimum. Lifting this information from the double coset space  $K \backslash T_n / \text{Stab}_T(A)$  to  $T_n$  we have proved

PROPOSITION 2.48. *Let  $T_n$  be the maximal torus of invertible diagonal matrices in  $GL_n$  and consider the restriction of the function  $GL_n \xrightarrow{p_A} \mathbb{R}_+$  to  $T_n$  for  $A \in M_n^m$ , then*

- (1) *Any critical point of  $p_A$  is a point where  $p_A$  obtains its minimal value.*
- (2) *If  $p_A$  obtains a minimal value, then*
  - *the set  $V$  where  $p_A$  obtains this minimum consists of a single  $K - \text{Stab}_T(A)$  coset in  $T_n$  and is connected.*
  - *the second order variation of  $p_A$  at a point of  $V$  in any direction not tangent to  $V$  is positive.*

The same proof applies to all maximal tori  $T$  of  $GL_n$  which are defined over  $\mathbb{R}$ . Recall the *Cartan decomposition* of  $GL_n$  which we proved before theorem 2.27 : any  $g \in GL_n$  can be written as  $g = udu'$  where  $u, u' \in U_n$  and  $d$  is a diagonal matrix with positive real entries. Using this fact we can now extend the above proposition to  $GL_n$ .

THEOREM 2.49. *Consider the function  $GL_n \xrightarrow{p_A} \mathbb{R}_+$  for  $A \in M_n^m$ .*

- (1) *Any critical point of  $p_A$  is a point where  $p_A$  obtains its minimal value.*
- (2) *If  $p_A$  obtains its minimal value, it does so on a single  $U_n - \text{Stab}(A)$ -coset.*

PROOF. (1) : Because for any  $h \in GL_n$  we have that  $p_{h.A}(g) = p_A(gh)$  we may assume that  $\mathbb{1}_n$  is the critical point of  $p_A$ . We have to prove that  $p_A(g) \geq p_A(\mathbb{1}_n)$  for all  $g \in GL_n$ . By the Cartan decomposition  $g = udu'$  whence  $g = u''t$  where  $u'' = uu' \in U_n$  and  $t = u'^{-1}du' \in T$  a maximal torus of  $GL_n$  defined over  $\mathbb{R}$ . Because the Hermitian inproduct is invariant under  $U_n$  we have that  $p_A(g) = p_A(t)$ . Because  $\mathbb{1}_n$  is a critical point for the restriction of  $p_A$  to  $T$  we have by proposition 2.48 that  $p_A(t) \geq p_A(\mathbb{1}_n)$ , proving the claim.

(2) : Because for any  $h \in GL_n$ ,  $p_{h.A}(g) = p_A(gh)$  and  $\text{Stab}(h.A) = h\text{Stab}(A)h^{-1}$  we may assume that  $p_A$  obtains its minimal value at  $\mathbb{1}_n$ . If  $V$  denotes the subset of  $GL_n$  where  $p_A$  obtains its minimal value we then have that  $U_n\text{Stab}(A) \subset V$  and we have to prove the reverse inclusion. Assume  $g \in V$  and write as before  $g = u''t$  with  $u'' \in U_n$  and  $t \in T$  a maximal torus defined over  $\mathbb{R}$ . Then, by unitary invariance of the inproduct,  $t$  is a point of  $T$  where the restriction of  $p_A$  to  $T$  obtains its minimal value  $p_A(\mathbb{1}_n)$ . By proposition 2.48 we conclude that  $t \in K_T\text{Stab}_T(A)$  where  $K_T = U_n \cap T$ . But then,

$$V \subset U_n \left( \bigcup_T K_T \text{Stab}_T(A) \right) \subset U_n \text{Stab}(A)$$

where  $T$  runs over all maximal tori of  $GL_n$  which are defined over  $\mathbb{R}$ , finishing the proof.  $\square$

PROPOSITION 2.50. *The function  $p_A : GL_n \longrightarrow \mathbb{R}_+$  obtains a minimal value if and only if  $\mathcal{O}(A)$  is a closed orbit in  $M_n^m$ , that is, determines a semi-simple representation.*

PROOF. If  $\mathcal{O}(A)$  is closed then  $p_A$  clearly obtains a minimal value. Conversely, assume that  $\mathcal{O}(A)$  is not closed, that is,  $A$  does not determine a semi-simple  $n$ -dimensional representation  $M$  of  $\mathbb{C}\langle x_1, \dots, x_m \rangle$ . By choosing a basis in  $M$  (that is, possibly going to another point in the orbit  $\mathcal{O}(A)$ ) we have a one-parameter subgroup  $\mathbb{C}^* \xrightarrow{\lambda} T_n \hookrightarrow GL_n$  corresponding to the Jordan-Hölder filtration of  $M$  with  $\lim_{t \rightarrow 0} \lambda(t)A = B$  with  $B$  corresponding to the semi-simplification of  $M$ . Now consider the restriction of  $p'_A$  to  $U_1 \setminus \mathbb{C}^* \simeq \mathbb{R}$ , then as before we can write it uniquely in the form

$$p'_A(x) = \sum_i a_i e^{l_i x} \quad a_i > 0, \quad l_1 < l_2 < \dots < l_z$$

for some real numbers  $l_i$  and some  $z$ . Because the above limit exists, the limit

$$\lim_{x \rightarrow -\infty} p'_A(x) \in \mathbb{R}$$

and hence none of the  $l_i$  are negative. Further, because  $\mathcal{O}(A) \neq \mathcal{O}(B)$  at least one of the  $l_i$  must be positive. Therefore,  $p'_A$  is a strictly increasing function on  $\mathbb{R}$  whence never obtains a minimal value, whence neither does  $p_A$ .  $\square$

Finally, we have to clarify the connection between the function  $p_A$  and the *real moment map*

$$\begin{cases} M_n^m \xrightarrow{\mu} (\text{Lie } U_n)^* & A \longrightarrow (h \mapsto (h.A, A)) \\ M_n^n \xrightarrow{\mu_{\mathbb{R}}} \text{Lie } U_n & A \longrightarrow i[A, A^*] \end{cases}$$

Assume  $A \in M_n^m$  is such that  $p_A$  has a critical point, which we may assume to be  $\mathfrak{U}_n$  by an argument as in the proof of theorem 2.49. Then, the differential in  $\mathfrak{U}_n$

$$(dp_A)_{\mathfrak{U}_n} : M_n = T_{\mathfrak{U}_n} GL_n \longrightarrow \mathbb{R} \quad \text{satisfies} \quad (dp_A)_{\mathfrak{U}_n}(h) = 0 \quad \forall h \in M_n$$

Let us work out this differential

$$\begin{aligned} p_A(\mathfrak{U}_n) + \epsilon(dp_A)_{\mathfrak{U}_n}(h) &= \text{tr}((A^* + \epsilon(A^*h^* - h^*A^*))(A + \epsilon(hA - Ah)) \\ &= \text{tr}(A^*A) + \epsilon \text{tr}(A^*hA - A^*Ah + A^*h^*A - h^*A^*A) \\ &= \text{tr}(A^*A) + \epsilon \text{tr}((AA^* - A^*A)(h - h^*)) \end{aligned}$$

But then, vanishing of the differential for all  $h \in M_n$  is equivalent by the nondegeneracy of the Killing form on  $\text{Lie } U_n$  to

$$AA^* - A^*A = \sum_{i=1}^m A_i A_i^* - A_i^* A_i = 0$$

that is, to  $A \in \mu_{\mathbb{R}}^{-1}(0)$ . This concludes the proof of the main result on the real moment map for  $M_n^m$ .

**THEOREM 2.51.** *There are natural one-to-one correspondences between*

- (1) *isomorphism classes of semi-simple  $n$ -dimensional representations of  $\mathbb{C}\langle x_1, \dots, x_m \rangle$ ,*
- (2) *closed  $GL_n$ -orbits in  $M_n^m$ ,*
- (3)  *$U_n$ -orbits in the subset  $\mu_{\mathbb{R}}^{-1}(0) = \{A \in M_n^m \mid \sum_{i=1}^m [A_i, A_i^*] = 0\}$ .*

Let  $A \in \mathfrak{alg} @_{\mathbf{n}}$  be an affine Cayley-Hamilton algebra of degree  $n$ , then we can embed the reduced variety of  $\text{rep}_n^{\text{tr}} A$  in  $M_n^m$  and obtain as a consequence :

**THEOREM 2.52.** *For  $A \in \mathfrak{alg} @_{\mathbf{n}}$ , there are natural one-to-one correspondences between*

- (1) *isomorphism classes of semi-simple  $n$ -dimensional trace preserving representations of  $A$ ,*
- (2) *closed  $GL_n$ -orbits in the representation variety  $\underline{rep}_n^{tr} A$ ,*
- (3)  *$U_n$ -orbits in the intersection  $\underline{rep}_n^{tr} A \cap \mu_{\mathbb{R}}^{-1}(0)$ .*

### References.

The generalization of the Hilbert criterium, theorem 2.17 is due to D. Birkes [7]. The connection between semi-simple representations and closed orbits is due to M. Artin [2]. The geometric reconstruction result follows from theorem 1.28 and is due to C. Procesi [65]. The results on the real moment map are due to G. Kempf and L. Ness [37]. The treatment of the Hilbert criterium and invariant theory follows the textbook of H-P. Kraft [45], that of the Gerstenhaber-Hesselink result owes to the exposition of M. Hazewinkel in [29].



## top @<sub>n</sub>

The natural topology for noncommutative geometry @<sub>n</sub> is the étale topology. Let  $A$  be an affine  $\mathbb{C}$ -algebra and  $\xi \in \underline{iss}_n A$  with corresponding semi-simple  $n$ -dimensional representation  $M_\xi$ . We say that  $\xi$  is a smooth point for the level  $n$  approximation  $A @_n$  if the representation space  $\underline{rep}_n A$  is smooth in  $M_\xi$ . The main result, theorem 4.24, asserts that we can recover the étale local structure of  $\underline{iss}_n A$  and of the algebra  $A @_n$  near a smooth point  $\xi$  from combinatorial data.

Let  $M_\xi = S_1^{\oplus e_1} \oplus \dots \oplus S_k^{\oplus e_k}$  be the decomposition of  $M_\xi$  into distinct simple components  $S_i$  of dimension  $d_i$  and occurring in  $M_\xi$  with multiplicity  $e_i$ . We construct a quiver  $Q_\xi$  with  $k$  vertices  $\{v_1, \dots, v_k\}$  (where  $v_i$  corresponds to the simple component  $S_i$ ) and the number of oriented arrows from  $v_i$  to  $v_j$  is given by

$$\# \textcircled{j} \leftarrow \textcircled{i} = \dim_{\mathbb{C}} \text{Ext}_A^1(S_i, S_j)$$

Further, we define the dimension vector  $\alpha_\xi = (e_1, \dots, e_k)$  describing the multiplicities of the simple components in  $M_\xi$ . Assigning to every vertex  $v_i$  the space  $\mathbb{C}^{e_i}$  and to every arrow  $\textcircled{j} \leftarrow \textcircled{i}$  a matrix  $\mathbb{C}^{e_i} \rightarrow \mathbb{C}^{e_j}$  we obtain the representation space  $\underline{rep}_{\alpha_\xi} Q_\xi$  on which the basechange group  $GL(\alpha_\xi) = GL_{e_1} \times \dots \times GL_{e_k}$  acts with quotient variety  $\underline{iss}_{\alpha_\xi} Q_\xi$  classifying  $\alpha_\xi$ -dimensional semi-simple  $Q_\xi$ -representations.

The coordinate ring  $\mathbb{N}_\alpha$  of the quotient variety  $\underline{iss}_{\alpha_\xi} Q_\xi$  is generated by traces along oriented cycles in the quiver  $Q_\xi$  and the ring  $\mathbb{T}_\alpha$  of  $GL(\alpha)$ -equivariant maps  $\underline{rep}_{\alpha_\xi} Q_\xi \rightarrow M_m(\mathbb{C})$  where  $m = \sum_i e_i$  has a block-decomposition with  $(i, j)$ -entry the  $\mathbb{N}_\alpha$ -module of oriented paths from  $v_i$  to  $v_j$  in  $Q_\xi$ . The completions of  $\mathbb{N}_\alpha$  and  $\mathbb{T}_\alpha$  with respect to the maximal graded ideal of  $\mathbb{N}_\alpha$  will be denoted by  $\widehat{\mathbb{N}}_\alpha$  and  $\widehat{\mathbb{T}}_\alpha$  respectively.

Let  $\mathfrak{m}$  be the maximal ideal of the coordinate ring  $N = \text{tr} A @_n$  of  $\underline{iss}_n A$  and denote  $T = A @_n$ , then we have the étale local description

$$\widehat{N}_{\mathfrak{m}} \simeq \widehat{\mathbb{N}}_\alpha \quad \text{and} \quad \widehat{T}_{\mathfrak{m}} \underset{\text{Morita}}{\sim} \widehat{\mathbb{T}}_\alpha$$

where the Morita equivalence is determined by the embedding  $GL(\alpha) \hookrightarrow GL_n$  given by the dimensions  $d_i$  of the simple components. An extension of this local description to points  $\xi \in \underline{iss}_n A$  which are not smooth for  $A @_n$  will involve the canonical  $A_\infty$ -structure on  $\text{Ext}_A^*(M_\xi, M_\xi)$ .

A smooth manifold in noncommutative geometry @<sub>n</sub>, that is a family  $(X_i)_{i \in I}$  of algebraic varieties locally controlled by a set  $\mathcal{A}$  of Quillen-smooth algebras, can therefore be studied locally by the combinatorics of quiver representations.



CHAPTER 3

## Etale Cohomology.

### 3.1. Etale topology.

A closed subvariety  $X \hookrightarrow \mathbb{C}^m$  can be equipped with the *Zariski topology* or with the much finer *analytic topology*. A major disadvantage of the coarseness of the Zariski topology is the failure to have an *implicit function* theorem in algebraic geometry. Etale morphisms are introduced to bypass this problem. These morphisms determine the *étale topology* which is no longer a topology determined by subsets but rather a *Grothendieck topology* determined by *covers*.

DEFINITION 3.1. A finite morphism  $A \xrightarrow{f} B$  of commutative  $\mathbb{C}$ -algebras is said to be *étale* if and only if  $B = A[t_1, \dots, t_k]/(f_1, \dots, f_k)$  such that the *Jacobian matrix*

$$\begin{bmatrix} \frac{\partial f_1}{\partial t_1} & \cdots & \frac{\partial f_1}{\partial t_k} \\ \vdots & & \vdots \\ \frac{\partial f_k}{\partial t_1} & \cdots & \frac{\partial f_k}{\partial t_k} \end{bmatrix}$$

has a determinant which is a unit in  $B$ .

With  $\text{spec } A$  we denote the *prime ideal spectrum* or the *affine scheme* of a commutative  $\mathbb{C}$ -algebra  $A$ . That is,  $\text{spec } A$  is the set of all *prime ideals* of  $A$  equipped with the *Zariski topology*, that is the open subset are of the form

$$\mathbb{X}(I) = \{P \in \text{spec } A \mid I \not\subset P\}$$

for some ideal  $I \triangleleft A$ . If  $A$  is an affine  $\mathbb{C}$ -algebra, the points of the corresponding affine variety correspond to the *maximal ideals* of  $A$  and the induced Zariski topology coincides with the one introduced before. In this chapter, however, not all  $\mathbb{C}$ -algebras will be affine.

EXAMPLE 3.2. Consider the morphism  $\mathbb{C}[x, x^{-1}] \hookrightarrow \mathbb{C}[x, x^{-1}][\sqrt[n]{x}]$  and the induced map on the affine schemes

$$\text{spec } \mathbb{C}[x, x^{-1}][\sqrt[n]{x}] \xrightarrow{\psi} \text{spec } \mathbb{C}[x, x^{-1}] = \mathbb{C} - \{0\}.$$

Clearly, every point  $\lambda \in \mathbb{C} - \{0\}$  has exactly  $n$  preimages  $\lambda_i = \zeta^i \sqrt[n]{\lambda}$ . Moreover, in a neighborhood of  $\lambda_i$ , the map  $\psi$  is a diffeomorphism. Still, we do not have an inverse map in algebraic geometry as  $\sqrt[n]{x}$  is not a polynomial map. However,  $\mathbb{C}[x, x^{-1}][\sqrt[n]{x}]$  is an étale extension of  $\mathbb{C}[x, x^{-1}]$ . In this way étale morphisms can be seen as an algebraic substitute for the failure of an inverse function theorem in algebraic geometry.

PROPOSITION 3.3. *Etale morphisms satisfy 'sorite', that is, they satisfy the commutative diagrams of figure 1. In these diagrams,  $et$  denotes an étale morphism,  $f.f.$  denotes a faithfully flat morphism and the dashed arrow is the étale morphism implied by 'sorite'.*





FIGURE 1. Sorite for étale morphisms

With these properties we can define a *Grothendieck topology* on the collection of all étale morphisms.

DEFINITION 3.4. The *étale site* of  $A$ , which we will denote by  $\mathcal{A}_{\text{ét}}$  is the category with

- objects : the étale extensions  $A \xrightarrow{f} B$  of  $A$
- morphisms : compatible  $A$ -algebra morphisms



By proposition 3.3 all morphisms in  $\mathcal{A}_{\text{ét}}$  are étale. We can turn  $\mathcal{A}_{\text{ét}}$  into a Grothendieck topology by defining

- cover : a collection  $\mathcal{C} = \{B \xrightarrow{f_i} B_i\}$  in  $\mathcal{A}_{\text{ét}}$  such that

$$\text{spec } B = \cup_i \text{Im} (\text{spec } B_i \xrightarrow{f} \text{Spec } B)$$

DEFINITION 3.5. An *étale presheaf* of groups on  $\mathcal{A}_{\text{ét}}$  is a functor

$$\mathbb{G} : \mathcal{A}_{\text{ét}} \longrightarrow \text{groups}$$

In analogy with usual (pre)sheaf notation we denote for each

- object  $B \in \mathcal{A}_{\text{ét}}$  the *global sections*  $\Gamma(B, \mathbb{G}) = \mathbb{G}(B)$
- morphism  $B \xrightarrow{\phi} C$  in  $\mathcal{A}_{\text{ét}}$  the *restriction map*  $\text{Res}_C^B = \mathbb{G}(\phi) : \mathbb{G}(B) \longrightarrow \mathbb{G}(C)$  and  $g|_C = \mathbb{G}(\phi)(g)$ .

An étale presheaf  $\mathbb{G}$  is an *étale sheaf* provided for every  $B \in \mathbf{A}_{\text{ét}}$  and every cover  $\{B \rightarrow B_i\}$  we have exactness of the *equalizer diagram*

$$0 \longrightarrow \mathbb{G}(B) \longrightarrow \prod_i \mathbb{G}(B_i) \rightrightarrows \prod_{i,j} \mathbb{G}(B_i \otimes_B B_j)$$

EXAMPLE 3.6. *Constant sheaf.*

If  $G$  is a group, then

$$\mathbb{G} : \mathbf{A}_{\text{ét}} \longrightarrow \text{groups} \quad B \mapsto G^{\oplus \pi_0(B)}$$

is a sheaf where  $\pi_0(B)$  is the number of connected components of  $\text{Spec } B$ .

EXAMPLE 3.7. *Multiplicative group  $\mathbb{G}_m$ .*

The functor

$$\mathbb{G}_m : \mathbf{A}_{\text{ét}} \longrightarrow \text{groups} \quad B \mapsto B^*$$

is a sheaf on  $\mathbf{A}_{\text{ét}}$ .

A sequence of sheaves of Abelian groups on  $\mathbf{A}_{\text{ét}}$  is said to be exact

$$\mathbb{G}' \xrightarrow{f} \mathbb{G} \xrightarrow{g} \mathbb{G}''$$

if for every  $B \in \mathbf{A}_{\text{ét}}$  and  $s \in \mathbb{G}(B)$  such that  $g(s) = 0 \in \mathbb{G}''(B)$  there is a cover  $\{B \rightarrow B_i\}$  in  $\mathbf{A}_{\text{ét}}$  and sections  $t_i \in \mathbb{G}'(B_i)$  such that  $f(t_i) = s|_{B_i}$ .

EXAMPLE 3.8. *Roots of unity  $\mu_n$ .*

We have a sheaf morphism

$$\mathbb{G}_m \xrightarrow{(-)^n} \mathbb{G}_m$$

and we denote the kernel with  $\mu_n$ . As  $A$  is a  $\mathbb{C}$ -algebra we can identify  $\mu_n$  with the constant sheaf  $\mathbb{Z}_n = \mathbb{Z}/n\mathbb{Z}$  via the isomorphism  $\zeta^i \mapsto i$  after choosing a primitive  $n$ -th root of unity  $\zeta \in \mathbb{C}$ .

LEMMA 3.9. *The Kummer sequence of sheaves of Abelian groups*

$$0 \longrightarrow \mu_n \longrightarrow \mathbb{G}_m \xrightarrow{(-)^n} \mathbb{G}_m \longrightarrow 0$$

is exact on  $\mathbf{A}_{\text{ét}}$  (but not necessarily on  $\text{spec } A$  with the Zariski topology).

PROOF. We only need to verify surjectivity. Let  $B \in \mathbf{A}_{\text{ét}}$  and  $b \in \mathbb{G}_m(B) = B^*$ . Consider the étale extension  $B' = B[t]/(t^n - b)$  of  $B$ , then  $b$  has an  $n$ -th root over in  $\mathbb{G}_m(B')$ . Observe that this  $n$ -th root does not have to belong to  $\mathbb{G}_m(B)$ .  $\square$

If  $\mathfrak{p}$  is a prime ideal of  $A$  we will denote with  $\mathbf{k}_{\mathfrak{p}}$  the algebraic closure of the field of fractions of  $A/\mathfrak{p}$ . An *étale neighborhood* of  $\mathfrak{p}$  is an étale extension  $B \in \mathbf{A}_{\text{ét}}$  such that the diagram below is commutative

$$\begin{array}{ccc} A & \xrightarrow{\text{nat}} & \mathbf{k}_{\mathfrak{p}} \\ \text{et} \downarrow & \nearrow & \\ B & & \end{array}$$

The analogue of the localization  $A_{\mathfrak{p}}$  for the étale topology is the *strict Henselization*

$$A_{\mathfrak{p}}^{\text{sh}} = \varinjlim B$$

where the limit is taken over all étale neighborhoods of  $\mathfrak{p}$ .

Recall that a local algebra  $L$  with maximal ideal  $m$  and residue map  $\pi : L \rightarrow L/m = k$  is said to be *Henselian* if the following condition holds. Let

$f \in L[t]$  be a monic polynomial such that  $\pi(f)$  factors as  $g_0 \cdot h_0$  in  $k[t]$ , then  $f$  factors as  $g \cdot h$  with  $\pi(g) = g_0$  and  $\pi(h) = h_0$ . If  $L$  is Henselian then tensoring with  $k$  induces an equivalence of categories between the étale  $A$ -algebras and the étale  $k$ -algebras.

An Henselian local algebra is said to be *strict Henselian* if and only if its residue field is algebraically closed. Thus, a strict Henselian ring has no proper finite étale extensions and can be viewed as a local algebra for the étale topology.

EXAMPLE 3.10. The algebraic functions  $\mathbb{C}\{x_1, \dots, x_d\}$

Consider the local algebra of  $\mathbb{C}[x_1, \dots, x_d]$  in the maximal ideal  $(x_1, \dots, x_d)$ , then the Henselization and strict Henselization are both equal to

$$\mathbb{C}\{x_1, \dots, x_d\}$$

the ring of *algebraic functions*. That is, the subalgebra of  $\mathbb{C}[[x_1, \dots, x_d]]$  of formal power-series consisting of those series  $\phi(x_1, \dots, x_d)$  which are algebraically dependent on the coordinate functions  $x_i$  over  $\mathbb{C}$ . In other words, those  $\phi$  for which there exists a non-zero polynomial  $f(x_i, y) \in \mathbb{C}[x_1, \dots, x_d, y]$  with  $f(x_1, \dots, x_d, \phi(x_1, \dots, x_d)) = 0$ .

These algebraic functions may be defined implicitly by polynomial equations. Consider a system of equations

$$f_i(x_1, \dots, x_d; y_1, \dots, y_m) = 0 \text{ for } f_i \in \mathbb{C}[x_i, y_j] \text{ and } 1 \leq i \leq m$$

Suppose there is a solution in  $\mathbb{C}$  with

$$x_i = 0 \text{ and } y_j = y_j^o$$

such that the Jacobian matrix is non-zero

$$\det \left( \frac{\partial f_i}{\partial y_j} (0, \dots, 0; y_1^o, \dots, y_m^o) \right) \neq 0$$

Then, the system can be solved uniquely for power series  $y_j(x_1, \dots, x_d)$  with  $y_j(0, \dots, 0) = y_j^o$  by solving inductively for the coefficients of the series. One can show that such implicitly defined series  $y_j(x_1, \dots, x_d)$  are algebraic functions and that, conversely, any algebraic function can be obtained in this way.

If  $\mathbb{G}$  is a sheaf on  $\mathbf{A}_{\text{ét}}$  and  $\mathfrak{p}$  is a prime ideal of  $A$ , we define the *stalk* of  $\mathbb{G}$  in  $\mathfrak{p}$  to be

$$\mathbb{G}_{\mathfrak{p}} = \varinjlim \mathbb{G}(B)$$

where the limit is taken over all étale neighborhoods of  $\mathfrak{p}$ . One can verify mono-epi- or isomorphisms of sheaves by checking it in all the stalks.

If  $A$  is an affine algebra defined over an algebraically closed field, then it suffices to verify it in the maximal ideals of  $A$ .

Before we define cohomology of sheaves on  $\mathbf{A}_{\text{ét}}$  let us recall the definition of *derived functors*. Let  $\mathcal{A}$  be an *Abelian category*. An object  $I$  of  $\mathcal{A}$  is said to be *injective* if the functor

$$\mathcal{A} \longrightarrow \text{abellan} \quad M \mapsto \text{Hom}_{\mathcal{A}}(M, I)$$

is exact. We say that  $\mathcal{A}$  has enough injectives if, for every object  $M$  in  $\mathcal{A}$ , there is a monomorphism  $M \hookrightarrow I$  into an injective object.

If  $\mathcal{A}$  has enough injectives and  $f : \mathcal{A} \longrightarrow \mathcal{B}$  is a left exact functor from  $\mathcal{A}$  into a second Abelian category  $\mathcal{B}$ , then there is an essentially unique sequence of functors

$$R^i f : \mathcal{A} \longrightarrow \mathcal{B} \quad i \geq 0$$

called the *right derived functors* of  $f$  satisfying the following properties

- $R^0 f = f$

- $R^i I = 0$  for  $I$  injective and  $i > 0$
- For every short exact sequence in  $\mathcal{A}$

$$0 \longrightarrow M' \longrightarrow M \longrightarrow M'' \longrightarrow 0$$

there are connecting morphisms  $\delta^i : R^i f(M'') \longrightarrow R^{i+1} f(M')$  for  $i \geq 0$  such that we have a long exact sequence

$$\dots \longrightarrow R^i f(M) \longrightarrow R^i f(M'') \xrightarrow{\delta^i} R^{i+1} f(M') \longrightarrow R^{i+1} f(M) \longrightarrow \dots$$

- For any morphism  $M \longrightarrow N$  there are morphisms  $R^i f(M) \longrightarrow R^i f(N)$  for  $i \geq 0$

In order to compute the objects  $R^i f(M)$  define an object  $N$  in  $\mathcal{A}$  to be *f-acyclic* if  $R^i f(N) = 0$  for all  $i > 0$ . If we have an *acyclic resolution* of  $M$

$$0 \longrightarrow M \longrightarrow N_0 \longrightarrow N_1 \longrightarrow N_2 \longrightarrow \dots$$

by *f-acyclic* object  $N_i$ , then the objects  $R^i f(M)$  are canonically isomorphic to the cohomology objects of the complex

$$0 \longrightarrow f(N_0) \longrightarrow f(N_1) \longrightarrow f(N_2) \longrightarrow \dots$$

One can show that all injectives are *f-acyclic* and hence that derived objects of  $M$  can be computed from an *injective resolution* of  $M$ .

Now, let  $\mathbf{S}^{ab}(\mathbf{A}_{\text{ét}})$  be the category of all sheaves of Abelian groups on  $\mathbf{A}_{\text{ét}}$ . This is an Abelian category having enough injectives whence we can form right derived functors of left exact functors. In particular, consider the global section functor

$$\Gamma : \mathbf{S}^{ab}(\mathbf{A}_{\text{ét}}) \longrightarrow \mathbf{Ab} \quad \mathbb{G} \mapsto \mathbb{G}(A)$$

which is left exact. The right derived functors of  $\Gamma$  will be called the *étale cohomology functors* and we denote

$$R^i \Gamma(\mathbb{G}) = H_{\text{ét}}^i(A, \mathbb{G})$$

In particular, if we have an exact sequence of sheaves of Abelian groups  $0 \longrightarrow \mathbb{G}' \longrightarrow \mathbb{G} \longrightarrow \mathbb{G}'' \longrightarrow 0$ , then we have a long exact cohomology sequence

$$\dots \longrightarrow H_{\text{ét}}^i(A, \mathbb{G}) \longrightarrow H_{\text{ét}}^i(A, \mathbb{G}'') \longrightarrow H_{\text{ét}}^{i+1}(A, \mathbb{G}') \longrightarrow \dots$$

If  $\mathbb{G}$  is a sheaf of non-Abelian groups (written multiplicatively), we cannot define cohomology groups. Still, one can define a *pointed set*  $H_{\text{ét}}^1(A, \mathbb{G})$  as follows. Take an étale cover  $\mathcal{C} = \{A \longrightarrow A_i\}$  of  $A$  and define a 1-cocycle for  $\mathcal{C}$  with values in  $\mathbb{G}$  to be a family

$$g_{ij} \in \mathbb{G}(A_{ij}) \text{ with } A_{ij} = A_i \otimes_A A_j$$

satisfying the cocycle condition

$$(g_{ij} | A_{ijk})(g_{jk} | A_{ijk}) = (g_{ik} | A_{ijk})$$

where  $A_{ijk} = A_i \otimes_A A_j \otimes_A A_k$ .

Two cocycles  $g$  and  $g'$  for  $\mathcal{C}$  are said to be cohomologous if there is a family  $h_i \in \mathbb{G}(A_i)$  such that for all  $i, j \in I$  we have

$$g'_{ij} = (h_i | A_{ij})g_{ij}(h_j | A_{ij})^{-1}$$

This is an equivalence relation and the set of cohomology classes is written as  $H_{\text{ét}}^1(\mathcal{C}, \mathbb{G})$ . It is a pointed set having as its distinguished element the cohomology class of  $g_{ij} = 1 \in \mathbb{G}(A_{ij})$  for all  $i, j \in I$ .

We then define the non-Abelian first *cohomology pointed set* as

$$H_{et}^1(A, \mathbb{G}) = \varinjlim H_{et}^1(\mathcal{C}, \mathbb{G})$$

where the limit is taken over all étale coverings of  $A$ . It coincides with the previous definition in case  $\mathbb{G}$  is Abelian.

A sequence  $1 \longrightarrow \mathbb{G}' \longrightarrow \mathbb{G} \longrightarrow \mathbb{G}'' \longrightarrow 1$  of sheaves of groups on  $\mathbf{A}_{et}$  is said to be exact if for every  $B \in \mathbf{A}_{et}$  we have

- $\mathbb{G}'(B) = \text{Ker } \mathbb{G}(B) \longrightarrow \mathbb{G}''(B)$
- For every  $g'' \in \mathbb{G}''(B)$  there is a cover  $\{B \longrightarrow B_i\}$  in  $\mathbf{A}_{et}$  and sections  $g_i \in \mathbb{G}(B_i)$  such that  $g_i$  maps to  $g''|_{B_i}$ .

PROPOSITION 3.11. *For an exact sequence of groups on  $\mathbf{A}_{et}$*

$$1 \longrightarrow \mathbb{G}' \longrightarrow \mathbb{G} \longrightarrow \mathbb{G}'' \longrightarrow 1$$

*there is associated an exact sequence of pointed sets*

$$\begin{aligned} 1 \longrightarrow \mathbb{G}'(A) \longrightarrow \mathbb{G}(A) \longrightarrow \mathbb{G}''(A) &\xrightarrow{\delta} H_{et}^1(A, \mathbb{G}') \longrightarrow \\ &\longrightarrow H_{et}^1(A, \mathbb{G}) \longrightarrow H_{et}^1(A, \mathbb{G}'') \cdots \cdots \longrightarrow H_{et}^2(A, \mathbb{G}') \end{aligned}$$

*where the last map exists when  $\mathbb{G}'$  is contained in the center of  $\mathbb{G}$  (and therefore is Abelian whence  $H^2$  is defined).*

PROOF. The connecting map  $\delta$  is defined as follows. Let  $g'' \in \mathbb{G}''(A)$  and let  $\mathcal{C} = \{A \longrightarrow A_i\}$  be an étale covering of  $A$  such that there are  $g_i \in \mathbb{G}(A_i)$  that map to  $g''|_{A_i}$  under the map  $\mathbb{G}(A_i) \longrightarrow \mathbb{G}''(A_i)$ . Then,  $\delta(g)$  is the class determined by the one cocycle

$$g_{ij} = (g_i|_{A_{ij}})^{-1}(g_j|_{A_{ij}})$$

with values in  $\mathbb{G}'$ . The last map can be defined in a similar manner, the other maps are natural and one verifies exactness.  $\square$

The main applications of this non-Abelian cohomology to non-commutative algebra is as follows. Let  $\Lambda$  be a not necessarily commutative  $A$ -algebra and  $M$  an  $A$ -module. Consider the sheaves of groups  $\text{Aut}(\Lambda)$  resp.  $\text{Aut}(M)$  on  $\mathbf{A}_{et}$  associated to the presheaves

$$B \mapsto \text{Aut}_{B\text{-alg}}(\Lambda \otimes_A B) \text{ resp. } B \mapsto \text{Aut}_{B\text{-mod}}(M \otimes_A B)$$

for all  $B \in \mathbf{A}_{et}$ . A *twisted form* of  $\Lambda$  (resp.  $M$ ) is an  $A$ -algebra  $\Lambda'$  (resp. an  $A$ -module  $M'$ ) such that there is an étale cover  $\mathcal{C} = \{A \longrightarrow A_i\}$  of  $A$  such that there are isomorphisms

$$\begin{cases} \Lambda \otimes_A A_i \xrightarrow{\phi_i} \Lambda' \otimes_A A_i \\ M \otimes_A A_i \xrightarrow{\psi_i} M' \otimes_A A_i \end{cases}$$

of  $A_i$ -algebras (resp.  $A_i$ -modules). The set of  $A$ -algebra isomorphism classes (resp.  $A$ -module isomorphism classes) of twisted forms of  $\Lambda$  (resp.  $M$ ) is denoted by  $Tw_A(\Lambda)$  (resp.  $Tw_A(M)$ ). To a twisted form  $\Lambda'$  one associates a cocycle on  $\mathcal{C}$

$$\alpha_{\Lambda'} = \alpha_{ij} = \phi_i^{-1} \circ \phi_j$$

with values in  $\text{Aut}(\Lambda)$ . Moreover, one verifies that two twisted forms are isomorphic as  $A$ -algebras if their cocycles are cohomologous. That is, there are embeddings

$$\begin{cases} Tw_A(\Lambda) \hookrightarrow H_{et}^1(A, \text{Aut}(\Lambda)) \\ Tw_A(M) \hookrightarrow H_{et}^1(A, \text{Aut}(M)) \end{cases}$$



In favorable situations one can even show bijectivity. In particular, this is the case if the automorphisms group is a smooth affine algebraic group-scheme.

For example, consider  $\Lambda = M_n(A)$ , then the automorphism group is  $PGL_n$  and twisted forms of  $\Lambda$  are classified by elements of the cohomology group

$$H_{et}^1(A, PGL_n)$$

These twisted forms are precisely the *Azumaya algebras* of rank  $n^2$  with center  $A$ . When  $A$  is an affine commutative  $\mathbb{C}$ -algebra and  $\Lambda$  is an  $A$ -algebra with center  $A$ , then  $\Lambda$  is an *Azumaya algebra* of rank  $n^2$  if and only if

$$\frac{\Lambda}{\lambda \mathfrak{m} \Lambda} \simeq M_n(\mathbb{C})$$

for every maximal ideal  $\mathfrak{m}$  of  $A$ .

We will often encounter Azumaya algebras as follows. Let  $A$  be this time a *noncommutative* affine  $\mathbb{C}$ -algebra and assume that the following two conditions are satisfied

- $A$  has a simple representation of dimension  $n$ ,
- $\underline{rep}_n A$  is an irreducible variety.

Then  $tr A @_{\mathfrak{n}} = \mathbb{C}[\underline{rep}_n A]^{GL_n}$  is a domain (whence  $\underline{iss}_n A$  is irreducible) and we have an onto trace preserving algebra map corresponding to the simple representation

$$A @_{\mathfrak{n}} \xrightarrow{\phi} M_n(\mathbb{C})$$

Lift the standard basis  $e_{ij}$  of  $M_n(\mathbb{C})$  to elements  $a_{ij} \in A @_{\mathfrak{n}}$  and consider the determinant  $d$  of the  $n^2 \times n^2$  matrix  $(tr(a_{ij}a_{kl}))_{ij,kl}$  with values in  $tr A @_{\mathfrak{n}}$ . Then  $d \neq 0$  and consider the Zariski open affine subset of  $\underline{iss}_n A$

$$\mathbb{X}(d) = \{A @_{\mathfrak{n}} \xrightarrow{\psi} M_n(\mathbb{C}) \mid \psi \text{ semisimple and } \det(tr(\psi(a_{ij})\psi(a_{kl}))) \neq 0\}$$

If  $\psi \in \mathbb{X}(d)$ , then  $\psi : A @_{\mathfrak{n}} \longrightarrow M_n(\mathbb{C})$  is onto as the  $\psi(a_{ij})$  form a basis of  $M_n(\mathbb{C})$  whence  $\psi$  determines a simple  $n$ -dimensional representation.

PROPOSITION 3.12. *With notations as above,*

- (1) *The localization of  $A @_{\mathfrak{n}}$  at the central multiplicative set  $\{1, d, d^2, \dots\}$  is an affine Azumaya algebra with center  $\mathbb{C}[\mathbb{X}(d)]$  which is the localization of  $tr A @_{\mathfrak{n}}$  at this multiplicative set.*
- (2) *The restriction of the quotient map  $\underline{rep}_n A \xrightarrow{\pi} \underline{iss}_n A$  to the open set  $\pi^{-1}(\mathbb{X}(d))$  is a principal  $PGL_n$ -fibration and determines an element in*

$$H_{et}^1(\mathbb{C}[\mathbb{X}(d)], PGL_n)$$

*giving the class of the Azumaya algebra.*

PROOF. (1) : If  $\mathfrak{m} = Ker \psi$  is the maximal ideal of  $\mathbb{C}[\mathbb{X}(d)]$  corresponding to the semisimple representation  $\psi : A @_{\mathfrak{n}} \longrightarrow M_n(\mathbb{C})$ , then we have seen that the quotient

$$\frac{A @_{\mathfrak{n}}}{A @_{\mathfrak{n}} \mathfrak{m} A @_{\mathfrak{n}}} \simeq M_n(\mathbb{C})$$

whence  $A @_{\mathfrak{n}} \otimes_{tr A @_{\mathfrak{n}}} \mathbb{C}[\mathbb{X}(d)]$  is an Azumaya algebra. (2) will follow from the theory of Knop-Luna slices and will be proved in section 4.5.  $\square$

An Azumaya algebra over a field is a central simple algebra. Under the above conditions we have that

$$A @_{\mathbf{n}} \otimes_{tr A} @_{\mathbf{n}} \mathbb{C}(\underline{iss}_n A)$$

is a central simple algebra over the functionfield of  $\underline{iss}_n A$  and hence determines a class in its Brauer group, which is an important birational invariant. In the following section we recall the cohomological description of Brauer groups of fields.

### 3.2. Central simple algebras

Let  $K$  be a field of characteristic zero, choose an algebraic closure  $\mathbb{K}$  with absolute Galois group  $G_K = Gal(\mathbb{K}/K)$ .

LEMMA 3.13. *The following are equivalent*

- (1)  $K \longrightarrow A$  is étale
- (2)  $A \otimes_K \mathbb{K} \simeq \mathbb{K} \times \dots \times \mathbb{K}$
- (3)  $A = \prod L_i$  where  $L_i/K$  is a finite field extension

PROOF. Assume (1), then  $A = K[x_1, \dots, x_n]/(f_1, \dots, f_n)$  where  $f_i$  have invertible Jacobian matrix. Then  $A \otimes \mathbb{K}$  is a smooth algebra (hence reduced) of dimension 0 so (2) holds.

Assume (2), then

$$Hom_{K\text{-alg}}(A, \mathbb{K}) \simeq Hom_{\mathbb{K}\text{-alg}}(A \otimes \mathbb{K}, \mathbb{K})$$

has  $dim_{\mathbb{K}}(A \otimes \mathbb{K})$  elements. On the other hand we have by the *Chinese remainder theorem* that

$$A/Jac A = \prod_i L_i$$

with  $L_i$  a finite field extension of  $K$ . However,

$$dim_{\mathbb{K}}(A \otimes \mathbb{K}) = \sum_i dim_K(L_i) = dim_K(A/Jac A) \leq dim_K(A)$$

and as both ends are equal  $A$  is reduced and hence  $A = \prod_i L_i$  whence (3).

Assume (3), then each  $L_i = K[x_i]/(f_i)$  with  $\partial f_i/\partial x_i$  invertible in  $L_i$ . But then  $A = \prod L_i$  is étale over  $K$  whence (1).  $\square$

To every finite étale extension  $A = \prod L_i$  we can associate the finite set  $rts(A) = Hom_{K\text{-alg}}(A, \mathbb{K})$  on which the Galois group  $G_K$  acts via a finite quotient group. If we write  $A = K[t]/(f)$ , then  $rts(A)$  is the *set of roots* in  $\mathbb{K}$  of the polynomial  $f$  with obvious action by  $G_K$ . Galois theory, in the interpretation of Grothendieck, can now be stated as

PROPOSITION 3.14. *The functor*

$$K_{\text{ét}} \xrightarrow{rts(-)} \text{finite } G_K\text{-sets}$$

*is an anti-equivalence of categories.*

We will now give a similar interpretation of the Abelian sheaves on  $K_{\text{ét}}$ . Let  $\mathbb{G}$  be a presheaf on  $K_{\text{ét}}$ . Define

$$M_{\mathbb{G}} = \varinjlim \mathbb{G}(L)$$

where the limit is taken over all subfields  $L \hookrightarrow \mathbb{K}$  which are finite over  $K$ . The Galois group  $G_K$  acts on  $\mathbb{G}(L)$  on the left through its action on  $L$  whenever  $L/K$

is Galois. Hence,  $G_K$  acts on  $M_{\mathbb{G}}$  and  $M_{\mathbb{G}} = \cup M_{\mathbb{G}}^H$  where  $H$  runs through the *open subgroups* (that is, containing a normal subgroup having a finite quotient) of  $G_K$ . That is,  $M_{\mathbb{G}}$  is a *continuous  $G_K$ -module*.

Conversely, given a continuous  $G_K$ -module  $M$  we can define a presheaf  $\mathbb{G}_M$  on  $\mathbf{K}_{\text{ét}}$  such that

- $\mathbb{G}_M(L) = M^H$  where  $H = G_L = \text{Gal}(\mathbb{K}/L)$ .
- $\mathbb{G}_M(\prod L_i) = \prod \mathbb{G}_M(L_i)$ .

One verifies that  $\mathbb{G}_M$  is a sheaf of Abelian groups on  $\mathbf{K}_{\text{ét}}$ .

**THEOREM 3.15.** *There is an equivalence of categories*

$$\mathbf{S}(\mathbf{K}_{\text{ét}}) \xleftrightarrow{\sim} G_K - \text{mod}$$

*induced by the correspondences  $\mathbb{G} \mapsto M_{\mathbb{G}}$  and  $M \mapsto \mathbb{G}_M$ . Here,  $G_K - \text{mod}$  is the category of continuous  $G_K$ -modules.*

**PROOF.** A  $G_K$ -morphism  $M \longrightarrow M'$  induces a morphism of sheaves  $\mathbb{G}_M \longrightarrow \mathbb{G}_{M'}$ . Conversely, if  $H$  is an open subgroup of  $G_K$  with  $L = \mathbb{K}^H$ , then if  $\mathbb{G} \xrightarrow{\phi} \mathbb{G}'$  is a sheafmorphism,  $\phi(L) : \mathbb{G}(L) \longrightarrow \mathbb{G}'(L)$  commutes with the action of  $G_K$  by functoriality of  $\phi$ . Therefore,  $\varinjlim \phi(L)$  is a  $G_K$ -morphism  $M_{\mathbb{G}} \longrightarrow M_{\mathbb{G}'}$ .

One verifies easily that  $\text{Hom}_{G_K}(M, M') \longrightarrow \text{Hom}(\mathbb{G}_M, \mathbb{G}_{M'})$  is an isomorphism and that the canonical map  $\mathbb{G} \longrightarrow \mathbb{G}_{M_{\mathbb{G}}}$  is an isomorphism.  $\square$

In particular, we have that  $\mathbb{G}(K) = \mathbb{G}(\mathbb{K})^{G_K}$  for every sheaf  $\mathbb{G}$  of Abelian groups on  $\mathbf{K}_{\text{ét}}$  and where  $\mathbb{G}(\mathbb{K}) = M_{\mathbb{G}}$ . Hence, the right derived functors of  $\Gamma$  and  $(-)^G$  coincide for Abelian sheaves.

The category  $G_K - \text{mod}$  of continuous  $G_K$ -modules is Abelian having enough injectives. Therefore, the left exact functor

$$(-)^G : G_K - \text{mod} \longrightarrow \text{abelian}$$

admits right derived functors. They are called the *Galois cohomology groups* and denoted

$$R^i M^G = H^i(G_K, M)$$

Therefore, we have.

**PROPOSITION 3.16.** *For any sheaf of Abelian groups  $\mathbb{G}$  on  $\mathbf{K}_{\text{ét}}$  we have a group isomorphism*

$$H_{\text{ét}}^i(K, \mathbb{G}) \simeq H^i(G_K, \mathbb{G}(\mathbb{K}))$$

Therefore, étale cohomology is a natural extension of Galois cohomology to arbitrary algebras.

The following definition-characterization of central simple algebras is classical, see for example [63].

**PROPOSITION 3.17.** *Let  $A$  be a finite dimensional  $K$ -algebra. The following are equivalent :*

- (1)  $A$  has no proper twosided ideals and the center of  $A$  is  $K$ .
- (2)  $A_{\mathbb{K}} = A \otimes_K \mathbb{K} \simeq M_n(\mathbb{K})$  for some  $n$ .
- (3)  $A_L = A \otimes_K L \simeq M_n(L)$  for some  $n$  and some finite Galois extension  $L/K$ .

- (4)  $A \simeq M_k(D)$  for some  $k$  where  $D$  is a division algebra of dimension  $l^2$  with center  $K$ .

The last part of this result suggests the following definition. Call two central simple algebras  $A$  and  $A'$  equivalent if and only if  $A \simeq M_k(\Delta)$  and  $A' \simeq M_l(\Delta)$  with  $\Delta$  a division algebra. From the second characterization it follows that the tensorproduct of two central simple  $K$ -algebras is again central simple. Therefore, we can equip the set of equivalence classes of central simple algebras with a product induced from the tensorproduct. This product has the class  $[K]$  as unit element and  $[\Delta]^{-1} = [\Delta^{opp}]$ , the opposite algebra as  $\Delta \otimes_K \Delta^{opp} \simeq End_K(\Delta) = M_{l^2}(K)$ . This group is called the *Brauer group* and is denoted  $Br(K)$ . We will quickly recall its cohomological description, all of which is classical.

$GL_r$  is an affine smooth algebraic group defined over  $K$  and is the automorphism group of a vectorspace of dimension  $r$ . It defines a sheaf of groups on  $K_{et}$  that we will denote by  $\mathcal{G}L_r$ . Using the fact that the first cohomology classifies twisted forms of vectorspaces of dimension  $r$  we have :

LEMMA 3.18.

$$H_{et}^1(K, \mathcal{G}L_r) = H^1(G_K, GL_r(\mathbb{K})) = 0$$

In particular, we have 'Hilbert's theorem 90'

$$H_{et}^1(K, \mathbb{G}_m) = H^1(G_K, \mathbb{K}^*) = 0$$

PROOF. The cohomology group classifies  $K$ -module isomorphism classes of twisted forms of  $r$ -dimensional vectorspaces over  $K$ . There is just one such class.  $\square$

$PGL_n$  is an affine smooth algebraic group defined over  $K$  and it is the automorphism group of the  $K$ -algebra  $M_n(K)$ . It defines a sheaf of groups on  $K_{et}$  denoted by  $\mathcal{P}GL_n$ . By proposition 3.17 we know that any central simple  $K$ -algebra  $\Delta$  of dimension  $n^2$  is a twisted form of  $M_n(K)$ . Therefore,

LEMMA 3.19. *The pointed set of  $K$ -algebra isomorphism classes of central simple algebras of dimension  $n^2$  over  $K$  coincides with the cohomology set*

$$H_{et}^1(K, \mathcal{P}GL_n) = H^1(G_K, PGL_n(\mathbb{K}))$$

THEOREM 3.20. *There is a natural inclusion*

$$H_{et}^1(K, \mathcal{P}GL_n) \hookrightarrow H_{et}^2(K, \mu_n) = Br_n(K)$$

where  $Br_n(K)$  is the  $n$ -torsion part of the Brauer group of  $K$ . Moreover,

$$Br(K) = H_{et}^2(K, \mathbb{G}_m)$$

is a torsion group.

PROOF. Consider the exact commutative diagram of sheaves of groups on  $K_{et}$  of figure 2. Taking cohomology of the second exact sequence we obtain

$$GL_n(K) \xrightarrow{\det} K^* \longrightarrow H_{et}^1(K, \mathcal{S}L_n) \longrightarrow H_{et}^1(K, \mathcal{G}L_n)$$

where the first map is surjective and the last term is zero, whence

$$H_{et}^1(K, \mathcal{S}L_n) = 0$$

Taking cohomology of the first vertical exact sequence we get

$$H_{et}^1(K, \mathcal{S}L_n) \longrightarrow H_{et}^1(K, \mathcal{P}GL_n) \longrightarrow H_{et}^2(K, \mu_n)$$

from which the first claim follows.




FIGURE 2. Brauer group diagram.

As for the second assertion, taking cohomology of the first exact sequence we get

$$H_{et}^1(K, \mathbb{G}_m) \longrightarrow H_{et}^2(K, \mu_n) \longrightarrow H_{et}^2(K, \mathbb{G}_m) \xrightarrow{n} H_{et}^2(K, \mathbb{G}_m)$$

By Hilbert 90, the first term vanishes and hence  $H_{et}^2(K, \mu_n)$  is equal to the  $n$ -torsion of the group

$$H_{et}^2(K, \mathbb{G}_m) = H^2(G_K, \mathbb{K}^*) = Br(K)$$

where the last equality follows from the crossed product result, see for example [63].  $\square$

So far, the field  $K$  was arbitrary. If  $K$  is of transcendence degree  $d$ , this will put restrictions on the 'size' of the Galois group  $G_K$ . In particular this will enable us to show in section 3.6 that  $H^i(G_K, \mu_n) = 0$  for  $i > d$ .

### 3.3. Quiver orders.

In this section and the next we will construct a large class of central simple algebras controlled by combinatorial data, using the setting of proposition 3.12.

Recall that a *quiver*  $Q$  is a directed graph determined by

- a finite set  $Q_v = \{v_1, \dots, v_k\}$  of *vertices*, and
- a finite set  $Q_a = \{a_1, \dots, a_l\}$  of *arrows* where we allow multiple arrows between vertices and loops in vertices.

Every arrow  $\textcircled{j} \xleftarrow{a} \textcircled{i}$  has a *starting vertex*  $s(a) = i$  and a *terminating vertex*  $t(a) = j$ . Multiplication in the *path algebra*  $\mathbb{C}Q$  is induced by (left) concatenation

of paths. More precisely,  $1 = v_1 + \dots + v_k$  is a decomposition of 1 into mutually orthogonal idempotents and further we define

- $v_j.a$  is always zero unless  $\textcircled{j} \xleftarrow{a} \textcircled{\phantom{j}}$  in which case it is the path  $a$ ,
- $a.v_i$  is always zero unless  $\textcircled{\phantom{i}} \xleftarrow{a} \textcircled{i}$  in which case it is the path  $a$ ,
- $a_i.a_j$  is always zero unless  $\textcircled{\phantom{i}} \xleftarrow{a_i} \textcircled{\phantom{j}} \xleftarrow{a_j} \textcircled{\phantom{i}}$  in which case it is the path  $a_i a_j$ .

Path algebras of quivers are the archetypical examples of *Quillen smooth algebras* which are the basic building blocks to construct noncommutative manifolds.

DEFINITION 3.21. A  $\mathbb{C}$ -algebra  $A$  is said to be *Quillen smooth* if it satisfies the following lifting property. Let  $T$  be a  $\mathbb{C}$ -algebra and  $I \triangleleft T$  a nilpotent ideal. If there is a  $\mathbb{C}$ -algebra morphism  $A \xrightarrow{\kappa} T/I$  then there exists a  $\mathbb{C}$ -algebra lift  $A \xrightarrow{\lambda} T$

$$\begin{array}{ccc} & A & \\ & \swarrow \exists \lambda & \downarrow \kappa \\ T & \longrightarrow & T/I \end{array}$$

making the diagram commutative.

This definition is rather restrictive. In particular, a commutative smooth algebra does not have to satisfy this lifting property.

EXAMPLE 3.22. consider the polynomial algebra  $\mathbb{C}[x_1, \dots, x_d]$  and the 4-dimensional non-commutative local algebra

$$T = \frac{\mathbb{C}\langle x, y \rangle}{(x^2, y^2, xy + yx)} = \mathbb{C} \oplus \mathbb{C}x \oplus \mathbb{C}y \oplus \mathbb{C}xy$$

Consider the one-dimensional nilpotent ideal  $I = \mathbb{C}(xy - yx)$  of  $T$ , then the 3-dimensional quotient  $\frac{T}{I}$  is commutative and we have a morphism  $\mathbb{C}[x_1, \dots, x_d] \xrightarrow{\phi} \frac{T}{I}$  by  $x_1 \mapsto x, x_2 \mapsto y$  and  $x_i \mapsto 0$  for  $i \geq 2$ . This morphism admits no lift to  $T$  as for any potential lift the commutator

$$[\tilde{\phi}(x), \tilde{\phi}(y)] \neq 0 \quad \text{in } T.$$

Therefore,  $\mathbb{C}[x_1, \dots, x_d]$  can only be Quillen smooth if  $d = 1$ .

Consider the commutative  $\mathbb{C}$ -algebra

$$C_k = \mathbb{C}[e_1, \dots, e_k] / (e_i^2 - e_i, e_i e_j, \sum_{i=1}^k e_i - 1).$$

$C_k$  is the universal  $\mathbb{C}$ -algebra in which 1 is decomposed into  $k$  orthogonal idempotents, that is, if  $R$  is any  $\mathbb{C}$ -algebra such that  $1 = r_1 + \dots + r_k$  with  $r_i \in R$  idempotents satisfying  $r_i r_j = 0$ , then there is an embedding  $C_k \hookrightarrow R$  sending  $e_i$  to  $r_i$ .

PROPOSITION 3.23.  $C_k$  is Quillen smooth. That is, if  $I$  be a nilpotent ideal of a  $\mathbb{C}$ -algebra  $T$  and if  $1 = \bar{e}_1 + \dots + \bar{e}_k$  is a decomposition of 1 into orthogonal idempotents  $\bar{e}_i \in T/I$ . Then, we can lift this decomposition to  $1 = e_1 + \dots + e_k$  for orthogonal idempotents  $e_i \in T$  such that  $\pi(e_i) = \bar{e}_i$  where  $T \xrightarrow{\pi} T/I$  is the canonical projection.

PROOF. Assume that  $I^l = 0$ , clearly any element  $1 - i$  with  $i \in I$  is invertible in  $T$  as

$$(1 - i)(1 + i + i^2 + \dots + i^{l-1}) = 1 - i^l = 1.$$

If  $\bar{e}$  is an idempotent of  $T/I$  and  $x \in T$  such that  $\pi(x) = \bar{e}$ . Then,  $x - x^2 \in I$  whence

$$0 = (x - x^2)^l = x^l - lx^{l+1} + \binom{l}{2}x^{l+2} - \dots + (-1)^l x^{2l}$$

and therefore  $x^l = ax^{l+1}$  where  $a = l - \binom{l}{2}x + \dots + (-1)^{l-1}x^{l-1}$  and so  $ax = xa$ .

If we take  $e = (ax)^l$ , then  $e$  is an idempotent in  $T$  as

$$e^2 = (ax)^{2l} = a^l(a^l x^{2l}) = a^l x^l = e$$

the next to last equality follows from  $x^l = ax^{l+1} = a^2 x^{l+2} = \dots = a^l x^{2l}$ . Moreover,

$$\pi(e) = \pi(a)^l \pi(x)^l = \pi(a)^l \pi(x)^{2l} = \pi(a^l x^{2l}) = \pi(x)^l = \bar{e}.$$

If  $\bar{f}$  is another idempotent in  $T/I$  such that  $\bar{e}\bar{f} = 0 = \bar{f}\bar{e}$  then as above we can lift  $\bar{f}$  to an idempotent  $f'$  of  $T$ . As  $f'e \in I$  we can form the element

$$f = (1 - e)(1 - f'e)^{-1}f'(1 - f'e).$$

Because  $f'(1 - f'e) = f'(1 - e)$  one verifies that  $f$  is idempotent,  $\pi(f) = \bar{f}$  and  $e.f = 0 = f.e$ . Assume by induction that we have already lifted the pairwise orthogonal idempotents  $\bar{e}_1, \dots, \bar{e}_{k-1}$  to pairwise orthogonal idempotents  $e_1, \dots, e_{k-1}$  of  $R$ , then  $e = e_1 + \dots + e_{k-1}$  is an idempotent of  $T$  such that  $\bar{e}\bar{e}_k = 0 = \bar{e}_k\bar{e}$ . Hence, we can lift  $\bar{e}_k$  to an idempotent  $e_k \in T$  such that  $ee_k = 0 = e_k e$ . But then also

$$e_i e_k = (e_i e) e_k = 0 = e_k (e e_i) = e_k e_i.$$

Finally, as  $e_1 + \dots + e_k - 1 = i \in I$  we have that

$$e_1 + \dots + e_k - 1 = (e_1 + \dots + e_k - 1)^l = i^l = 0$$

finishing the proof.  $\square$

PROPOSITION 3.24. *For any quiver  $Q$ , the path algebra  $\mathbb{C}Q$  is Quillen smooth.*

PROOF. Take an algebra  $T$  with a nilpotent twosided ideal  $I \triangleleft T$  and consider

$$\begin{array}{ccc} T & \xrightarrow{\quad} & \frac{T}{I} \\ & \searrow \tilde{\phi} & \uparrow \phi \\ & & \mathbb{C}Q \end{array}$$

The decomposition  $1 = \phi(v_1) + \dots + \phi(v_k)$  into mutually orthogonal idempotents in  $\frac{T}{I}$  can be lifted up the nilpotent ideal  $I$  to a decomposition  $1 = \tilde{\phi}(v_1) + \dots + \tilde{\phi}(v_k)$  into mutually orthogonal idempotents in  $T$ . But then, taking for every arrow  $a$


$$\textcircled{j} \xleftarrow{a} \textcircled{i} \quad \text{an arbitrary element} \quad \tilde{\phi}(a) \in \tilde{\phi}(v_j)(\phi(a) + I)\tilde{\phi}(v_i)$$

gives a required lifted algebra morphism  $\mathbb{C}Q \xrightarrow{\tilde{\phi}} T$ .  $\square$

The description of the quiver  $Q$  can be encoded in an integral  $k \times k$  matrix

$$\chi_Q = \begin{bmatrix} \chi_{11} & \cdots & \chi_{1k} \\ \vdots & & \vdots \\ \chi_{k1} & \cdots & \chi_{kk} \end{bmatrix} \quad \text{with} \quad \chi_{ij} = \delta_{ij} - \# \{ \textcircled{i} \leftarrow \textcircled{i} \}$$

EXAMPLE 3.25. Consider the quiver  $Q$



Then, with the indicated ordering of the vertices we have that the integral matrix is

$$\chi_Q = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & -1 \\ 0 & 0 & 0 \end{bmatrix}$$

and the path algebra of  $Q$  is isomorphic to the block-matrix algebra

$$\mathbb{C}Q' \simeq \begin{bmatrix} \mathbb{C} & \mathbb{C} \oplus \mathbb{C} & 0 \\ 0 & \mathbb{C} & 0 \\ 0 & \mathbb{C}[x] & \mathbb{C}[x] \end{bmatrix}$$

where  $x$  is the loop in vertex  $v_3$ .

The subspace  $\mathbb{C}Qv_i$  has as basis the paths starting in vertex  $v_i$  and because  $\mathbb{C}Q = \bigoplus_i \mathbb{C}Qv_i$ ,  $\mathbb{C}Qv_i$  is a projective left ideal of  $\mathbb{C}Q$ . Similarly,  $v_i\mathbb{C}Q$  has as basis the paths ending at  $v_i$  and is a projective right ideal of  $\mathbb{C}Q$ . The subspace  $v_i\mathbb{C}Qv_j$  has as basis the paths starting at  $v_j$  and ending at  $v_i$  and  $\mathbb{C}Qv_i\mathbb{C}Q$  is the twosided ideal of  $\mathbb{C}Q$  having as basis all paths passing through  $v_i$ . If  $0 \neq f \in \mathbb{C}Qv_i$  and  $0 \neq g \in v_i\mathbb{C}Q$ , then  $f.g \neq 0$  for let  $p$  be a longest path occurring in  $f$  and  $q$  a longest path in  $g$ , then the coefficient of  $p.q$  in  $f.g$  cannot be zero. As a consequence we have

LEMMA 3.26. *The projective left ideals  $\mathbb{C}Qv_i$  are indecomposable and pairwise non-isomorphic.*

PROOF. If  $\mathbb{C}Qv_i$  is not indecomposable, then there exists a projection idempotent  $f \in \text{Hom}_{\mathbb{C}Q}(\mathbb{C}Qv_i, \mathbb{C}Qv_i) \simeq v_i\mathbb{C}Qv_i$ . But then,  $f^2 = f = f.v_i$  whence  $f.(f - v_i) = 0$ , contradicting the remark above. Further, for any left  $\mathbb{C}Q$ -module  $M$  we have that  $\text{Hom}_{\mathbb{C}Q}(\mathbb{C}Qv_i, M) \simeq v_iM$ . So, if  $\mathbb{C}Qv_i \simeq \mathbb{C}Qv_j$  then the isomorphism gives elements  $f \in v_i\mathbb{C}Qv_j$  and  $g \in v_j\mathbb{C}Qv_i$  such that  $f.g = v_i$  and  $g.f = v_j$ . But then,  $v_i \in \mathbb{C}Qv_j\mathbb{C}Q$ , a contradiction unless  $i = j$  as this space has basis all paths passing through  $v_j$ .  $\square$

EXAMPLE 3.27. Let  $Q$  be a quiver, then the following properties hold :

- (1)  $\mathbb{C}Q$  is finite dimensional if and only if  $Q$  has no oriented cycles.
- (2)  $\mathbb{C}Q$  is prime (that is,  $I.J \neq 0$  for all twosided ideals  $I, J \neq 0$ ) if and only if  $Q$  is *strongly connected*, that is, for all vertices  $v_i$  and  $v_j$  there is a path from  $v_i$  to  $v_j$ .
- (3)  $\mathbb{C}Q$  is Noetherian (that is, satisfies the ascending chain condition on left (or right) ideals) if and only if for every vertex  $v_i$  belonging to an oriented cycle there is only one arrow starting at  $v_i$  and only one arrow terminating at  $v_i$ .
- (4) The radical of  $\mathbb{C}Q$  has as basis all paths from  $v_i$  to  $v_j$  for which there is no path from  $v_j$  to  $v_i$ .
- (5) The center of  $\mathbb{C}Q$  is of the form  $\mathbb{C} \times \dots \times \mathbb{C} \times \mathbb{C}[x] \times \dots \times \mathbb{C}[x]$  with one factor for each connected component  $C$  of  $Q$  (that is, connected component for the underlying graph forgetting the orientation) and this factor is isomorphic to  $\mathbb{C}[x]$  if and only if  $C$  is one oriented cycle.



Recall that a *representation*  $V$  of the quiver  $Q$  is given by

- a finite dimensional  $\mathbb{C}$ -vector space  $V_i$  for each vertex  $v_i \in Q_v$ , and
- a linear map  $V_j \xleftarrow{V_a} V_i$  for every arrow  $\textcircled{j} \xleftarrow{a} \textcircled{i}$  in  $Q_a$ .

If  $\dim V_i = d_i$  we call the integral vector  $\alpha = (d_1, \dots, d_k) \in \mathbb{N}^k$  the *dimension vector* of  $V$  and denote it with  $\dim V$ . A *morphism*  $V \xrightarrow{\phi} W$  between two representations  $V$  and  $W$  of  $Q$  is determined by a set of linear maps

$$V_i \xrightarrow{\phi_i} W_i \quad \text{for all vertices } v_i \in Q_v$$

satisfying the following compatibility conditions for every arrow  $\textcircled{j} \xleftarrow{a} \textcircled{i}$  in  $Q_a$

$$\begin{array}{ccc} V_i & \xrightarrow{V_a} & V_j \\ \downarrow \phi_i & & \downarrow \phi_j \\ W_i & \xrightarrow{W_a} & W_j \end{array}$$

Clearly, *composition* of morphisms  $V \xrightarrow{\phi} W \xrightarrow{\psi} X$  is given by the rule that  $(\psi \circ \phi)_i = \psi_i \circ \phi_i$  and one readily verifies that this is again a morphism of representations of  $Q$ . In this way we form a *category*  $\text{rep } Q$  of all finite dimensional representations of the quiver  $Q$ .

**PROPOSITION 3.28.** *The category  $\text{rep } Q$  is equivalent to the category of finite dimensional  $\mathbb{C}Q$ -representations  $\mathbb{C}Q\text{-mod}$ .*

**PROOF.** Let  $M$  be an  $n$ -dimensional  $\mathbb{C}Q$ -representation. Then, we construct a representation  $V$  of  $Q$  by taking

- $V_i = v_i M$ , and for any arrow  $\textcircled{j} \xleftarrow{a} \textcircled{i}$  in  $Q_a$  define
- $V_a : V_i \longrightarrow V_j$  by  $V_a(x) = v_j a x$ .

Observe that the dimension vector  $\dim(V) = (d_1, \dots, d_k)$  satisfies  $\sum d_i = n$ . If  $\phi : M \longrightarrow N$  is  $\mathbb{C}Q$ -linear, then we have a linear map  $V_i = v_i M \xrightarrow{\phi_i} W_i = v_i N$  which clearly satisfies the compatibility condition.

Conversely, let  $V$  be a representation of  $Q$  with dimension vector  $\dim(V) = (d_1, \dots, d_k)$ . Then, consider the  $n = \sum d_i$ -dimensional space  $M = \bigoplus_i V_i$  which we turn into a  $\mathbb{C}Q$ -representation as follows. Consider the canonical injection and projection maps  $V_j \xrightarrow{i_j} M \xrightarrow{\pi_j} V_j$ . Then, define the action of  $\mathbb{C}Q$  by fixing the action of the algebra generators  $v_j$  and  $a_l$  to be

$$\begin{cases} v_j m &= i_j(\pi_j(m)) \\ a_l m &= i_j(V_a(\pi_i(m))) \end{cases}$$

for all arrows  $\textcircled{j} \xleftarrow{a_l} \textcircled{i}$ . A computation verifies that these two operations are inverse to each other and induce an equivalence of categories.  $\square$

Let  $A$  be a  $\mathbb{C}$ -algebra and let  $M$  and  $N$  be two  $A$ -representations of dimensions say  $m$  and  $n$ . An  $A$ -representation  $P$  of dimension  $m+n$  is said to be an *extension of  $N$  by  $M$*  if there exists a short exact sequence of left  $A$ -modules

$$e : \quad 0 \longrightarrow M \longrightarrow P \longrightarrow N \longrightarrow 0$$

We define an equivalence relation on extensions  $(P, e)$  of  $N$  by  $M : (P, e) \cong (P', e')$  if and only if there is an isomorphism  $P \xrightarrow{\phi} P'$  of left  $A$ -modules such that the diagram below is commutative

$$\begin{array}{ccccccccc}
 e : & 0 & \longrightarrow & M & \longrightarrow & P & \longrightarrow & N & \longrightarrow & 0 \\
 & & & \downarrow \text{id}_M & & \downarrow \phi & & \downarrow \text{id}_N & & \\
 e' : & 0 & \longrightarrow & M & \longrightarrow & P' & \longrightarrow & N & \longrightarrow & 0
 \end{array}$$

The set of equivalence classes of extensions of  $N$  by  $M$  will be denoted by  $Ext_A^1(N, M)$ .

An alternative description of  $Ext_A^1(N, M)$  is as follows. Let  $\rho : A \longrightarrow M_m$  and  $\sigma : A \longrightarrow M_n$  be the representations defining  $M$  and  $N$ . For an extension  $(P, e)$  we can identify the  $\mathbb{C}$ -vectorspace with  $M \oplus N$  and the  $A$ -module structure on  $P$  gives a algebra map  $\mu : A \longrightarrow M_{m+n}$  and we can represent the action of  $a$  on  $P$  by left multiplication of the block-matrix

$$\mu(a) = \begin{bmatrix} \rho(a) & \lambda(a) \\ 0 & \sigma(a) \end{bmatrix},$$

where  $\lambda(a)$  is an  $m \times n$  matrix and hence defines a linear map

$$\lambda : A \longrightarrow Hom_{\mathbb{C}}(N, M).$$

The condition that  $\mu$  is an algebra morphism is equivalent to the condition

$$\lambda(aa') = \rho(a)\lambda(a') + \lambda(a)\sigma(a')$$

and we denote the set of all liner maps  $\lambda : A \longrightarrow Hom_{\mathbb{C}}(N, M)$  by  $Z(N, M)$  and call it the space of *cycle*. The extensions of  $N$  by  $M$  corresponding to two cycles  $\lambda$  and  $\lambda'$  from  $Z(N, M)$  are equivalent if and only if we have an  $A$ -module isomorphism in block form

$$\begin{bmatrix} \text{id}_M & \beta \\ 0 & \text{id}_N \end{bmatrix} \quad \text{with } \beta \in Hom_{\mathbb{C}}(N, M)$$

between them.  $A$ -linearity of this map translates into the matrix relation

$$\begin{bmatrix} \text{id}_M & \beta \\ 0 & \text{id}_N \end{bmatrix} \cdot \begin{bmatrix} \rho(a) & \lambda(a) \\ 0 & \sigma(a) \end{bmatrix} = \begin{bmatrix} \rho(a) & \lambda'(a) \\ 0 & \sigma(a) \end{bmatrix} \cdot \begin{bmatrix} \text{id}_M & \beta \\ 0 & \text{id}_N \end{bmatrix} \quad \text{for all } a \in A$$

or equivalently, that  $\lambda(a) - \lambda'(a) = \rho(a)\beta - \beta\sigma(a)$  for all  $a \in A$ . We will now define the subspace of  $Z(N, M)$  of *boundaries*  $B(N, M)$

$$\{\delta \in Hom_{\mathbb{C}}(N, M) \mid \exists \beta \in Hom_{\mathbb{C}}(N, M) : \forall a \in A : \delta(a) = \rho(a)\beta - \beta\sigma(a)\}.$$

We then have the description  $Ext_A^1(N, M) = \frac{Z(N, M)}{B(N, M)}$ .

The *Euler form* of the quiver  $Q$  is the bilinear form on  $\mathbb{Z}^k$

$$\chi_Q(\cdot, \cdot) : \mathbb{Z}^k \times \mathbb{Z}^k \longrightarrow \mathbb{Z} \quad \text{defined by } \chi_Q(\alpha, \beta) = \alpha \cdot \chi_Q \cdot \beta^T$$

for all row vectors  $\alpha, \beta \in \mathbb{Z}^k$ .

**THEOREM 3.29.** *Let  $V$  and  $W$  be two representations of  $Q$ , then*

$$\dim_{\mathbb{C}} Hom_{\mathbb{C}Q}(V, W) - \dim_{\mathbb{C}} Ext_{\mathbb{C}Q}^1(V, W) = \chi_Q(\dim(V), \dim(W))$$

PROOF. We claim that there exists an exact sequence of  $\mathbb{C}$ -vectorspaces

$$0 \longrightarrow \text{Hom}_{\mathbb{C}Q}(V, W) \xrightarrow{\gamma} \bigoplus_{v_i \in Q_v} \text{Hom}_{\mathbb{C}}(V_i, W_i) \xrightarrow{d_W^V} \\ \xrightarrow{d_W^V} \bigoplus_{a \in Q_a} \text{Hom}_{\mathbb{C}}(V_{s(a)}, W_{t(a)}) \xrightarrow{\epsilon} \text{Ext}_{\mathbb{C}Q}^1(V, W) \longrightarrow 0$$

Here,  $\gamma(\phi) = (\phi_1, \dots, \phi_k)$  and  $d_W^V$  maps a family of linear maps  $(f_1, \dots, f_k)$  to the linear maps  $\mu_a = f_{t(a)}V_a - W_a f_{s(a)}$  for any arrow  $a$  in  $Q$ , that is, to the obstruction of the following diagram to be commutative

$$\begin{array}{ccc} V_{s(a)} & \xrightarrow{V_a} & V_{t(a)} \\ \downarrow f_{s(a)} & \searrow \mu_a & \downarrow f_{t(a)} \\ W_{s(a)} & \xrightarrow{W_a} & W_{t(a)} \end{array}$$

By the definition of morphisms between representations of  $Q$  it is clear that the kernel of  $d_W^V$  coincides with  $\text{Hom}_{\mathbb{C}Q}(V, W)$ .

Further, the map  $\epsilon$  is defined by sending a family of maps  $(g_1, \dots, g_s) = (g_a)_{a \in Q_a}$  to the equivalence class of the exact sequence

$$0 \longrightarrow W \xrightarrow{i} E \xrightarrow{p} V \longrightarrow 0$$

where for all  $v_i \in Q_v$  we have  $E_i = W_i \oplus V_i$  and the inclusion  $i$  and projection map  $p$  are the obvious ones and for each generator  $a \in Q_a$  the action of  $a$  on  $E$  is defined by the matrix

$$E_a = \begin{bmatrix} W_a & g_a \\ 0 & V_a \end{bmatrix} : E_{s(a)} = W_{s(a)} \oplus V_{s(a)} \longrightarrow W_{t(a)} \oplus V_{t(a)} = E_{t(a)}$$

Clearly, this makes  $E$  into a  $\mathbb{C}Q$ -module and one verifies that the above short exact sequence is one of  $\mathbb{C}Q$ -modules. Remains to prove that the cokernel of  $d_W^V$  can be identified with  $\text{Ext}_{\mathbb{C}Q}^1(V, W)$ .

A set of algebra generators of  $\mathbb{C}Q$  is given by  $\{v_1, \dots, v_k, a_1, \dots, a_l\}$ . A cycle is given by a linear map  $\lambda : \mathbb{C}Q \longrightarrow \text{Hom}_{\mathbb{C}}(V, W)$  such that for all  $f, f' \in \mathbb{C}Q$  we have the condition

$$\lambda(ff') = \rho(f)\lambda(f') + \lambda(f)\sigma(f')$$

where  $\rho$  determines the action on  $W$  and  $\sigma$  that on  $V$ . First, consider  $v_i$  then the condition says  $\lambda(v_i^2) = \lambda(v_i) = p_i^W \lambda(v_i) + \lambda(v_i) p_i^V$  whence  $\lambda(v_i) : V_i \longrightarrow W_i$  but then applying again the condition we see that  $\lambda(v_i) = 2\lambda(v_i)$  so  $\lambda(v_i) = 0$ . Similarly, using the condition on  $a = v_{t(a)}a = av_{s(a)}$  we deduce that  $\lambda(a) : V_{s(a)} \longrightarrow W_{t(a)}$ . That is, we can identify  $\bigoplus_{a \in Q_a} \text{Hom}_{\mathbb{C}}(V_{s(a)}, W_{t(a)})$  with  $Z(V, W)$  under the map  $\epsilon$ . Moreover, the image of  $\delta$  gives rise to a family of morphisms  $\lambda(a) = f_{t(a)}V_a - W_a f_{s(a)}$  for a linear map  $f = (f_i) : V \longrightarrow W$  so this image coincides precisely to the subspace of boundaries  $B(V, W)$  proving that indeed the cokernel of  $d_W^V$  is  $\text{Ext}_{\mathbb{C}Q}^1(V, W)$  finishing the proof of exactness of the long sequence of vectorspaces. But then, if  $\dim(V) = (r_1, \dots, r_k)$  and  $\dim(W) = (s_1, \dots, s_k)$ , we have that

$\dim \operatorname{Hom}(V, W) - \dim \operatorname{Ext}^1(V, W)$  is equal to

$$\begin{aligned} & \sum_{v_i \in Q_v} \dim \operatorname{Hom}_{\mathbb{C}}(V_i, W_i) - \sum_{a \in Q_a} \dim \operatorname{Hom}_{\mathbb{C}}(V_{s(a)}, W_{t(a)}) \\ &= \sum_{v_i \in Q_v} r_i s_i - \sum_{a \in Q_a} r_{s(a)} s_{t(a)} \\ &= (r_1, \dots, r_k) M_Q (s_1, \dots, s_k)^{\tau} = \chi_Q(\dim(V), \dim(W)) \end{aligned}$$

finishing the proof.  $\square$

Fix a dimension vector  $\alpha = (d_1, \dots, d_k) \in \mathbb{N}^k$  and consider the set  $\operatorname{rep}_{\alpha} Q$  of all representations  $V$  of  $Q$  such that  $\dim(V) = \alpha$ . Because  $V$  is completely determined by the linear maps

$$V_a : V_{s(a)} = \mathbb{C}^{d_{s(a)}} \longrightarrow \mathbb{C}^{d_{t(a)}} = V_{t(a)}$$

we see that  $\operatorname{rep}_{\alpha} Q$  is the affine space

$$\operatorname{rep}_{\alpha} Q = \bigoplus_{\textcircled{j} \xleftarrow{a} \textcircled{i}} M_{d_j \times d_i}(\mathbb{C}) \simeq \mathbb{C}^r$$

where  $r = \sum_{a \in Q_a} d_{s(a)} d_{t(a)}$ . On this affine space we have an action of the algebraic group  $GL(\alpha) = GL_{d_1} \times \dots \times GL_{d_k}$  by conjugation. That is, if  $g = (g_1, \dots, g_k) \in GL(\alpha)$  and if  $V = (V_a)_{a \in Q_a}$  then  $g.V$  is determined by the matrices

$$(g.V)_a = g_{t(a)} V_a g_{s(a)}^{-1}.$$

If  $V$  and  $W$  in  $\operatorname{rep}_{\alpha} Q$  are isomorphic as representations of  $Q$ , such an isomorphism is determined by invertible matrices  $g_i : V_i \longrightarrow W_i \in GL_{d_i}$  such that for every arrow  $\textcircled{j} \xleftarrow{a} \textcircled{i}$  we have a commutative diagram

$$\begin{array}{ccc} V_i & \xrightarrow{V_a} & V_j \\ \downarrow g_i & & \downarrow g_j \\ W_i & \xrightarrow{W_a} & W_j \end{array}$$

or equivalently,  $g_j V_a = W_a g_i$ . That is, two representations are isomorphic if and only if they belong to the same orbit under  $GL(\alpha)$ . In particular, we see that

$$\operatorname{Stab}_{GL(\alpha)} V \simeq \operatorname{Aut}_{\mathbb{C}Q} V$$

and the latter is an open subvariety of the affine space  $\operatorname{End}_{\mathbb{C}Q}(V) = \operatorname{Hom}_{\mathbb{C}Q}(V, V)$  whence they have the same dimension. The dimension of the orbit  $\mathcal{O}(V)$  of  $V$  in  $\operatorname{rep}_{\alpha} Q$  is equal to

$$\dim \mathcal{O}(V) = \dim GL(\alpha) - \dim \operatorname{Stab}_{GL(\alpha)} V.$$

But then we have a geometric reformulation of the above theorem.

LEMMA 3.30. *Let  $V \in \operatorname{rep}_{\alpha} Q$ , then*

$$\dim \operatorname{rep}_{\alpha} Q - \dim \mathcal{O}(V) = \dim \operatorname{End}_{\mathbb{C}Q}(V) - \chi_Q(\alpha, \alpha) = \dim \operatorname{Ext}_{\mathbb{C}Q}^1(V, V)$$

PROOF. We have seen that  $\dim \text{rep}_\alpha Q - \dim \mathcal{O}(V)$  is equal to

$$\sum_a d_{s(a)} d_{t(a)} - \left( \sum_i d_i^2 - \dim \text{End}_{\mathbb{C}Q}(V) \right) = \dim \text{End}_{\mathbb{C}Q}(V) - \chi_Q(\alpha, \alpha)$$

and the foregoing theorem asserts that the latter term is equal to  $\dim \text{Ext}_{\mathbb{C}Q}^1(V, V)$ .  $\square$

In particular it follows that the orbit  $\mathcal{O}(V)$  is *open* in  $\text{rep}_\alpha Q$  if and only if  $V$  has no self-extensions. Moreover, as  $\text{rep}_\alpha Q$  is irreducible there can be at most one isomorphism class of a representation without self-extensions.

For every dimension vector  $\alpha = (d_1, \dots, d_k)$  we will construct a *quiver order*  $\mathbb{T}_\alpha Q$  which is a Cayley-Hamilton algebra of degree  $n$  where  $n = d_1 + \dots + d_k$ . First, we describe the  $n$ -dimensional representations of the Quillen-smooth algebra  $C_k$ .

PROPOSITION 3.31. *Let  $C_k = \mathbb{C}[e_1, \dots, e_k]/(e_i^2 - e_i, e_i e_j, \sum_{i=1}^k e_i - 1)$ , then  $\underline{\text{rep}}_n C_k$  is reduced and is the disjoint union of the homogeneous varieties*

$$\underline{\text{rep}}_n C_k = \bigcup_{\alpha} GL_n / (GL_{d_1} \times \dots \times GL_{d_k})$$

where the union is taken over all  $\alpha = (d_1, \dots, d_k)$  such that  $n = \sum_i d_i$ .

PROOF. As  $C_k$  is Quillen smooth we will see in section 4.4 that all its representation spaces  $\underline{\text{rep}}_n C_k$  are smooth varieties hence in particular reduced. Therefore, it suffices to describe the points. For any  $n$ -dimensional representation

$$C_k \xrightarrow{\phi} M_n(\mathbb{C})$$

the image is a commutative semi-simple algebra with orthogonal idempotents  $f_i = \phi(e_i)$  of rank  $d_i$ . Because  $\sum_i e_i = \mathbb{1}_n$  we must have that  $\sum_i d_i = n$ . Alternatively, the corresponding  $n$ -dimensional representation  $M = \bigoplus_i M_i$  where  $M_i = e_i \mathbb{C}^n$  has dimension  $d_i$ . The stabilizer subgroup of  $M$  is equal to  $GL(\alpha) = GL_{d_1} \times \dots \times GL_{d_k}$ , proving the claim.  $\square$

The algebra embedding  $C_k \xrightarrow{\phi} \mathbb{C}Q$  obtained by  $\phi(e_i) = v_i$  determines a morphism

$$\underline{\text{rep}}_n \mathbb{C}Q \xrightarrow{\pi} \underline{\text{rep}}_n C_k = \bigcup_{\alpha} \mathcal{O}(\alpha) = \bigcup_{\alpha} GL_n / GL(\alpha)$$

where the disjoint union is taken over all the dimension vectors  $\alpha = (d_1, \dots, d_k)$  such that  $n = \sum d_i$ . Consider the point  $p_\alpha \in \mathcal{O}(\alpha)$  determined by sending the idempotents  $e_i$  to the canonical diagonal idempotents

$$\sum_{j=\sum_{i=1}^{i-1} d_i+1}^{\sum_{i=1}^i d_i} e_{jj} \in M_n(\mathbb{C})$$

We denote by  $C_k(\alpha)$  this semi-simple commutative subalgebra of  $M_n(\mathbb{C})$ . As  $\text{rep}_\alpha Q$  can be identified with the variety of  $n$ -dimensional representations of  $\mathbb{C}Q$  in block form determined by these idempotents we see that  $\text{rep}_\alpha Q = \pi^{-1}(p)$ .

We define the *quiver trace algebra*  $\mathbb{T}Q$  to be the path algebra of  $Q$  over the polynomial algebra  $R$  in the variables  $t_p$  where  $p$  is a word in the arrows  $a_j \in Q_a$  and is determined only up to cyclic permutation. As a consequence we only retain the variables  $t_p$  where  $p$  is an oriented cycle in  $Q$  (as all the others have a cyclic

permutation which is the zero element in  $\mathbb{C}Q$ ). We define a formal trace map  $tr$  on  $\mathbb{T}Q$  by  $tr(p) = t_p$  if  $p$  is an oriented cycle in  $Q$  and  $tr(p) = 0$  otherwise.

For a fixed dimension vector  $\alpha = (d_1, \dots, d_k)$  with  $\sum_i d_i = n$  we define  $\mathbb{T}_\alpha Q$  to be the quotient

$$\mathbb{T}_\alpha Q = \frac{\mathbb{T}Q}{(\chi_a^{(n)}(a), tr(v_i) - d_i)}$$

by dividing out the substitution invariant twosided ideal generated by all the evaluations of the formal Cayley-Hamilton algebras of degree  $n$ ,  $\chi_a^{(n)}(a)$  for  $a \in \mathbb{T}Q$  together with the additional relations that  $tr(v_i) = d_i$ .  $\mathbb{T}_\alpha Q$  is a Cayley-Hamilton algebra of degree  $n$  with a decomposition  $1 = e_1 + \dots + e_k$  into orthogonal idempotents such that  $tr(e_i) = d_i$ .

More generally, let  $A$  be a Cayley-Hamilton algebra of degree  $n$  with decomposition  $1 = a_1 + \dots + a_n$  into orthogonal idempotents such that  $tr(a_i) = d_i \in \mathbb{N}_+$  and  $\sum d_i = n$ . Then, we have a trace preserving embedding  $C_k(\alpha) \xrightarrow{i} A$  making  $A$  into a  $C_k(\alpha) = \times_{i=1}^k \mathbb{C}$ -algebra. We have a trace preserving embedding  $C_k(\alpha) \xrightarrow{i'} M_n(\mathbb{C})$  by sending the idempotent  $e_i$  to the diagonal idempotent  $E_i \in M_n(\mathbb{C})$  with ones on the diagonal from position  $\sum_{j=1}^{i-1} d_j + 1$  to  $\sum_{j=1}^i d_j$ . This calls for the introduction of a *restricted representation space* of all trace preserving algebra morphisms  $\chi$  such that the diagram below is commutative

$$\begin{array}{ccc} A & \xrightarrow{\chi} & M_n(\mathbb{C}) \\ \uparrow i & \nearrow \chi & \\ C_k(\alpha) & & \end{array}$$

that is, such that  $\chi(a_i) = E_i$ . This again determines an affine scheme  $\underline{rep}_\alpha^{res} A$  which is in fact a closed subscheme of  $\underline{rep}_n^{tr} A$ . The functorial description of the restricted module scheme is as follows. Let  $C$  be any commutative  $\mathbb{C}$ -algebra, then  $M_n(C)$  is a  $C_k(\alpha)$ -algebra and the idempotents  $E_i$  allow for a block decomposition

$$M_n(C) = \oplus_{i,j} E_i M_n(C) E_j = \begin{bmatrix} E_1 M_n(C) E_1 & \dots & E_1 M_n(C) E_k \\ \vdots & & \vdots \\ E_k M_n(C) E_1 & \dots & E_k M_n(C) E_k \end{bmatrix}.$$

The scheme  $\underline{rep}_\alpha^{res} A$  assigns to the algebra  $C$  the set of all trace preserving algebra maps

$$A \xrightarrow{\phi} M_n(B) \quad \text{such that } \phi(a_i) = E_i.$$

Equivalently, the idempotents  $a_i$  decompose  $A$  into block form  $A = \oplus_{i,j} a_i A a_j$  and then  $\underline{rep}_\alpha^{res} A(C)$  are the trace preserving algebra morphisms  $A \longrightarrow M_n(B)$  compatible with the block decompositions.

Still another description of the restricted representation scheme is therefore that  $\underline{rep}_\alpha^{res} A$  is the scheme theoretic fiber  $\pi^{-1}(p_\alpha)$  of the point  $p_\alpha$  under the  $GL_n$ -equivariant morphism

$$\underline{rep}_n^{tr} A \xrightarrow{\pi} \underline{rep}_n^{tr} C_k(\alpha).$$

Hence, the stabilizer subgroup of  $p$  acts on  $\underline{rep}_\alpha^{res} A$ . This stabilizer is the subgroup  $GL(\alpha) = GL_{m_1} \times \dots \times GL_{m_k}$  embedded in  $GL_n$  along the diagonal

$$GL(\alpha) = \begin{bmatrix} GL_{m_1} & & \\ & \ddots & \\ & & GL_{m_k} \end{bmatrix} \hookrightarrow GL_n.$$

Clearly,  $GL(\alpha)$  acts via this embedding by conjugation on  $M_n(\mathbb{C})$ .

**THEOREM 3.32.** *Let  $A$  be a Cayley-Hamilton algebra of degree  $n$  such that  $1 = a_1 + \dots + a_k$  is a decomposition into orthogonal idempotents with  $tr(a_i) = m_i \in \mathbb{N}_+$ . Then,  $A$  is isomorphic to the ring of  $GL(\alpha)$ -equivariant maps*

$$\underline{rep}_\alpha^{res} A \longrightarrow M_n.$$

**PROOF.** We know that  $A$  is the ring of  $GL_n$ -equivariant maps  $\underline{rep}_n^{tr} A \longrightarrow M_n$ . Further, we have a  $GL_n$ -equivariant map

$$\underline{rep}_n^{tr} A \xrightarrow{\pi} \underline{rep}_n tr C_k(\alpha) = GL_n.p \simeq GL_n/GL(\alpha)$$

Thus, the  $GL_n$ -equivariant maps from  $\underline{rep}_n^{tr} A$  to  $M_n$  coincide with the  $Stab(p) = GL(\alpha)$ -equivariant maps from the fiber  $\pi^{-1}(p) = \underline{rep}_\alpha^{res} A$  to  $M_n$ .  $\square$

That is, we have a block matrix decomposition for  $A$ . Indeed, we have

$$A \simeq (\mathbb{C}[\underline{rep}_\alpha^{res} A] \otimes M_n(\mathbb{C}))^{GL(\alpha)}$$

and this isomorphism is clearly compatible with the block decomposition and thus we have for all  $i, j$  that

$$a_i A a_j \simeq (\mathbb{C}[\underline{rep}_\alpha^{res} A] \otimes M_{m_i \times m_j}(\mathbb{C}))^{GL(\alpha)}$$

where  $M_{m_i \times m_j}(\mathbb{C})$  is the space of rectangular  $m_i \times m_j$  matrices  $M$  with coefficients in  $\mathbb{C}$  on which  $GL(\alpha)$  acts via

$$g.M = g_i M g_j^{-1} \quad \text{where } g = (g_1, \dots, g_k) \in GL(\alpha).$$

If we specialize this result to the case of quiver orders we have

$$\underline{rep}_\alpha^{res} \mathbb{T}_\alpha Q \simeq \underline{rep}_\alpha Q$$

as  $GL(\alpha)$ -varieties and we deduce

**THEOREM 3.33.** *With notations as before,*

- (1)  $\mathbb{T}_\alpha Q$  is the algebra of  $GL(\alpha)$ -equivariant maps from  $\underline{rep}_\alpha Q$  to  $M_n$ , that is,

$$\mathbb{T}_\alpha Q = M_n(\mathbb{C}[\underline{rep}_\alpha Q])^{GL(\alpha)}$$

- (2) The quiver necklace algebra

$$\mathbb{N}_\alpha Q = \mathbb{C}[\underline{rep}_\alpha Q]^{GL(\alpha)}$$

is generated by traces along oriented cycles in the quiver  $Q$  of length bounded by  $n^2 + 1$ .

A concrete realization of these algebras is as follows. To an arrow  $\textcircled{j} \xleftarrow{a} \textcircled{i}$  corresponds a  $d_j \times d_i$  matrix of variables from  $\mathbb{C}[\text{rep}_\alpha Q]$

$$\boxed{M_a} = \begin{bmatrix} x_{11}(a) & \dots & \dots & x_{1d_i}(a) \\ \vdots & & & \vdots \\ x_{d_j1}(a) & \dots & \dots & x_{d_jd_i}(a) \end{bmatrix}$$

where  $x_{ij}(a)$  are the coordinate functions of the entries of  $V_a$  of a representation  $V \in \text{rep}_\alpha Q$ . Let  $p = a_1 a_2 \dots a_r$  be an oriented cycle in  $Q$ , then we can compute the following matrix

$$M_p = M_{a_r} \dots M_{a_2} M_{a_1}$$

over  $\mathbb{C}[\text{rep}_\alpha Q]$ . As we have that  $s(a_r) = t(a_1) = v_i$ , this is a square  $d_i \times d_i$  matrix with coefficients in  $\mathbb{C}[\text{rep}_\alpha Q]$  and we can take its ordinary trace

$$\text{Tr}(M_p) \in \mathbb{C}[\text{rep}_\alpha Q].$$

Then,  $\mathbb{N}_\alpha Q$  is the  $\mathbb{C}$ -subalgebra of  $\mathbb{C}[\text{rep}_\alpha Q]$  generated by these elements. Consider the block structure of  $M_n(\mathbb{C}[\text{rep}_\alpha Q])$  with respect to the idempotents  $e_i$

$$\begin{bmatrix} M_{d_1}(S) & \dots & \dots & M_{d_1 \times d_k}(S) \\ \vdots & & & \vdots \\ \vdots & M_{d_j \times d_i}(S) & & \vdots \\ M_{d_k \times d_i}(S) & \dots & \dots & M_{d_k}(S) \end{bmatrix}$$

where  $S = \mathbb{C}[\text{rep}_\alpha Q]$ . Then, we can also view the matrix  $M_a$  for an arrow  $\textcircled{j} \xleftarrow{a} \textcircled{i}$  as a block matrix in  $M_n(\mathbb{C}[\text{rep}_\alpha Q])$

$$\begin{bmatrix} 0 & \dots & \dots & 0 \\ \vdots & & & \vdots \\ \vdots & \boxed{M_a} & & \vdots \\ 0 & \dots & \dots & 0 \end{bmatrix}$$


Then,  $\mathbb{T}_\alpha Q$  is the  $C_k(\alpha)$ -subalgebra of  $M_n(\mathbb{C}[\text{rep}_\alpha Q])$  generated by  $\mathbb{N}_\alpha Q$  and these block matrices for all arrows  $a \in Q_a$ .  $\mathbb{T}_\alpha Q$  itself has a block decomposition

$$\mathbb{T}_\alpha Q = \begin{bmatrix} P_{11} & \dots & \dots & P_{1k} \\ \vdots & & & \vdots \\ \vdots & P_{ij} & & \vdots \\ P_{k1} & \dots & \dots & P_{kk} \end{bmatrix}$$

where  $P_{ij}$  is the  $\mathbb{N}_\alpha Q$ -module spanned by all matrices  $M_p$  where  $p$  is a path from  $v_i$  to  $v_j$  of length bounded by  $n^2$ .




EXAMPLE 3.34. Consider the path algebra  $\mathbb{M}$  of the quiver



and take as dimension vector  $\alpha = (n, 1)$ . The total dimension is in this case  $\bar{n} = n + 1$  and we fix the embedding  $C_2 = \mathbb{C} \times \mathbb{C} \hookrightarrow \mathbb{M}$  given by the decomposition  $1 = e + f$ . Then, the above realization of  $T_\alpha \mathbb{M}$  consists in taking the following  $\bar{n} \times \bar{n}$  matrices

$$\begin{aligned}
 e_n &= \begin{bmatrix} 1 & & & 0 \\ & \ddots & & \vdots \\ & & 1 & 0 \\ 0 & \dots & 0 & 0 \end{bmatrix} & f_n &= \begin{bmatrix} 0 & \dots & 0 & 0 \\ \vdots & & \vdots & \vdots \\ 0 & \dots & 0 & 0 \\ 0 & \dots & 0 & 1 \end{bmatrix} & x_n &= \begin{bmatrix} x_{11} & \dots & x_{1n} & 0 \\ \vdots & & \vdots & \vdots \\ x_{n1} & \dots & x_{nn} & 0 \\ 0 & \dots & 0 & 0 \end{bmatrix} \\
 y_n &= \begin{bmatrix} y_{11} & \dots & y_{1n} & 0 \\ \vdots & & \vdots & \vdots \\ y_{n1} & \dots & y_{nn} & 0 \\ 0 & \dots & 0 & 0 \end{bmatrix} & u_n &= \begin{bmatrix} 0 & \dots & 0 & u_1 \\ \vdots & & \vdots & \vdots \\ 0 & \dots & 0 & u_n \\ 0 & \dots & 0 & 0 \end{bmatrix} & v_n &= \begin{bmatrix} 0 & \dots & 0 & 0 \\ \vdots & & \vdots & \vdots \\ 0 & \dots & 0 & 0 \\ v_1 & \dots & v_n & 0 \end{bmatrix}
 \end{aligned}$$

In order to determine the ring of  $GL(\alpha)$ -polynomial invariants of  $rep_\alpha \mathbb{M}$  we have to consider the traces along oriented cycles in the quiver. Any nontrivial such cycle must pass through the vertex  $e$  and then we can decompose the cycle into factors  $x, y$  and  $uv$  (observe that if we wanted to describe circuits based at the vertex  $f$  they are of the form  $c = vc'u$  with  $c'$  a circuit based at  $e$  and we can use the cyclic property of traces to bring it into the claimed form). That is, all relevant oriented cycles in the quiver can be represented by a necklace word  $w$



where each bead is one of the elements

$$\blacksquare = x \quad \square = y \quad \text{and} \quad \blacktriangledown = uv$$

In calculating the trace, we first have to replace each occurrence of  $x, y, u$  or  $v$  by the relevant  $\bar{n} \times \bar{n}$ -matrix above. This results in replacing each of the beads in the necklace by one of the following  $n \times n$  matrices

$$\blacksquare = \begin{bmatrix} x_{11} & \dots & x_{1n} \\ \vdots & & \vdots \\ x_{n1} & \dots & x_{nn} \end{bmatrix} \quad \square = \begin{bmatrix} y_{11} & \dots & y_{1n} \\ \vdots & & \vdots \\ y_{n1} & \dots & y_{nn} \end{bmatrix} \quad \blacktriangledown = \begin{bmatrix} u_1 v_1 & \dots & u_1 v_n \\ \vdots & & \vdots \\ u_n v_1 & \dots & u_n v_n \end{bmatrix}$$

and taking the trace of the  $n \times n$  matrix obtained after multiplying these bead-matrices cyclicly in the indicated orientation. This concludes the description of the invariant ring  $\mathbb{N}_\alpha \mathbb{Q}$ . The algebra  $T_\alpha \mathbb{M}$  of  $GL(\alpha)$ -equivariant maps from  $rep_\alpha \mathbb{M}$  to  $M_{\bar{n}}$  is then the subalgebra of  $M_{\bar{n}}(\mathbb{C}[rep_\alpha \mathbb{M}])$  generated as  $C_2(\alpha)$ -algebra (using the idempotent  $\bar{n} \times \bar{n}$  matrices corresponding to  $e$  and  $f$ ) by  $\mathbb{N}_\alpha \mathbb{M}$  and the  $\bar{n} \times \bar{n}$ -matrices corresponding to  $x, y, u$  and  $v$ .

### 3.4. Simple roots.

In this section we will use proposition 3.12 to construct quiver orders  $\mathbb{T}_\alpha Q$  which determine central simple algebras over the functionfield of the quotient variety  $iss_\alpha Q = rep_\alpha Q/GL(\alpha)$ . With  $PGL(\alpha)$  we denote the groupscheme corresponding to the algebraic group

$$PGL(\alpha) = GL(\alpha)/\mathbb{C}^*(\mathbb{1}_{d_1}, \dots, \mathbb{1}_{d_k})$$

If  $C$  is a commutative  $\mathbb{C}$ -algebra, then using the embedding  $PGL(\alpha) \hookrightarrow PGL_n$ , the pointed cohomology set

$$H_{et}^1(C, PGL(\alpha)) \hookrightarrow H_{et}^1(C, PGL_n)$$

classifies Azumaya algebras  $A$  over  $C$  with a distinguished embedding  $C_k \hookrightarrow A$  that are split by an étale cover such that on this cover the embedding of  $C_k$  in matrices is conjugate to the standard embedding  $C_k(\alpha)$ . Modifying the argument of proposition 3.12 we have

**PROPOSITION 3.35.** *If  $\alpha$  is the dimension vector of a simple representation of  $Q$ , then*

$$\mathbb{T}_\alpha Q \otimes_{\mathbb{N}_\alpha Q} \mathbb{C}(iss_\alpha Q)$$

*is a central simple algebra over the function field of the quotient variety  $iss_\alpha Q$ .*

Remains to classify the *simple roots*  $\alpha$ , that is, the dimension vectors of simple representations of the quiver  $Q$ . Consider the vertex set  $Q_v = \{v_1, \dots, v_k\}$ . To a subset  $S \hookrightarrow Q_v$  we associate the full subquiver  $Q_S$  of  $Q$ , that is,  $Q_S$  has as set of vertices the subset  $S$  and as set of arrows all arrows  $\textcircled{i} \xleftarrow{a} \textcircled{j}$  in  $Q_a$  such that  $v_i$  and  $v_j$  belong to  $S$ . A full subquiver  $Q_S$  is said to be *strongly connected* if and only if for all  $v_i, v_j \in V$  there is an oriented cycle in  $Q_S$  passing through  $v_i$  and  $v_j$ . We can partition

$$Q_v = S_1 \sqcup \dots \sqcup S_s$$

such that the  $Q_{S_i}$  are maximal strongly connected components of  $Q$ . Clearly, the direction of arrows in  $Q$  between vertices in  $S_i$  and  $S_j$  is the same by the maximality assumption and can be used to define an orientation between  $S_i$  and  $S_j$ . The *strongly connected component quiver*  $SC(Q)$  is then the quiver on  $s$  vertices  $\{w_1, \dots, w_s\}$  with  $w_i$  corresponding to  $S_i$  and there is one arrow from  $w_i$  to  $w_j$  if and only if there is an arrow in  $Q$  from a vertex in  $S_i$  to a vertex in  $S_j$ . Observe that when the underlying graph of  $Q$  is connected, then so is the underlying graph of  $SC(Q)$  and  $SC(Q)$  is a quiver without oriented cycles.

Vertices with specific in- and out-going arrows are given names as in figure 3

If  $\alpha = (d_1, \dots, d_k)$  is a dimension vector, we define the *support* of  $\alpha$  to be  $supp(\alpha) = \{v_i \in Q_v \mid d_i \neq 0\}$ .

**LEMMA 3.36.** *If  $\alpha$  is the dimension vector of a simple representation of  $Q$ , then  $Q_{supp(\alpha)}$  is a strongly connected subquiver.*

**PROOF.** If not, we consider the strongly connected component quiver  $SC(Q_{supp(\alpha)})$  and by assumption there must be a sink in it corresponding to a proper subset  $S \subsetneq Q_v$ . If  $V \in rep_\alpha Q$  we can then construct a representation  $W$  by

- $W_i = V_i$  for  $v_i \in S$  and  $W_i = 0$  if  $v_i \notin S$ ,
- $W_a = V_a$  for an arrow  $a$  in  $Q_S$  and  $W_a = 0$  otherwise.




FIGURE 3. Vertex terminology

One verifies that  $W$  is a proper subrepresentation of  $V$ , so  $V$  cannot be simple, a contradiction.  $\square$

The second necessary condition involves the Euler form of  $Q$ . With  $\epsilon_i$  be denote the dimension vector of the simple representation having a one-dimensional space at vertex  $v_i$  and zero elsewhere and all arrows zero matrices.

LEMMA 3.37. *If  $\alpha$  is the dimension vector of a simple representation of  $Q$ , then*

$$\begin{cases} \chi_Q(\alpha, \epsilon_i) \leq 0 \\ \chi_Q(\epsilon_i, \alpha) \leq 0 \end{cases}$$

for all  $v_i \in \text{supp}(\alpha)$ .

PROOF. Let  $V$  be a simple representation of  $Q$  with dimension vector  $\alpha = (d_1, \dots, d_k)$ . One verifies that

$$\chi_Q(\epsilon_i, \alpha) = d_i - \sum_{\substack{j \leftarrow a \\ i}} d_j$$

Assume that  $\chi_Q(\epsilon_i, \alpha) > 0$ , then the natural linear map

$$\bigoplus_{\substack{j \leftarrow a \\ i}} V_a : V_i \longrightarrow \bigoplus_{\substack{j \leftarrow a \\ i}} V_j$$

has a nontrivial kernel, say  $K$ . But then we consider the representation  $W$  of  $Q$  determined by

- $W_i = K$  and  $W_j = 0$  for all  $j \neq i$ ,
- $W_a = 0$  for all  $a \in Q_a$ .

It is clear that  $W$  is a proper subrepresentation of  $V$ , a contradiction.

Similarly, assume that  $\chi_Q(\alpha, \epsilon_i) = d_i - \sum d_j > 0$ , then the linear map

$$\textcircled{i} \xleftarrow{a} \textcircled{j}$$


$$\bigoplus_{\textcircled{i} \xleftarrow{a} \textcircled{j}} V_a : \bigoplus_{\textcircled{i} \xleftarrow{a} \textcircled{j}} V_j \longrightarrow V_i$$

has an image  $I$  which is a proper subspace of  $V_i$ . The representation  $W$  of  $Q$  determined by

- $W_i = I$  and  $W_j = V_j$  for  $j \neq i$ ,
- $W_a = V_a$  for all  $a \in Q_a$ .

is a proper subrepresentation of  $V$ , a contradiction finishing the proof. □

EXAMPLE 3.38. The necessary conditions of the foregoing two lemmas are not sufficient. Consider the extended Dynkin quiver of type  $\tilde{A}_k$  with cyclic orientation.



and dimension vector  $\alpha = (a, \dots, a)$ . For a simple representation all arrow matrices must be invertible but then, under the action of  $GL(\alpha)$ , they can be diagonalized. Hence, the only simple representations (which are not the trivial simples concentrated in a vertex) have dimension vector  $(1, \dots, 1)$ .

Nevertheless, we will show that these are the only exceptions. A vertex  $v_i$  is said to be *large* with respect to a dimension vector  $\alpha = (d_1, \dots, d_k)$  whenever  $d_i$  is maximal among the  $d_j$ . The vertex  $v_i$  is said to be *good* if  $v_i$  is large and has no direct successor which is a large prism nor a direct predecessor which is a large focus.


LEMMA 3.39. *Let  $Q$  be a strongly connected quiver, not of type  $\tilde{A}_k$ , then one of the following hold*

- (1)  $Q$  has a good vertex, or,
- (2)  $Q$  has a large prism having no direct large prism successors, or
- (3)  $Q$  has a large focus having no direct large focus predecessors.

PROOF. If neither of the cases hold, we would have an oriented cycle in  $Q$  consisting of prisms (or consisting of focusses). Assume  $(v_{i_1}, \dots, v_{i_l})$  is a cycle of prisms, then the unique incoming arrow of  $v_{i_j}$  belongs to the cycle. As  $Q \neq \tilde{A}_k$  there is at least one extra vertex  $v_a$  not belonging to the cycle. But then, there can be no oriented path from  $v_a$  to any of the  $v_{i_j}$ , contradicting the assumption that  $Q$  is strongly connected. □

If we are in one of the two last cases, let  $a$  be the maximum among the components of the dimension vector  $\alpha$  and assume that  $\alpha$  satisfies  $\chi_Q(\alpha, \epsilon_i) \leq 0$  and

$\chi_Q(\epsilon_i, \alpha) \leq 0$  for all  $1 \leq i \leq k$ , then we have the following subquiver in  $Q$




We can reduce to a quiver situation with strictly less vertices.

LEMMA 3.40. *Assume  $Q$  is strongly connected and we have a vertex  $v_i$  which is a prism with unique predecessor the vertex  $v_j$ , which is a focus. Consider the dimension vector  $\alpha = (d_1, \dots, d_k)$  with  $d_i = d_j = a \neq 0$ . Then,  $\alpha$  is the dimension of a simple representation of  $Q$  if and only if*

$$\alpha' = (d_1, \dots, d_{i-1}, d_{i+1}, \dots, d_k) \in \mathbb{N}^{k-1}$$


*is the dimension vector of a simple representation of the quiver  $Q'$  on  $k-1$  vertices, obtained from  $Q$  by identifying the vertices  $v_i$  and  $v_j$ , that is, the above subquiver in  $Q$  is simplified to the one below in  $Q'$*



PROOF. If  $b$  is the unique arrow from  $v_j$  to  $v_i$  and if  $V \in \text{rep}_\alpha Q$  is a simple representation then  $V_b$  is an isomorphism, so we can identify  $V_i$  with  $V_j$  and obtain a simple representation of  $Q'$ . Conversely, if  $V' \in \text{rep}_{\alpha'} Q'$  is a simple representation, define  $V \in \text{rep}_\alpha Q$  by  $V_i = V'_j$  and  $V_z = V'_z$  for  $z \neq i$ ,  $V_{b'} = V'_{b'}$  for all arrows  $b' \neq b$  and  $V_b = \mathbb{1}_a$ . Clearly,  $V$  is a simple representation of  $Q$ .  $\square$

THEOREM 3.41.  $\alpha = (d_1, \dots, d_k)$  is the dimension vector of a simple representation of  $Q$  if and only if one of the following two cases holds

- (1)  $\text{supp}(\alpha) = \tilde{A}_k$ , the extended Dynkin quiver on  $k$  vertices with cyclic orientation and  $d_i = 1$  for all  $1 \leq i \leq k$



(2)  $\text{supp}(\alpha) \neq \tilde{A}_k$ . Then,  $\text{supp}(\alpha)$  is strongly connected and for all  $1 \leq i \leq k$  we have

$$\begin{cases} \chi_Q(\alpha, \epsilon_i) \leq 0 \\ \chi_Q(\epsilon_i, \alpha) \leq 0 \end{cases}$$

PROOF. We will use induction, both on the number of vertices  $k$  in  $\text{supp}(\alpha)$  and on the total dimension  $n = \sum_i d_i$  of the representation. If  $\text{supp}(\alpha)$  does not possess a good vertex, then the above lemma finishes the proof by induction on  $k$ . Observe that the Euler-form conditions are preserved in passing from  $Q$  to  $Q'$  as  $d_i = d_j$ .

Hence, assume  $v_i$  is a good vertex in  $\text{supp}(\alpha)$ . If  $d_i = 1$  then all  $d_j = 1$  for  $v_j \in \text{supp}(\alpha)$  and we can construct a simple representation by taking  $V_b = 1$  for all arrows  $b$  in  $\text{supp}(\alpha)$ . Simplicity follows from the fact that  $\text{supp}(\alpha)$  is strongly connected.

If  $d_i > 1$ , consider the dimension vector  $\alpha' = (d_1, \dots, d_{i-1}, d_i - 1, d_{i+1}, \dots, d_k)$ . Clearly,  $\text{supp}(\alpha') = \text{supp}(\alpha)$  is strongly connected and we claim that the Euler-form conditions still hold for  $\alpha'$ . The only vertices  $v_l$  where things might go wrong are direct predecessors or direct successors of  $v_i$ . Assume for one of them  $\chi_Q(\epsilon_l, \alpha) > 0$  holds, then

$$d_l = d'_l > \sum_{\textcircled{m} \xleftarrow{a} \textcircled{i}} d'_m \geq d'_i = d_i - 1$$

But then,  $d_l = d_i$  whence  $v_l$  is a large vertex of  $\alpha$  and has to be also a focus with end vertex  $v_i$  (if not,  $d_l > d_i$ ), contradicting goodness of  $v_i$ .

Hence, by induction on  $n$  we may assume that there is a simple representation  $W \in \text{rep}_{\alpha'} Q$ . Consider the space  $\text{rep}_W$  of representations  $V \in \text{rep}_{\alpha} Q$  such that  $V \mid \alpha' = W$ . That is, for every arrow

$$\begin{array}{l} \textcircled{j} \xleftarrow{a} \textcircled{i} \quad V_a = \begin{array}{|c|} \hline W_a \\ \hline v_1 \quad \dots \quad v_{d_j} \\ \hline \end{array} \\ \\ \textcircled{i} \xleftarrow{a} \textcircled{j} \quad V_a = \begin{array}{|c|c|} \hline & v_1 \\ \hline W_a & \vdots \\ \hline & v_{d_j} \\ \hline \end{array} \end{array}$$

Hence,  $\text{rep}_W$  is an affine space consisting of all representations degenerating to  $W \oplus S_i$  where  $S_i$  is the simple one-dimensional representation concentrated in  $v_i$ . As  $\chi_Q(\alpha', \epsilon_i) < 0$  and  $\chi_Q(\epsilon_i, \alpha') < 0$  we have that  $\text{Ext}^1(W, S_i) \neq 0 \neq \text{Ext}^1(S_i, W)$  so there is an open subset of representations which are not isomorphic to  $W \oplus S_i$ .

As there are simple representations of  $Q$  having a one-dimensional component at each vertex in  $\text{supp}(\alpha)$  and as the subset of simple representations in  $\text{rep}_{\alpha'} Q$  is open, we can choose  $W$  such that  $\text{rep}_W$  contains representations  $V$  such that a trace of an oriented cycle differs from that of  $W \oplus S_i$ . Hence, by the description of the invariant ring  $\mathbb{C}[\text{rep}_{\alpha} Q]^{GL(\alpha)}$  as being generated by traces along oriented cycles and by the identification of points in the quotient variety as isomorphism classes of semi-simple representations, it follows that the Jordan-Hölder factors of  $V$  are different from  $W$  and  $S_i$ . In view of the definition of  $\text{rep}_W$ , this can only happen if  $V$  is a simple representation, finishing the proof of the theorem.  $\square$

Still, the central simple algebras constructed from quivers are very special examples as we will see in section 4.8. In the next sections we will show that the Brauer groups of function fields are huge objects but first we need to recall some facts about spectral sequences.

### 3.5. Spectral sequences

Let  $\mathcal{A}, \mathcal{B}$  and  $\mathcal{C}$  be Abelian categories such that  $\mathcal{A}$  and  $\mathcal{B}$  have enough injectives and consider left exact functors

$$\mathcal{A} \xrightarrow{f} \mathcal{B} \xrightarrow{g} \mathcal{C}$$

Let the functors be such that  $f$  maps injectives of  $\mathcal{A}$  to  $g$ -acyclic objects in  $\mathcal{B}$ , that is  $R^i g(f I) = 0$  for all  $i > 0$ . Then, there are connections between the objects

$$R^p g(R^q f(A)) \text{ and } R^n gf(A)$$

for all objects  $A \in \mathcal{A}$ . These connections can be summarized by giving a spectral sequence

**THEOREM 3.42.** *Let  $\mathcal{A}, \mathcal{B}, \mathcal{C}$  be Abelian categories with  $\mathcal{A}, \mathcal{B}$  having enough injectives and left exact functors*

$$\mathcal{A} \xrightarrow{f} \mathcal{B} \xrightarrow{g} \mathcal{C}$$

*such that  $f$  takes injectives to  $g$ -acyclics.*

*Then, for any object  $A \in \mathcal{A}$  there is a spectral sequence*

$$E_2^{p,q} = R^p g(R^q f(A)) \implies R^n gf(A)$$

*In particular, there is an exact sequence*

$$0 \longrightarrow R^1 g(f(A)) \longrightarrow R^1 gf(A) \longrightarrow g(R^1 f(A)) \longrightarrow R^2 g(f(A)) \longrightarrow \dots$$

*Moreover, if  $f$  is an exact functor, then we have*

$$R^p gf(A) \simeq R^p g(f(A))$$

A spectral sequence  $E_2^{p,q} \implies E^n$  (or  $E_1^{p,q} \implies E^n$ ) consists of the following data


- (1) A family of objects  $E_r^{p,q}$  in an Abelian category for  $p, q, r \in \mathbb{Z}$  such that  $p, q \geq 0$  and  $r \geq 2$  (or  $r \geq 1$ ).
- (2) A family of morphisms in the Abelian category

$$d_r^{p,q} : E_r^{p,q} \longrightarrow E_r^{p+r, q-r+1}$$


satisfying the complex condition

$$d_r^{p+r, q-r+1} \circ d_r^{p,q} = 0$$

and where we assume that  $d_r^{p,q} = 0$  if any of the numbers  $p, q, p + r$  or  $q - r + 1$  is  $< 1$ . At level one we have the following



At level two we have the following



- (3) The objects  $E_{r+1}^{p,q}$  on level  $r + 1$  are derived from those on level  $r$  by taking the cohomology objects of the complexes, that is,

$$E_{r+1}^p = Ker d_r^{p,q} / Im d_r^{p-r,q+r-1}$$

At each place  $(p, q)$  this process converges as there is an integer  $r_0$  depending on  $(p, q)$  such that for all  $r \geq r_0$  we have  $d_r^{p,q} = 0 = d_r^{p-r,q+r-1}$ . We then define

$$E_\infty^{p,q} = E_{r_0}^{p,q} (= E_{r_0+1}^{p,q} = \dots)$$

Observe that there are injective maps  $E_\infty^{0,q} \hookrightarrow E_2^{0,q}$ .

- (4) A family of objects  $E^n$  for integers  $n \geq 0$  and for each we have a filtration


$$0 \subset E_n^n \subset E_{n-1}^n \subset \dots \subset E_1^n \subset E_0^n = E^n$$

such that the successive quotients are given by

$$E_p^n / E_{p+1}^n = E_\infty^{p,n-p}$$



That is, the terms  $E_\infty^{p,q}$  are the composition terms of the *limiting terms*  $E^{p+q}$ . Pictorially,



For small  $n$  one can make the relation between  $E^n$  and the terms  $E_2^{p,q}$  explicit. First note that

$$E_2^{0,0} = E_\infty^{0,0} = E^0$$

Also,  $E_1^1 = E_\infty^{1,0} = E_2^{1,0}$  and  $E^1/E_1^1 = E_\infty^{0,1} = Ker d_2^{0,1}$ . This gives an exact sequence

$$0 \longrightarrow E_2^{1,0} \longrightarrow E^1 \longrightarrow E_2^{0,1} \xrightarrow{d_2^{0,1}} E_2^{2,0}$$

Further,  $E^2 \supset E_1^2 \supset E_2^2$  where

$$E_2^2 = E_\infty^{2,0} = E_2^{2,0} / Im d_2^{0,1}$$

and  $E_1^2/E_2^2 = E_\infty^{1,1} = Ker d_2^{1,1}$  whence we can extend the above sequence to

$$\dots \longrightarrow E_2^{0,1} \xrightarrow{d_2^{0,1}} E_2^{2,0} \longrightarrow E_1^2 \longrightarrow E_2^{1,1} \xrightarrow{d_2^{1,1}} E_2^{3,0}$$

as  $E^2/E_1^2 = E_\infty^{0,2} \hookrightarrow E_2^{0,2}$  we have that  $E_1^2 = Ker (E^2 \longrightarrow E_2^{0,2})$ . If we specialize to the spectral sequence  $E_2^{p,q} = R^p g(R^q f(A)) \implies R^n gf(A)$  we obtain the exact sequence

$$\begin{aligned} 0 \longrightarrow R^1 g(f(A)) \longrightarrow R^1 gf(A) \longrightarrow g(R^1 f(A)) \longrightarrow R^2 g(f(A)) \longrightarrow \\ \longrightarrow E_1^2 \longrightarrow R^1 g(R^1 f(A)) \longrightarrow R^3 g(f(A)) \end{aligned}$$

where  $E_1^2 = Ker (R^2 gf(A) \longrightarrow g(R^2 f(A)))$ .

An important example of a spectral sequence is the *Leray spectral sequence*. Assume we have an algebra morphism  $A \xrightarrow{f} A'$  and a sheaf of groups  $\mathbb{G}$  on  $A'_{et}$ . We define the *direct image* of  $\mathbb{G}$  under  $f$  to be the sheaf of groups  $f_* \mathbb{G}$  on  $A_{et}$  defined by

$$f_* \mathbb{G}(B) = \mathbb{G}(B \otimes_A A')$$

for all  $B \in A_{et}$  (recall that  $B \otimes_A A' \in A'_{et}$  so the right hand side is well defined).

This gives us a left exact functor

$$f_* : \mathbf{S}^{ab}(A'_{et}) \longrightarrow \mathbf{S}^{ab}(A_{et})$$

and therefore we have right derived functors of it  $R^i f_*$ . If  $\mathbb{G}$  is an Abelian sheaf on  $A'_{et}$ , then  $R^i f_* \mathbb{G}$  is a sheaf on  $A_{et}$ . One verifies that its stalk in a prime ideal  $\mathfrak{p}$  is equal to

$$(R^i f_* \mathbb{G})_{\mathfrak{p}} = H_{et}^i(A_{\mathfrak{p}}^{sh} \otimes_A A', \mathbb{G})$$

where the right hand side is the direct limit of cohomology groups taken over all étale neighborhoods of  $\mathfrak{p}$ . We can relate cohomology of  $\mathbb{G}$  and  $f_*\mathbb{G}$  by the following

**THEOREM 3.43.** (*Leray spectral sequence*) *If  $\mathbb{G}$  is a sheaf of Abelian groups on  $A'_{\text{ét}}$  and  $A \xrightarrow{f} A'$  an algebra morphism, then there is a spectral sequence*

$$E_2^{p,q} = H_{\text{ét}}^p(A, R^q f_*\mathbb{G}) \implies H_{\text{ét}}^n(A, \mathbb{G})$$

*In particular, if  $R^j f_*\mathbb{G} = 0$  for all  $j > 0$ , then for all  $i \geq 0$  we have isomorphisms*

$$H_{\text{ét}}^i(A, f_*\mathbb{G}) \simeq H_{\text{ét}}^i(A', \mathbb{G})$$

### 3.6. Tsen and Tate fields

**DEFINITION 3.44.** A field  $K$  is said to be a  $Tsen^d$ -field if every homogeneous form of degree  $deg$  with coefficients in  $K$  and  $n > deg^d$  variables has a non-trivial zero in  $K$ .

For example, an algebraically closed field  $\mathbb{K}$  is a  $Tsen^0$ -field as any form in  $n$ -variables defines a hypersurface in  $\mathbb{P}_{\mathbb{K}}^{n-1}$ . In fact, algebraic geometry tells us a stronger story

**LEMMA 3.45.** *Let  $\mathbb{K}$  be algebraically closed. If  $f_1, \dots, f_r$  are forms in  $n$  variables over  $\mathbb{K}$  and  $n > r$ , then these forms have a common non-trivial zero in  $\mathbb{K}$ .*

**PROOF.** Each  $f_i$  defines a hypersurface  $V(f_i) \hookrightarrow \mathbb{P}_{\mathbb{K}}^{n-1}$ . The intersection of  $r$  hypersurfaces has dimension  $\geq n - 1 - r$  from which the claim follows.  $\square$

We want to extend this fact to higher Tsen-fields. The proof of the following result is technical unenlightening inequality manipulation, see for example [74].

**PROPOSITION 3.46.** *Let  $K$  be a  $Tsen^d$ -field and  $f_1, \dots, f_r$  forms in  $n$  variables of degree  $deg$ . If  $n > r deg^d$ , then they have a non-trivial common zero in  $K$ .*

For our purposes the main interest in Tsen-fields comes from :

**THEOREM 3.47.** *Let  $K$  be of transcendence degree  $d$  over an algebraically closed field  $\mathbb{C}$ , then  $K$  is a  $Tsen^d$ -field.*

**PROOF.** First we claim that the purely transcendental field  $\mathbb{C}(t_1, \dots, t_d)$  is a  $Tsen^d$ -field. By induction we have to show that if  $L$  is  $Tsen^k$ , then  $L(t)$  is  $Tsen^{k+1}$ .

By homogeneity we may assume that  $f(x_1, \dots, x_n)$  is a form of degree  $deg$  with coefficients in  $L[t]$  and  $n > deg^{k+1}$ . For fixed  $s$  we introduce new variables  $y_{ij}^{(s)}$  with  $i \leq n$  and  $0 \leq j \leq s$  such that

$$x_i = y_{i0}^{(s)} + y_{i1}^{(s)}t + \dots + y_{is}^{(s)}t^s$$

If  $r$  is the maximal degree of the coefficients occurring in  $f$ , then we can write

$$f(x_i) = f_0(y_{ij}^{(s)}) + f_1(y_{ij}^{(s)})t + \dots + f_{deg.s+r}(y_{ij}^{(s)})t^{deg.s+r}$$

where each  $f_j$  is a form of degree  $deg$  in  $n(s+1)$ -variables. By the proposition above, these forms have a common zero in  $L$  provided

$$n(s+1) > deg^k(ds+r+1) \iff (n - deg^{k+1})s > deg^k(r+1) - n$$

which can be satisfied by taking  $s$  large enough. the common non-trivial zero in  $L$  of the  $f_j$ , gives a non-trivial zero of  $f$  in  $L[t]$ .

By assumption,  $K$  is an algebraic extension of  $\mathbb{C}(t_1, \dots, t_d)$  which by the above argument is  $Tsen^d$ . As the coefficients of any form over  $K$  lie in a finite extension  $E$  of  $\mathbb{C}(t_1, \dots, t_d)$  it suffices to prove that  $E$  is  $Tsen^d$ .

Let  $f(x_1, \dots, x_n)$  be a form of degree  $deg$  in  $E$  with  $n > deg^d$ . Introduce new variables  $y_{ij}$  with

$$x_i = y_{i1}e_1 + \dots + y_{ik}e_k$$

where  $e_i$  is a basis of  $E$  over  $\mathbb{C}(t_1, \dots, t_d)$ . Then,

$$f(x_i) = f_1(y_{ij})e_1 + \dots + f_k(y_{ij})e_k$$

where the  $f_i$  are forms of degree  $deg$  in  $k \cdot n$  variables over  $\mathbb{C}(t_1, \dots, t_d)$ . Because  $\mathbb{C}(t_1, \dots, t_d)$  is  $Tsen^d$ , these forms have a common zero as  $k \cdot n > k \cdot deg^d$ . Finding a non-trivial zero of  $f$  in  $E$  is equivalent to finding a common non-trivial zero to the  $f_1, \dots, f_k$  in  $\mathbb{C}(t_1, \dots, t_d)$ , done.  $\square$

A direct application of this result is *Tsen's theorem* :

**THEOREM 3.48.** *Let  $K$  be the functionfield of a curve  $C$  defined over an algebraically closed field. Then, the only central simple  $K$ -algebras are  $M_n(K)$ . That is,  $Br(K) = 1$ .*

**PROOF.** Assume there exists a central division algebra  $\Delta$  of dimension  $n^2$  over  $K$ . There is a finite Galois extension  $L/K$  such that  $\Delta \otimes L = M_n(L)$ . If  $x_1, \dots, x_{n^2}$  is a  $K$ -basis for  $\Delta$ , then the reduced norm of any  $x \in \Delta$ ,

$$N(x) = \det(x \otimes 1)$$

is a form in  $n^2$  variables of degree  $n$ . Moreover, as  $x \otimes 1$  is invariant under the action of  $Gal(L/K)$  the coefficients of this form actually lie in  $K$ .

By the main result,  $K$  is a  $Tsen^1$ -field and  $N(x)$  has a non-trivial zero whenever  $n^2 > n$ . As the reduced norm is multiplicative, this contradicts  $N(x)N(x^{-1}) = 1$ . Hence,  $n = 1$  and the only central division algebra is  $K$  itself.  $\square$

If  $K$  is the functionfield of a surface, we also have an immediate application :

**PROPOSITION 3.49.** *Let  $K$  be the functionfield of a surface defined over an algebraically closed field. If  $\Delta$  is a central simple  $K$ -algebra of dimension  $n^2$ , then the reduced norm map*

$$N : \Delta \longrightarrow K$$

*is surjective.*

**PROOF.** Let  $e_1, \dots, e_{n^2}$  be a  $K$ -basis of  $\Delta$  and  $k \in K$ , then

$$N\left(\sum x_i e_i\right) - kx_{n^2+1}$$

is a form of degree  $n$  in  $n^2 + 1$  variables. Since  $K$  is a  $Tsen^2$  field, it has a non-trivial solution  $(x_i^0)$ , but then,  $\delta = (\sum x_i^0 e_i)x_{n^2+1}^{-1}$  has reduced norm equal to  $k$ .  $\square$

From the cohomological description of the Brauer group it is clear that we need to have some control on the absolute Galois group  $G_K = Gal(\mathbb{K}/K)$ . We will see that finite transcendence degree forces some cohomology groups to vanish.

**DEFINITION 3.50.** The *cohomological dimension* of a group  $G$ ,  $cd(G) \leq d$  if and only if  $H^r(G, A) = 0$  for all  $r > d$  and all torsion modules  $A \in G\text{-mod}$ .

DEFINITION 3.51. A field  $K$  is said to be a  $Tate^d$ -field if the absolute Galois group  $G_K = Gal(\mathbb{K}/K)$  satisfies  $cd(G) \leq d$ .

First, we will reduce the condition  $cd(G) \leq d$  to a more manageable one. To start, one can show that a profinite group  $G$  (that is, a projective limit of finite groups, see [74] for more details) has  $cd(G) \leq d$  if and only if

$$H^{d+1}(G, A) = 0 \text{ for all torsion } G\text{-modules } A$$

Further, as all Galois cohomology groups of profinite groups are torsion, we can decompose the cohomology in its  $p$ -primary parts and relate their vanishing to the cohomological dimension of the  $p$ -Sylow subgroups  $G_p$  of  $G$ . This problem can then be verified by computing cohomology of finite simple  $G_p$ -modules of  $p$ -power order, but for a profinite  $p$ -group there is just one such module namely  $\mathbb{Z}/p\mathbb{Z}$  with the trivial action.

Combining these facts we have the following manageable criterium on cohomological dimension.

PROPOSITION 3.52.  $cd(G) \leq d$  if  $H^{d+1}(G, \mathbb{Z}/p\mathbb{Z}) = 0$  for the simple  $G$ -modules with trivial action  $\mathbb{Z}/p\mathbb{Z}$ .

We will need the following spectral sequence in Galois cohomology

PROPOSITION 3.53. (Hochschild-Serre spectral sequence) If  $N$  is a closed normal subgroup of a profinite group  $G$ , then

$$E_2^{p,q} = H^p(G/N, H^q(N, A)) \implies H^n(G, A)$$

holds for every continuous  $G$ -module  $A$ .

Now, we are in a position to state and prove *Tate's theorem*

THEOREM 3.54. Let  $K$  be of transcendence degree  $d$  over an algebraically closed field, then  $K$  is a  $Tate^d$ -field.

PROOF. Let  $\mathbb{C}$  denote the algebraically closed basefield, then  $K$  is algebraic over  $\mathbb{C}(t_1, \dots, t_d)$  and therefore

$$G_K \hookrightarrow G_{\mathbb{C}(t_1, \dots, t_d)}$$

Thus,  $K$  is  $Tate^d$  if  $\mathbb{C}(t_1, \dots, t_d)$  is  $Tate^d$ . By induction it suffices to prove

$$\text{If } cd(G_L) \leq k \text{ then } cd(G_{L(t)}) \leq k + 1$$

Let  $\mathbb{L}$  be the algebraic closure of  $L$  and  $\mathbb{M}$  the algebraic closure of  $L(t)$ . As  $L(t)$  and  $\mathbb{L}$  are linearly disjoint over  $L$  we have the following diagram of extensions and Galois groups

$$\begin{array}{ccccc} \mathbb{L} & \hookrightarrow & \mathbb{L}(t) & \xrightarrow{G_{\mathbb{L}(t)}} & \mathbb{M} \\ \uparrow G_L & & \uparrow G_L & \nearrow G_{\mathbb{L}(t)} & \\ L & \hookrightarrow & L(t) & & \end{array}$$

where  $G_{L(t)}/G_{\mathbb{L}(t)} \simeq G_L$ .

We claim that  $cd(G_{\mathbb{L}(t)}) \leq 1$ . Consider the exact sequence of  $G_{L(t)}$ -modules

$$0 \longrightarrow \mu_p \longrightarrow \mathbb{M}^* \xrightarrow{(-)^p} \mathbb{M}^* \longrightarrow 0$$

where  $\mu_p$  is the subgroup (of  $\mathbb{C}^*$ ) of  $p$ -roots of unity. As  $G_{L(t)}$  acts trivially on  $\mu_p$  it is after a choice of primitive  $p$ -th root of one isomorphic to  $\mathbb{Z}/p\mathbb{Z}$ . Taking cohomology with respect to the subgroup  $G_{\mathbb{L}(t)}$  we obtain

$$0 = H^1(G_{\mathbb{L}(t)}, \mathbb{M}^*) \longrightarrow H^2(G_{\mathbb{L}(t)}, \mathbb{Z}/p\mathbb{Z}) \longrightarrow H^2(G_{\mathbb{L}(t)}, \mathbb{M}^*) = Br(\mathbb{L}(t))$$

But the last term vanishes by Tsen's theorem as  $\mathbb{L}(t)$  is the functionfield of a curve defined over the algebraically closed field  $\mathbb{L}$ . Therefore,  $H^2(G_{\mathbb{L}(t)}, \mathbb{Z}/p\mathbb{Z}) = 0$  for all simple modules  $\mathbb{Z}/p\mathbb{Z}$ , whence  $cd(G_{\mathbb{L}(t)}) \leq 1$ .


By the inductive assumption we have  $cd(G_L) \leq k$  and now we are going to use exactness of the sequence

$$0 \longrightarrow G_L \longrightarrow G_{L(t)} \longrightarrow G_{\mathbb{L}(t)} \longrightarrow 0$$

to prove that  $cd(G_{L(t)}) \leq k + 1$ . For, let  $A$  be a torsion  $G_{L(t)}$ -module and consider the Hochschild-Serre spectral sequence

$$E_2^{p,q} = H^p(G_L, H^q(G_{\mathbb{L}(t)}, A)) \implies H^n(G_{L(t)}, A)$$

By the restrictions on the cohomological dimensions of  $G_L$  and  $G_{\mathbb{L}(t)}$  the level two term has following shape



where the only non-zero groups are lying in the lower rectangular region. Therefore, all  $E_\infty^{p,q} = 0$  for  $p+q > k+1$ . Now, all the composition factors of  $H^{k+2}(G_{L(t)}, A)$  are lying on the indicated diagonal line and hence are zero. Thus,  $H^{k+2}(G_{L(t)}, A) = 0$  for all torsion  $G_{L(t)}$ -modules  $A$  and hence  $cd(G_{L(t)}) \leq k + 1$ .  $\square$

**THEOREM 3.55.** *If  $\mathbf{A}$  is a constant sheaf of an Abelian torsion group  $A$  on  $K_{\text{et}}$ , then*

$$H_{\text{et}}^i(K, \mathbf{A}) = 0$$

whenever  $i > \text{trdeg}_{\mathbb{C}}(K)$ .

### 3.7. Coniveau spectral sequence

Consider the setting

$$\begin{array}{ccc} A & \xrightarrow{i} & K \\ \downarrow \pi & & \\ k & & \end{array}$$

where  $A$  is a discrete valuation ring in  $K$  with residue field  $A/m = k$ . As always, we will assume that  $A$  is a  $\mathbb{C}$ -algebra. By now we have a grip on the Galois cohomology groups

$$H_{et}^i(K, \mu_n^{\otimes l}) \text{ and } H_{et}^i(k, \mu_n^{\otimes l})$$

and we will use this information to compute the étale cohomology groups

$$H_{et}^i(A, \mu_n^{\otimes l})$$

Here,  $\mu_n^{\otimes l} = \underbrace{\mu_n \otimes \dots \otimes \mu_n}_l$  where the tensorproduct is as sheafs of invertible  $\mathbb{Z}_n = \mathbb{Z}/n\mathbb{Z}$ -modules.

We will consider the Leray spectral sequence for  $i$  and hence have to compute the derived sheaves of the direct image

- LEMMA 3.56. (1)  $R^0 i_* \mu_n^{\otimes l} \simeq \mu_n^{\otimes l}$  on  $\mathbf{A}_{et}$ .  
 (2)  $R^1 i_* \mu_n^{\otimes l} \simeq \mu_n^{\otimes l-1}$  concentrated in  $m$ .  
 (3)  $R^j i_* \mu_n^{\otimes l} \simeq 0$  whenever  $j \geq 2$ .

PROOF. The strict Henselizations of  $A$  at the two primes  $\{0, m\}$  are resp.

$$A_0^{sh} \simeq \mathbb{K} \text{ and } A_m^{sh} \simeq \mathbf{k}\{t\}$$

where  $\mathbb{K}$  (resp.  $\mathbf{k}$ ) is the algebraic closure of  $K$  (resp.  $k$ ). Therefore,

$$(R^j i_* \mu_n^{\otimes l})_0 = H_{et}^j(\mathbb{K}, \mu_n^{\otimes l})$$

which is zero for  $i \geq 1$  and  $\mu_n^{\otimes l}$  for  $j = 0$ . Further,  $A_m^{sh} \otimes_A K$  is the field of fractions of  $\mathbf{k}\{t\}$  and hence is of transcendence degree one over the algebraically closed field  $\mathbf{k}$ , whence

$$(R^j i_* \mu_n^{\otimes l})_m = H_{et}^j(L, \mu_n^{\otimes l})$$

which is zero for  $j \geq 2$  because  $L$  is Tate<sup>1</sup>.

For the field-tower  $K \subset L \subset \mathbb{K}$  we have that  $G_L = \hat{\mathbb{Z}} = \varprojlim \mu_m$  because the only Galois extensions of  $L$  are the Kummer extensions obtained by adjoining  $\sqrt[m]{t}$ . But then,

$$H_{et}^1(L, \mu_n^{\otimes l}) = H^1(\hat{\mathbb{Z}}, \mu_n^{\otimes l}(\mathbb{K})) = Hom(\hat{\mathbb{Z}}, \mu_n^{\otimes l}(\mathbb{K})) = \mu_n^{\otimes l-1}$$

from which the claims follow.  $\square$

THEOREM 3.57. *We have a long exact sequence*

$$\begin{aligned} 0 \longrightarrow H^1(A, \mu_n^{\otimes l}) \longrightarrow H^1(K, \mu_n^{\otimes l}) \longrightarrow H^0(k, \mu_n^{\otimes l-1}) \longrightarrow \\ H^2(A, \mu_n^{\otimes l}) \longrightarrow H^2(K, \mu_n^{\otimes l}) \longrightarrow H^1(k, \mu_n^{\otimes l-1}) \longrightarrow \dots \end{aligned}$$

PROOF. By the foregoing lemma, the second term of the Leray spectral sequence for  $i_* \mu_n^{\otimes l}$  looks like

0	0	0	...
$H^0(k, \mu_n^{\otimes l-1})$	$H^1(k, \mu_n^{\otimes l-1})$	$H^2(k, \mu_n^{\otimes l-1})$	...
$H^0(A, \mu_n^{\otimes l})$	$H^1(A, \mu_n^{\otimes l})$	$H^2(A, \mu_n^{\otimes l})$	...

with connecting morphisms

$$H_{et}^{i-1}(k, \mu_n^{\otimes l-1}) \xrightarrow{\alpha_i} H_{et}^{i+1}(A, \mu_n^{\otimes l})$$

The spectral sequences converges to its limiting term which looks like

0	0	0	...
$Ker \alpha_1$	$Ker \alpha_2$	$Ker \alpha_3$	...
$H^0(A, \mu_n^{\otimes l})$	$H^1(A, \mu_n^{\otimes l})$	$Coker \alpha_1$	...

and the Leray sequence gives the short exact sequences

$$\begin{aligned} 0 &\longrightarrow H_{et}^1(A, \mu_n^{\otimes l}) \longrightarrow H_{et}^1(K, \mu_n^{\otimes l}) \longrightarrow Ker \alpha_1 \longrightarrow 0 \\ 0 &\longrightarrow Coker \alpha_1 \longrightarrow H_{et}^2(K, \mu_n^{\otimes l}) \longrightarrow Ker \alpha_2 \longrightarrow 0 \\ 0 &\longrightarrow Coker \alpha_{i-1} \longrightarrow H_{et}^i(K, \mu_n^{\otimes l}) \longrightarrow Ker \alpha_i \longrightarrow 0 \end{aligned}$$

and gluing these sequences gives us the required result. □

In particular, if  $A$  is a discrete valuation ring of  $K$  with residue field  $k$  we have for each  $i$  a connecting morphism

$$H_{et}^i(K, \mu_n^{\otimes l}) \xrightarrow{\partial_{i,A}} H_{et}^{i-1}(k, \mu_n^{\otimes l-1})$$

Like any other topology, the étale topology can be defined locally on any scheme  $X$ . That is, we call a morphism of schemes

$$Y \xrightarrow{f} X$$

an étale extension (resp. cover) if locally  $f$  has the form

$$f^a | U_i : A_i = \Gamma(U_i, \mathcal{O}_X) \longrightarrow B_i = \Gamma(f^{-1}(U_i), \mathcal{O}_Y)$$

with  $A_i \longrightarrow B_i$  an étale extension (resp. cover) of algebras.




FIGURE 4. Coniveau spectral sequence

Again, we can construct the étale site of  $X$  locally and denote it with  $X_{et}$ . Presheaves and sheaves of groups on  $X_{et}$  are defined similarly and the right derived functors of the left exact global sections functor

$$\Gamma : \mathbf{S}^{ab}(X_{et}) \longrightarrow \text{abelian}$$

will be called the cohomology functors and we denote

$$R^i \Gamma(\mathbb{G}) = H_{et}^i(X, \mathbb{G})$$

From now on we restrict to the case when  $X$  is a smooth, irreducible projective variety of dimension  $d$  over  $\mathbb{C}$ . In this case, we can initiate the computation of the cohomology groups  $H_{et}^i(X, \mu_n^{\otimes l})$  via Galois cohomology of functionfields of subvarieties using the coniveau spectral sequence

**THEOREM 3.58.** *Let  $X$  be a smooth irreducible variety over  $\mathbb{C}$ . Let  $X^{(p)}$  denote the set of irreducible subvarieties  $x$  of  $X$  of codimension  $p$  with functionfield  $\mathbb{C}(x)$ , then there exists a coniveau spectral sequence*

$$E_1^{p,q} = \bigoplus_{x \in X^{(p)}} H_{et}^{q-p}(\mathbb{C}(x), \mu_n^{\otimes l-p}) \implies H_{et}^{p+q}(X, \mu_n^{\otimes l})$$

In contrast to the spectral sequences used before, the existence of the coniveau spectral sequence by no means follows from general principles. In it, a lot of heavy machinery on étale cohomology of schemes is encoded. In particular,

- cohomology groups with support of a closed subscheme, see for example [61, p. 91-94], and
- cohomological purity and duality, see [61, p. 241-252]

a detailed exposition of which would take us too far afield. For more details we refer the reader to [12].

Using the results on cohomological dimension and vanishing of Galois cohomology of  $\mu_n^{\otimes k}$  when the index is larger than the transcendence degree, we see that the coniveau spectral sequence has shape as in figure 4 where the only non-zero terms are in the indicated region.



Let us understand the connecting morphisms at the first level, a typical instance of which is

$$\bigoplus_{x \in X^{(p)}} H^i(\mathbb{C}(x), \mu_n^{\oplus l-p}) \longrightarrow \bigoplus_{y \in X^{(p+1)}} H^{i-1}(\mathbb{C}(y), \mu_n^{\oplus l-p-1})$$

and consider one of the closed irreducible subvarieties  $x$  of  $X$  of codimension  $p$  and one of those  $y$  of codimension  $p + 1$ . Then, either  $y$  is not contained in  $x$  in which case the component map

$$H^i(\mathbb{C}(x), \mu_n^{\oplus l-p}) \longrightarrow H^{i-1}(\mathbb{C}(y), \mu_n^{\oplus l-p-1})$$

is the zero map. Or,  $y$  is contained in  $x$  and hence defines a codimension one subvariety of  $x$ . That is,  $y$  defines a discrete valuation on  $\mathbb{C}(x)$  with residue field  $\mathbb{C}(y)$ . In this case, the above component map is the connecting morphism defined above.

In particular, let  $K$  be the functionfield of  $X$ . Then we can define the unramified cohomology groups

$$F_n^{i,l}(K/\mathbb{C}) = Ker H^i(K, \mu_n^{\otimes l}) \xrightarrow{\oplus \partial_{i,A}} \oplus H^{i-1}(k_A, \mu_n^{\otimes l-1})$$

where the sum is taken over all discrete valuation rings  $A$  of  $K$  (or equivalently, the irreducible codimension one subvarieties of  $X$ ) with residue field  $k_A$ . By definition, this is a (stable) birational invariant of  $X$ . In particular, if  $X$  is (stably) rational over  $\mathbb{C}$ , then

$$F_n^{i,l}(K/\mathbb{C}) = 0 \text{ for all } i, l \geq 0$$

### 3.8. The Artin-Mumford exact sequence

In this section  $S$  will be a smooth irreducible projective surface.

DEFINITION 3.59.  $S$  is called *simply connected* if every étale cover  $Y \longrightarrow S$  is trivial, that is,  $Y$  is isomorphic to a finite disjoint union of copies of  $S$ .

The first term of the coniveau spectral sequence of  $S$  has following shape

$\vdots$	$\vdots$	$\vdots$	$\vdots$	
0	0	0	0	$\dots$
$H^2(\mathbb{C}(S), \mu_n)$	$\oplus_C H^1(\mathbb{C}(S), \mathbb{Z}_n)$	$\oplus_P \mu_n^{-1}$	0	$\dots$
$H^1(\mathbb{C}(S), \mu_n)$	$\oplus_C \mathbb{Z}_n$	0	0	$\dots$
$\mu_n$	0	0	0	$\dots$

where  $C$  runs over all irreducible curves on  $S$  and  $P$  over all points of  $S$ .

LEMMA 3.60. For any smooth  $S$  we have  $H^1(\mathbb{C}(S), \mu_n) \longrightarrow \oplus_C \mathbb{Z}_n$ . If  $S$  is simply connected,  $H_{\text{ét}}^1(S, \mu_n) = 0$ .

PROOF. Using the Kummer sequence  $1 \longrightarrow \mu_n \longrightarrow \mathbb{G}_m \xrightarrow{(-)} \mathbb{G}_m \longrightarrow 1$  and Hilbert 90 we obtain that

$$H_{\text{ét}}^1(\mathbb{C}(S), \mu_n) = \mathbb{C}(S)^* / \mathbb{C}(S)^{*n}$$

The first claim follows from the exact diagram describing divisors of rational functions

$$\begin{array}{ccccccc}
 & & \mu_n & \simeq & \mu_n & & 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & \mathbb{C}^* & \longrightarrow & \mathbb{C}(S)^* & \xrightarrow{\text{div}} & \bigoplus_C \mathbb{Z} \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & & & (-)^n & & n \cdot \\
 0 & \longrightarrow & \mathbb{C}^* & \longrightarrow & \mathbb{C}(S)^* & \xrightarrow{\text{div}} & \bigoplus_C \mathbb{Z} \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & \bigoplus_C \mathbb{Z}_n & \simeq & \bigoplus_C \mathbb{Z}_n
 \end{array}$$

By the coniveau spectral sequence we have that  $H_{\text{ét}}^1(S, \mu_n)$  is equal to the kernel of the morphism

$$H_{\text{ét}}^1(\mathbb{C}(S), \mu_n) \xrightarrow{\gamma} \bigoplus_C \mathbb{Z}_n$$

and in particular,  $H^1(S, \mu_n) \hookrightarrow H^1(\mathbb{C}(S), \mu_n)$ .

As for the second claim, an element in  $H^1(S, \mu_n)$  determines a cyclic extension  $L = \mathbb{C}(S) \sqrt[n]{f}$  with  $f \in \mathbb{C}(S)^* / \mathbb{C}(S)^{*n}$  such that in each fieldcomponent  $L_i$  of  $L$  there is an étale cover  $T_i \longrightarrow S$  with  $\mathbb{C}(T_i) = L_i$ . By assumption no non-trivial étale covers exist whence  $f = 1 \in \mathbb{C}(S)^* / \mathbb{C}(S)^{*n}$ .  $\square$

If we invoke another major tool in étale cohomology of schemes, *Poincaré duality*, see for example [61, VI, §11], we obtain the following information on the cohomology groups for  $S$ .

PROPOSITION 3.61. (*Poincaré duality for  $S$* ) *If  $S$  is simply connected, then*

- (1)  $H_{\text{ét}}^0(S, \mu_n) = \mu_n$
- (2)  $H_{\text{ét}}^1(S, \mu_n) = 0$
- (3)  $H_{\text{ét}}^3(S, \mu_n) = 0$
- (4)  $H_{\text{ét}}^4(S, \mu_n) = \mu_n^{-1}$

PROOF. The third claim follows from the second as both groups are dual to each other. The last claim follows from the fact that for any smooth irreducible projective variety  $X$  of dimension  $d$  one has that

$$H_{\text{ét}}^{2d}(X, \mu_n) \simeq \mu_n^{\otimes 1-d}$$

$\square$

We are now in a position to state and prove the important

**THEOREM 3.62.** (*Artin-Mumford exact sequence*) *If  $S$  is a simply connected smooth projective surface, then the sequence*

$$0 \longrightarrow Br_n(S) \longrightarrow Br_n(\mathbb{C}(S)) \longrightarrow \bigoplus_C \mathbb{C}(C)^*/\mathbb{C}(C)^{*n} \longrightarrow \\ \longrightarrow \bigoplus_P \mu_n^{-1} \longrightarrow \mu_n^{-1} \longrightarrow 0$$

is exact.

**PROOF.** The top complex in the first term of the coniveau spectral sequence for  $S$  was

$$H^2(\mathbb{C}(S), \mu_n) \xrightarrow{\alpha} \bigoplus_C H^1(\mathbb{C}(C), \mathbb{Z}_n) \xrightarrow{\beta} \bigoplus_P \mu_n$$

The second term of the spectral sequence (which is also the limiting term) has the following form

$\vdots$	$\vdots$	$\vdots$	$\vdots$	
0	0	0	0	...
$Ker \alpha$	$Ker \beta / Im \alpha$	$Coker \beta$	0	...
$Ker \gamma$	$Coker \gamma$	0	0	...
$\mu_n$	0	0	0	...

By the foregoing lemma we know that  $Coker \gamma = 0$ . By Poincare duality we know that  $Ker \beta = Im \alpha$  and  $Coker \beta = \mu_n^{-1}$ . Hence, the top complex was exact in its middle term and can be extended to an exact sequence

$$0 \longrightarrow H^2(S, \mu_n) \longrightarrow H^2(\mathbb{C}(S), \mu_n) \longrightarrow \bigoplus_C H^1(\mathbb{C}(C), \mathbb{Z}_n) \longrightarrow \\ \bigoplus_P \mu_n^{-1} \longrightarrow \mu_n^{-1} \longrightarrow 0$$

As  $\mathbb{Z}_n \simeq \mu_n$  the third term is equal to  $\bigoplus_C \mathbb{C}(C)^*/\mathbb{C}(C)^{*n}$  by the argument given before and the second term we remember to be  $Br_n(\mathbb{C}(S))$ . The identification of  $Br_n(S)$  with  $H^2(S, \mu_n)$  will be explained below.  $\square$

Some immediate consequences can be drawn from this : For a smooth simply connected surface  $S$ ,  $Br_n(S)$  is a birational invariant (it is the birational invariant  $F_n^{2,1}(\mathbb{C}(S)/\mathbb{C})$  of the foregoing section. In particular, if  $S = \mathbb{P}^2$  we have that  $Br_n(\mathbb{P}^2) = 0$  and as

$$0 \longrightarrow Br_n \mathbb{C}(x, y) \longrightarrow \bigoplus_C \mathbb{C}(C)^*/\mathbb{C}(C)^{*n} \longrightarrow \bigoplus_P \mu_n^{-1} \longrightarrow \mu_n \longrightarrow 0$$

we obtain a description of  $Br_n \mathbb{C}(x, y)$  by a certain geo-combinatorial package which we call a  $\mathbb{Z}_n$ -wrinkle over  $\mathbb{P}^2$ . A  $\mathbb{Z}_n$ -wrinkle is determined by

- A finite collection  $\mathcal{C} = \{C_1, \dots, C_k\}$  of *irreducible curves* in  $\mathbb{P}^2$ , that is,  $C_i = V(F_i)$  for an irreducible form in  $\mathbb{C}[X, Y, Z]$  of degree  $d_i$ .
- A finite collection  $\mathcal{P} = \{P_1, \dots, P_l\}$  of *points* of  $\mathbb{P}^2$  where each  $P_i$  is either an intersection point of two or more  $C_i$  or a singular point of some  $C_i$ .

- For each  $P \in \mathcal{P}$  the branch-data  $b_P = (b_1, \dots, b_{i_P})$  with  $b_i \in \mathbb{Z}_n = \mathbb{Z}/n\mathbb{Z}$  and  $\{1, \dots, i_P\}$  the different branches of  $\mathcal{C}$  in  $P$ . These numbers must satisfy the admissibility condition

$$\sum_i b_i = 0 \in \mathbb{Z}_n$$

for every  $P \in \mathcal{P}$


- for each  $C \in \mathcal{C}$  we fix a cyclic  $\mathbb{Z}_n$ -cover of smooth curves

$$D \longrightarrow \tilde{C}$$

of the desingularization  $\tilde{C}$  of  $C$  which is compatible with the branch-data. That is, if  $Q \in \tilde{C}$  corresponds to a  $\mathcal{C}$ -branch  $b_i$  in  $P$ , then  $D$  is ramified in  $Q$  with stabilizer subgroup

$$\text{Stab}_Q = \langle b_i \rangle \subset \mathbb{Z}_n$$

For example, a portion of a  $\mathbb{Z}_4$ -wrinkle can have the following picture



It is clear that the cover-data is the most untractable part of a  $\mathbb{Z}_n$ -wrinkle, so we want to have some control on the covers  $D \longrightarrow \tilde{C}$ . Let  $\{Q_1, \dots, Q_z\}$  be the points of  $\tilde{C}$  where the cover ramifies with branch numbers  $\{b_1, \dots, b_z\}$ , then  $D$  is determined by a continuous module structure (that is, a cofinite subgroup acts trivially) of

$$\pi_1(\tilde{C} - \{Q_1, \dots, Q_z\}) \text{ on } \mathbb{Z}_n$$

where the fundamental group of the Riemann surface  $\tilde{C}$  with  $z$  punctures is known (topologically) to be equal to the group

$$\langle u_1, v_1, \dots, u_g, v_g, x_1, \dots, x_z \rangle / ([u_1, v_1] \dots [u_g, v_g] x_1 \dots x_z)$$

where  $g$  is the genus of  $\tilde{C}$ . The action of  $x_i$  on  $\mathbb{Z}_n$  is determined by multiplication with  $b_i$ . In fact, we need to use the étale fundamental group, see [61], but this group has the same finite continuous modules as the topological fundamental group.

EXAMPLE 3.63. Covers of  $\mathbb{P}^1$  and elliptic curves.

- (1) If  $\tilde{C} = \mathbb{P}^1$  then  $g = 0$  and hence  $\pi_1(\mathbb{P}^1 - \{Q_1, \dots, Q_z\})$  is zero if  $z \leq 1$  (whence no covers exist) and is  $\mathbb{Z}$  if  $z = 2$ . Hence, there exists a unique cover  $D \longrightarrow \mathbb{P}^1$  with branch-data  $(1, -1)$  in say  $(0, \infty)$  namely with  $D$  the normalization of  $\mathbb{P}^1$  in  $\mathbb{C}(\sqrt{x})$ .
- (2) If  $\tilde{C} = E$  an elliptic curve, then  $g = 1$ . Hence,  $\pi_1(C) = \mathbb{Z} \oplus \mathbb{Z}$  and there exist unramified  $\mathbb{Z}_n$ -covers. They are given by the isogenies

$$E' \longrightarrow E$$

where  $E'$  is another elliptic curve and  $E = E'/\langle\tau\rangle$  where  $\tau$  is an  $n$ -torsion point on  $E'$ .

Any  $n$ -fold cover  $D \longrightarrow \tilde{C}$  is determined by a function  $f \in \mathbb{C}(C)^*/\mathbb{C}(C)^{*n}$ . This allows us to put a group-structure on the equivalence classes of  $\mathbb{Z}_n$ -wrinkles. In particular, we call a wrinkle *trivial* provided all coverings  $D_i \longrightarrow \tilde{C}_i$  are trivial (that is,  $D_i$  is the disjoint union of  $n$  copies of  $\tilde{C}_i$ ). The Artin-Mumford theorem for  $\mathbb{P}^2$  can now be stated as

**THEOREM 3.64.** *If  $\Delta$  is a central simple  $\mathbb{C}(x, y)$ -algebra of dimension  $n^2$ , then  $\Delta$  determines uniquely a  $\mathbb{Z}_n$ -wrinkle on  $\mathbb{P}^2$ . Conversely, any  $\mathbb{Z}_n$ -wrinkle on  $\mathbb{P}^2$  determines a unique division  $\mathbb{C}(x, y)$ - algebra whose class in the Brauer group has order  $n$ .*

**EXAMPLE 3.65.** If  $S$  is not necessarily simply connected, any class in  $Br(\mathbb{C}(S))_n$  still determines a  $\mathbb{Z}_n$ -wrinkle.

**EXAMPLE 3.66.** If  $X$  is a smooth irreducible rational projective variety of dimension  $d$ , the obstruction to classifying  $Br(\mathbb{C}(X))_n$  by  $\mathbb{Z}_n$ -wrinkles is given by  $H_{et}^3(X, \mu_n)$ .

We will give a ringtheoretical interpretation of the maps in the Artin-Mumford sequence. Observe that nearly all maps are those of the top complex of the first term in the coniveau spectral sequence for  $S$ . We gave an explicit description of them using discrete valuation rings. The statements below follow from this description.

Let us consider a discrete valuation ring  $A$  with field of fractions  $K$  and residue field  $k$ . Let  $\Delta$  be a central simple  $K$ -algebra of dimension  $n^2$ .

**DEFINITION 3.67.** An  $A$ -subalgebra  $\Lambda$  of  $\Delta$  will be called an  $A$ -order if it is a free  $A$ -module of rank  $n^2$  with  $\Lambda \cdot K = \Delta$ . An  $A$ -order is said to be *maximal* if it is not properly contained in any other order.

In order to study maximal orders in  $\Delta$  (they will turn out to be all conjugated), we consider the completion  $\hat{A}$  with respect to the  $m$ -adic filtration where  $m = At$  with  $t$  a uniformizing parameter of  $A$ .  $\hat{K}$  will denote the field of fractions of  $\hat{A}$  and  $\hat{\Delta} = \Delta \otimes_K \hat{K}$ .

Because  $\hat{\Delta}$  is a central simple  $\hat{K}$ -algebra of dimension  $n^2$  it is of the form

$$\hat{\Delta} = M_t(D)$$

where  $D$  is a division algebra with center  $\hat{K}$  of dimension  $s^2$  and hence  $n = s \cdot t$ . We call  $t$  the capacity of  $\Delta$  at  $A$ .

In  $D$  we can construct a unique maximal  $\hat{A}$ -order  $\Gamma$ , namely the integral closure of  $\hat{A}$  in  $D$ . We can view  $\Gamma$  as a discrete valuation ring extending the valuation  $v$  defined by  $A$  on  $K$ . If  $v : \hat{K} \longrightarrow \mathbb{Z}$ , then this extended valuation

$$w : D \longrightarrow n^{-2}\mathbb{Z} \text{ is defined as } w(a) = (\hat{K}(a) : \hat{K})^{-1}v(N_{\hat{K}(a)/\hat{K}}(a))$$

for every  $a \in D$  where  $\hat{K}(a)$  is the subfield generated by  $a$  and  $N$  is the norm map of fields.

The image of  $w$  is a subgroup of the form  $e^{-1}\mathbb{Z} \hookrightarrow n^{-2}\mathbb{Z}$ . The number  $e = e(D/\hat{K})$  is called the *ramification index* of  $D$  over  $\hat{K}$ . We can use it to normalize the valuation  $w$  to

$$v_D : D \longrightarrow \mathbb{Z} \text{ defined by } v_D(a) = \frac{e}{n^2}v(N_{D/\hat{K}}(a))$$

With these conventions we have that  $v_D(t) = e$ .

The maximal order  $\Gamma$  is then the subalgebra of all elements  $a \in D$  with  $v_D(a) \geq 0$ . It has a unique maximal ideal generated by a prime element  $T$  and we have that  $\bar{\Gamma} = \frac{\Gamma}{T\Gamma}$  is a division algebra finite dimensional over  $\hat{A}/t\hat{A} = k$  (but not necessarily having  $k$  as its center).

The *inertial degree* of  $D$  over  $\hat{K}$  is defined to be the number  $f = f(D/\hat{K}) = (\bar{\Gamma} : k)$  and one shows that

$$s^2 = e.f \text{ and } e \mid s \text{ whence } s \mid f$$

After this detour, we can now take  $\Lambda = M_t(\Gamma)$  as a maximal  $\hat{A}$ -order in  $\hat{\Delta}$ . One shows that all other maximal  $\hat{A}$ -orders are conjugated to  $\Lambda$ .  $\Lambda$  has a unique maximal ideal  $M$  with  $\bar{\Lambda} = M_t(\bar{\Gamma})$ .

DEFINITION 3.68. With notations as above, we call the numbers  $e = e(D/\hat{K})$ ,  $f = f(D/\hat{K})$  and  $t$  resp. the *ramification*, *inertia and capacity* of the central simple algebra  $\Delta$  at  $A$ . If  $e = 1$  we call  $\Lambda$  an Azumaya algebra over  $A$ , or equivalently, if  $\Lambda/t\Lambda$  is a central simple  $k$ -algebra of dimension  $n^2$ .

Now let us consider the case of a discrete valuation ring  $A$  in  $K$  such that the residue field  $k$  is  $Tsen^1$ . The center of the division algebra  $\bar{\Gamma}$  is a finite dimensional field extension of  $k$  and hence is also  $Tsen^1$  whence has trivial Brauer group and therefore must coincide with  $\bar{\Gamma}$ . Hence,

$$\bar{\Gamma} = k(\bar{a})$$

a commutative field, for some  $a \in \Gamma$ . But then,  $f \leq s$  and we have  $e = f = s$  and  $k(\bar{a})$  is a cyclic degree  $s$  field extension of  $k$ .

Because  $s \mid n$ , the cyclic extension  $k(\bar{a})$  determines an element of  $H_{et}^1(k, \mathbb{Z}_n)$ .

DEFINITION 3.69. Let  $Z$  be a normal domain with field of fractions  $K$  and let  $\Delta$  be a central simple  $K$ -algebra of dimension  $n^2$ . A  $Z$ -order  $B$  is a subalgebra which is a finitely generated  $Z$ -module. It is called maximal if it is not properly contained in any other order. One can show that  $B$  is a maximal  $Z$ -order if and only if  $\Lambda = B_p$  is a maximal order over the discrete valuation ring  $A = Z_p$  for every height one prime ideal  $p$  of  $Z$ .

Return to the situation of an irreducible smooth projective surface  $S$ . If  $\Delta$  is a central simple  $\mathbb{C}(S)$ -algebra of dimension  $n^2$ , we define a maximal order as a sheaf  $\mathcal{A}$  of  $\mathcal{O}_S$ -orders in  $\Delta$  which for an open affine cover  $U_i \hookrightarrow S$  is such that

$$\mathcal{A}_i = \Gamma(U_i, \mathcal{A}) \text{ is a maximal } Z_i = \Gamma(U_i, \mathcal{O}_S) \text{ order in } \Delta$$

Any irreducible curve  $C$  on  $S$  defines a discrete valuation ring on  $\mathbb{C}(S)$  with residue field  $\mathbb{C}(C)$  which is  $Tsen^1$ . Hence, the above argument can be applied to obtain from  $\mathcal{A}$  a cyclic extension of  $\mathbb{C}(C)$ , that is, an element of  $\mathbb{C}(C)^*/\mathbb{C}(C)^{*n}$ .

DEFINITION 3.70. We call the union of the curves  $C$  such that  $\mathcal{A}$  determines a non-trivial cyclic extension of  $\mathbb{C}(C)$  the *ramification divisor* of  $\Delta$  (or of  $\mathcal{A}$ ).

The map in the Artin-Mumford exact sequence

$$Br_n(\mathbb{C}(S)) \longrightarrow \bigoplus_C H_{et}^1(\mathbb{C}(C), \mu_n)$$

assigns to the class of  $\Delta$  the cyclic extensions introduced above.

DEFINITION 3.71. An  $S$ -Azumaya algebra (of index  $n$ ) is a sheaf of maximal orders in a central simple  $\mathbb{C}(S)$ -algebra  $\Delta$  of dimension  $n^2$  such that it is Azumaya at each curve  $C$ , that is, such that  $[\Delta]$  lies in the kernel of the above map.

Observe that this definition of Azumaya algebra coincides with the one given in the discussion of twisted forms of matrices. One can show that if  $\mathcal{A}$  and  $\mathcal{A}'$  are  $S$ -Azumaya algebras of index  $n$  resp.  $n'$ , then  $\mathcal{A} \otimes_{\mathcal{O}_S} \mathcal{A}'$  is an Azumaya algebra of index  $n.n'$ . We call an Azumaya algebra trivial if it is of the form  $End(\mathcal{P})$  where  $\mathcal{P}$  is a vectorbundle over  $S$ . The equivalence classes of  $S$ -Azumaya algebras can be given a group-structure called the Brauer-group  $Br(S)$  of the surface  $S$ .

### References.

More details on étale cohomology can be found in the textbook of J.S. Milne [61]. The classification of simple dimension vectors and invariants of quivers is due to L. Le Bruyn and C. Procesi [58]. The material of Tsen and Tate fields is based on the lecture notes of S. Shatz [74]. For more details on the coniveau spectral sequence we refer to the paper [12]. The description of the Brauer group of the functionfield of a surface is due to M. Artin and D. Mumford [5].





## CHAPTER 4

### Etale Slices.

#### 4.1. $C^\infty$ slices.

Let  $A$  be an affine  $\mathbb{C}$ -algebra and  $\xi \in \underline{iss}_n A$  a point in the quotient space corresponding to an  $n$ -dimensional semi-simple representation  $M_\xi$  of  $A$ . In this chapter we will present a method to study the étale local structure of  $\underline{iss}_n A$  near  $\xi$  and the étale local  $GL_n$ -structure of the representation variety  $\underline{rep}_n A$  near the closed orbit  $\mathcal{O}(M_\xi) = GL_n \cdot M_\xi$ . In this section we will outline the main idea in the setting of differential geometry.

Let  $M$  be a compact  $C^\infty$ -manifold on which a compact Lie group  $G$  acts differentially. By a usual averaging process we can define a  $G$ -invariant Riemannian metric on  $M$ . For a point  $m \in M$  we define

- The  $G$ -orbit  $\mathcal{O}(m) = G \cdot m$  of  $m$  in  $M$ ,
- the stabilizer subgroup  $H = \text{Stab}_G(m) = \{g \in G \mid g \cdot m = m\}$  and
- the normal space  $N_m$  defined to be the orthogonal complement to the tangent space in  $m$  to the orbit in the tangent space to  $M$ . That is, we have a decomposition of  $H$ -vectorspaces

$$T_m M = T_m \mathcal{O}(m) \oplus N_m$$

The normal spaces  $N_x$  when  $x$  varies over the points of the orbit  $\mathcal{O}(m)$  define a vectorbundle  $\mathcal{N} \xrightarrow{p} \mathcal{O}(m)$  over the orbit. We can identify the bundle with the associated fiber bundle


$$\mathcal{N} \simeq G \times^H N_m$$

Any point  $n \in \mathcal{N}$  in the normal bundle determines a geodesic

$$\gamma_n : \mathbb{R} \longrightarrow M \quad \text{defined by} \quad \begin{cases} \gamma_n(0) & = p(n) \\ \frac{d\gamma_n}{dt}(0) & = n \end{cases}$$

Using this geodesic we can define a  $G$ -equivariant exponential map from the normal bundle  $\mathcal{N}$  to the manifold  $M$  via

$$\mathcal{N} \xrightarrow{\text{exp}} M \quad \text{where} \quad \text{exp}(n) = \gamma_n(1)$$




Now, take  $\varepsilon > 0$  and define the  $C^\infty$  slice  $S_\varepsilon$  to be

$$S_\varepsilon = \{n \in N_m \mid \|n\| < \varepsilon\}$$

then  $G \times^H S_\varepsilon$  is a  $G$ -stable neighborhood of the zero section in the normal bundle  $\mathcal{N} = G \times^H N_m$ . But then we have a  $G$ -equivariant exponential

$$G \times^H S_\varepsilon \xrightarrow{\text{exp}} M$$

which for small enough  $\varepsilon$  gives a diffeomorphism with a  $G$ -stable tubular neighborhood  $U$  of the orbit  $\mathcal{O}(m)$  in  $M$ .



If we assume moreover that the action of  $G$  on  $M$  and the action of  $H$  on  $N_m$  are such that the orbit-spaces are manifolds  $M/G$  and  $N_m/H$ , then we have the situation

$$\begin{array}{ccccc} G \times^H S_\varepsilon & \xrightarrow[\simeq]{\text{exp}} & U & \hookrightarrow & M \\ \downarrow & & \downarrow & & \downarrow \\ S_\varepsilon/H & \xrightarrow[\simeq]{} & U/G & \hookrightarrow & M/G \end{array}$$

giving a local diffeomorphism between a neighborhood of  $\bar{0}$  in  $N_m/H$  and a neighborhood of the point  $\bar{m}$  in  $M/G$  corresponding to the orbit  $\mathcal{O}(m)$ .

Returning to the setting of the orbit  $\mathcal{O}(M_\xi)$  in  $\underline{rep}_n A$  we would equally like to define a  $GL_n$ -equivariant morphism from an associated fiber bundle

$$GL_n \times^{GL(\alpha)} N_\xi \xrightarrow{e} \underline{rep}_n A$$

where  $GL(\xi)$  is the stabilizer subgroup of  $M_\xi$  and  $N_\xi$  is a normal space to the orbit  $\mathcal{O}(M_\xi)$ . Because we do not have an exponential-map in the setting of algebraic geometry, the map  $e$  will have to be an étale map. Before we come to the description of these étale slices we will first study the tangent spaces to  $\underline{rep}_n A$  and give a ringtheoretical interpretation of the normal space  $N_\xi$ .

## 4.2. Normal spaces.

In this section we will determine the normal space to the orbit in a representation space. We recall some standard facts about tangent spaces first. Let  $\underline{X}$  be a not necessarily reduced affine variety with coordinate ring  $\mathbb{C}[\underline{X}] = \mathbb{C}[x_1, \dots, x_n]/I$ . If the origin  $o = (0, \dots, 0) \in \mathbb{V}(I)$ , elements of  $I$  have no constant terms and we can write any  $p \in I$  as

$$p = \sum_{i=1}^{\infty} p^{(i)} \quad \text{with } p^{(i)} \text{ homogeneous of degree } i.$$

The *order*  $ord(p)$  is the least integer  $r \geq 1$  such that  $p^{(r)} \neq 0$ . Define the following two ideals in  $\mathbb{C}[x_1, \dots, x_n]$

$$I_l = \{p^{(1)} \mid p \in I\} \quad \text{and } I_m = \{p^{(r)} \mid p \in I \text{ and } ord(p) = r\}.$$

The subscripts  $l$  (respectively  $m$ ) stand for *linear terms* (respectively, terms of *minimal* degree).

The *tangent space to  $\underline{X}$  in  $o$* ,  $T_o(\underline{X})$  is by definition the subscheme of  $\mathbb{C}^n$  determined by  $I_l$ . Observe that

$$I_l = (a_{11}x_1 + \dots + a_{1n}x_n, \dots, a_{l1}x_1 + \dots + a_{ln}x_n)$$

for some  $l \times n$  matrix  $A = (a_{ij})_{i,j}$  of rank  $l$ . That is, we can express all  $x_k$  as linear combinations of some  $\{x_{i_1}, \dots, x_{i_{n-l}}\}$ , but then clearly

$$\mathbb{C}[T_o(\underline{X})] = \mathbb{C}[x_1, \dots, x_n]/I_l = \mathbb{C}[x_{i_1}, \dots, x_{i_{n-l}}]$$

In particular,  $T_o(\underline{X})$  is reduced and is a linear subspace of dimension  $n - l$  in  $\mathbb{C}^n$  through the point  $o$ .

Next, consider an arbitrary geometric point  $x$  of  $\underline{X}$  with coordinates  $(a_1, \dots, a_n)$ . We can translate  $x$  to the origin  $o$  and the translate of  $\underline{X}$  is then the scheme defined by the ideal

$$(f_1(x_1 + a_1, \dots, x_n + a_n), \dots, f_k(x_1 + a_1, \dots, x_n + a_n))$$

Now, the linear term of the translated polynomial  $f_i(x_1 + a_1, \dots, x_n + a_n)$  is equal to

$$\frac{\partial f_i}{\partial x_1}(a_1, \dots, a_n)x_1 + \dots + \frac{\partial f_i}{\partial x_n}(a_1, \dots, a_n)x_n$$

and hence the *tangent space to  $\underline{X}$  in  $x$* ,  $T_x(\underline{X})$  is the linear subspace of  $\mathbb{C}^n$  defined by the set of zeroes of the linear terms

$$T_x(\underline{X}) = \mathbb{V}\left(\sum_{j=1}^n \frac{\partial f_1}{\partial x_j}(x)x_j, \dots, \sum_{j=1}^n \frac{\partial f_k}{\partial x_j}(x)x_j\right) \hookrightarrow \mathbb{C}^n.$$

In particular, the dimension of this linear subspace can be computed from the *Jacobian matrix* in  $x$  associated with the polynomials  $(f_1, \dots, f_k)$

$$\dim T_x(\underline{X}) = n - rk \begin{bmatrix} \frac{\partial f_1}{\partial x_1}(x) & \dots & \frac{\partial f_1}{\partial x_n}(x) \\ \vdots & & \vdots \\ \frac{\partial f_k}{\partial x_1}(x) & \dots & \frac{\partial f_k}{\partial x_n}(x) \end{bmatrix}.$$

Let  $\mathbb{C}[\varepsilon]$  be the *algebra of dual numbers*, that is,  $\mathbb{C}[\varepsilon] \simeq \mathbb{C}[y]/(y^2)$ . Consider a  $\mathbb{C}$ -algebra morphism

$$\mathbb{C}[x_1, \dots, x_n] \xrightarrow{\phi} \mathbb{C}[\varepsilon] \quad \text{defined by } x_i \mapsto a_i + c_i \varepsilon.$$

Because  $\varepsilon^2 = 0$  it is easy to verify that the image of a polynomial  $f(x_1, \dots, x_n)$  under  $\phi$  is of the form

$$\phi(f(x_1, \dots, x_n)) = f(a_1, \dots, a_n) + \sum_{j=1}^n \frac{\partial f}{\partial x_j}(a_1, \dots, a_n) c_j \varepsilon$$

Therefore,  $\phi$  factors through  $I$ , that is  $\phi(f_i) = 0$  for all  $1 \leq i \leq k$ , if and only if  $(c_1, \dots, c_n) \in T_x(\underline{X})$ . Hence, we can also identify the tangent space to  $\underline{X}$  in  $x$  with the algebra morphisms  $\mathbb{C}[\underline{X}] \xrightarrow{\phi} \mathbb{C}[\varepsilon]$  whose composition with the projection  $\pi : \mathbb{C}[\varepsilon] \longrightarrow \mathbb{C}$  (sending  $\varepsilon$  to zero) is the evaluation in  $x = (a_1, \dots, a_n)$ . That is, let  $ev_x \in \underline{X}(\mathbb{C})$  be the point corresponding to evaluation in  $x$ , then

$$T_x(\underline{X}) = \{\phi \in \underline{X}(\mathbb{C}[\varepsilon]) \mid \underline{X}(\pi)(\phi) = ev_x\}.$$

The following two examples compute the tangent spaces to the (trace preserving) representation varieties.

**EXAMPLE 4.1.** Let  $A$  be an affine  $\mathbb{C}$ -algebra generated by  $\{a_1, \dots, a_m\}$  and  $\rho : A \longrightarrow M_n(\mathbb{C})$  an algebra morphism, that is,  $\rho \in \text{rep}_n A$ . We call a linear map  $A \xrightarrow{D} M_n(\mathbb{C})$  a  $\rho$ -*derivation* if and only if for all  $a, a' \in A$  we have that

$$D(aa') = D(a) \cdot \rho(a') + \rho(a) \cdot D(a').$$

We denote the vectorspace of all  $\rho$ -derivations of  $A$  by  $\text{Der}_\rho(A)$ . Observe that any  $\rho$ -derivation is determined by its image on the generators  $a_i$ , hence  $\text{Der}_\rho(A) \subset M_n^m$ . We claim that

$$T_\rho(\text{rep}_n A) = \text{Der}_\rho(A).$$

Indeed, we know that  $\text{rep}_n A(\mathbb{C}[\varepsilon])$  is the set of algebra morphisms

$$A \xrightarrow{\phi} M_n(\mathbb{C}[\varepsilon])$$

By the functorial characterization of tangentspaces we have that  $T_\rho(\text{rep}_n A)$  is equal to

$$\{D : A \longrightarrow M_n(\mathbb{C}) \text{ linear} \mid \rho + D\varepsilon : A \longrightarrow M_n(\mathbb{C}[\varepsilon]) \text{ is an algebra map}\}.$$

Because  $\rho$  is an algebra morphism, the algebra map condition

$$\rho(aa') + D(aa')\varepsilon = (\rho(a) + D(a)\varepsilon) \cdot (\rho(a') + D(a')\varepsilon)$$

is equivalent to  $D$  being a  $\rho$ -derivation.

**EXAMPLE 4.2.** Let  $A$  be a Cayley-Hamilton algebra of degree  $n$  with trace map  $tr_A$  and trace generated by  $\{a_1, \dots, a_m\}$ . Let  $\rho \in \text{rep}_n^{tr} A$ , that is,  $\rho : A \longrightarrow M_n(\mathbb{C})$  is a *trace preserving* algebra morphism. Because  $\text{rep}_n^{tr} A(\mathbb{C}[\varepsilon])$  is the set of all trace preserving algebra morphisms  $A \longrightarrow M_n(\mathbb{C}[\varepsilon])$  (with the usual trace map  $tr$  on  $M_n(\mathbb{C}[\varepsilon])$ ) and the previous example one verifies that

$$T_\rho(\text{rep}_n^{tr} A) = \text{Der}_\rho^{tr}(A) \subset \text{Der}_\rho(A)$$

the subset of *trace preserving  $\rho$ -derivations*  $D$ , that is, those satisfying

$$D \circ tr_A = tr \circ D$$

$$\begin{array}{ccc} A & \xrightarrow{D} & M_n(\mathbb{C}) \\ \downarrow tr_A & & \downarrow tr \\ A & \xrightarrow{D} & M_n(\mathbb{C}) \end{array}$$

Again, using this property and the fact that  $A$  is *trace* generated by  $\{a_1, \dots, a_m\}$  a trace preserving  $\rho$ -derivation is determined by its image on the  $a_i$  so is a subspace of  $M_n^m$ .

The *tangent cone to  $\underline{X}$  in  $o$* ,  $TC_o(\underline{X})$ , is by definition the subscheme of  $\mathbb{C}^n$  determined by  $I_m$ , that is,

$$\mathbb{C}[TC_o(\underline{X})] = \mathbb{C}[x_1, \dots, x_n]/I_m.$$

It is called a *cone* because if  $c$  is a point of the underlying variety of  $TC_o(\underline{X})$ , then the line  $l = \overrightarrow{oc}$  is contained in this variety because  $I_m$  is a graded ideal. Further, observe that as  $I_l \subset I_m$ , the tangent cone is a closed subscheme of the tangent space at  $\underline{X}$  in  $o$ . Again, if  $x$  is an arbitrary geometric point of  $\underline{X}$  we define the *tangent cone to  $\underline{X}$  in  $x$* ,  $TC_x(\underline{X})$  as the tangent cone  $TC_o(\underline{X}')$  where  $\underline{X}'$  is the translated scheme of  $\underline{X}$  under the translation taking  $x$  to  $o$ . Both the tangent space and tangent cone contain *local information* of the scheme  $\underline{X}$  in a neighborhood of  $x$ .

Let  $m_x$  be the maximal ideal of  $\mathbb{C}[\underline{X}]$  corresponding to  $x$  (that is, the ideal of polynomial functions vanishing in  $x$ ). Then, its complement  $\mathcal{S}_x = \mathbb{C}[\underline{X}] - m_x$  is a multiplicatively closed subset and the *local algebra*  $\mathcal{O}_x(\underline{X})$  is the corresponding localization  $\mathbb{C}[\underline{X}]_{\mathcal{S}_x}$ . It has a unique maximal ideal  $\mathfrak{m}_x$  with residue field  $\mathcal{O}_x(\underline{X})/\mathfrak{m}_x = \mathbb{C}$ . We equip the local algebra  $\mathcal{O}_x = \mathcal{O}_x(\underline{X})$  with the  $\mathfrak{m}_x$ -adic filtration that is the increasing  $\mathbb{Z}$ -filtration

$$\mathcal{F}_x : \quad \dots \subset \mathfrak{m}^i \subset \mathfrak{m}^{i-1} \subset \dots \subset \mathfrak{m} \subset \mathcal{O}_x = \mathcal{O}_x = \dots = \mathcal{O}_x = \dots$$

with *associated graded algebra*

$$gr(\mathcal{O}_x) = \quad \dots \oplus \frac{\mathfrak{m}_x^i}{\mathfrak{m}_x^{i+1}} \oplus \frac{\mathfrak{m}_x^{i-1}}{\mathfrak{m}_x^i} \oplus \dots \oplus \frac{\mathfrak{m}_x}{\mathfrak{m}_x^2} \oplus \mathbb{C} \oplus 0 \oplus \dots \oplus 0 \oplus \dots$$

PROPOSITION 4.3. *If  $x$  is a geometric point of the affine scheme  $\underline{X}$ , then*

- (1)  $\mathbb{C}[T_x(\underline{X})]$  is isomorphic to the polynomial algebra  $\mathbb{C}[\frac{\mathfrak{m}_x}{\mathfrak{m}_x^2}]$ .
- (2)  $\mathbb{C}[TC_x(\underline{X})]$  is isomorphic to the associated graded algebra  $gr(\mathcal{O}_x(\underline{X}))$ .

PROOF. After translating we may assume that  $x = o$  lies in  $\mathbb{V}(I) \hookrightarrow \mathbb{C}^n$ . That is,

$$\mathbb{C}[\underline{X}] = \mathbb{C}[x_1, \dots, x_n]/I \quad \text{and} \quad m_x = (x_1, \dots, x_n)/I.$$

(1) : Under these identifications we have

$$\begin{aligned} \frac{\mathfrak{m}_x}{\mathfrak{m}_x^2} &\simeq \frac{m_x}{m_x^2} \\ &\simeq \frac{(x_1, \dots, x_n)}{(x_1, \dots, x_n)^2 + I} \\ &\simeq \frac{(x_1, \dots, x_n)}{(x_1, \dots, x_n)^2 + I_l} \end{aligned}$$

and as  $I_l$  is generated by linear terms it follows that the polynomial algebra on  $\frac{\mathfrak{m}_x}{\mathfrak{m}_x^2}$  is isomorphic to the quotient algebra  $\mathbb{C}[x_1, \dots, x_n]/I_l$  which is by definition the coordinate ring of the tangent space.

(2) : Again using the above identifications we have

$$\begin{aligned}
gr(\mathcal{O}_x) &\simeq \bigoplus_{i=0}^{\infty} \frac{\mathfrak{m}_x^i}{\mathfrak{m}_x^{i+1}} \\
&\simeq \bigoplus_{i=0}^{\infty} \frac{\mathfrak{m}_x^i}{\mathfrak{m}_x^{i+1}} \\
&\simeq \bigoplus_{i=0}^{\infty} \frac{(x_1, \dots, x_n)^i}{(x_1, \dots, x_n)^{i+1} + (I \cap (x_1, \dots, x_n)^i)} \\
&\simeq \bigoplus_{i=0}^{\infty} \frac{(x_1, \dots, x_n)^i}{(x_1, \dots, x_n)^{i+1} + I_m(i)}
\end{aligned}$$

where  $I_m(i)$  is the homogeneous part of  $I_m$  of degree  $i$ . On the other hand, the  $i$ -th homogeneous part of  $\mathbb{C}[x_1, \dots, x_n]/I_m$  is equal to

$$\frac{(x_1, \dots, x_n)^i}{(x_1, \dots, x_n)^{i+1} + I_m(i)}$$

we obtain the required isomorphism.  $\square$

This gives a third interpretation of the tangent space as

$$T_x(\underline{X}) = \text{Hom}_{\mathbb{C}}\left(\frac{\mathfrak{m}_x}{\mathfrak{m}_x^2}, \mathbb{C}\right) = \text{Hom}_{\mathbb{C}}\left(\frac{\mathfrak{m}_x}{\mathfrak{m}_x^2}, \mathbb{C}\right).$$

Hence, we can also view the tangent space  $T_x(\underline{X})$  as the space of *point derivations*  $Der_x(\mathcal{O}_x)$  on  $\mathcal{O}_x(\underline{X})$  (or of the point derivations  $Der_x(\mathbb{C}[\underline{X}])$  on  $\mathbb{C}[\underline{X}]$ ). That is,  $\mathbb{C}$ -linear maps  $D : \mathcal{O}_x \longrightarrow \mathbb{C}$  (or  $D : \mathbb{C}[\underline{X}] \longrightarrow \mathbb{C}$ ) such that for all functions  $f, g$  we have

$$D(fg) = D(f)g(x) + f(x)D(g).$$

If we define the *local dimension of an affine scheme  $\underline{X}$  in a geometric point  $x$*   $\dim_x \underline{X}$  to be the maximal dimension of irreducible components of the reduced variety  $X$  passing through  $x$ , then

$$\dim_x \underline{X} = \dim_o TC_x(\underline{X}).$$

We say that  $\underline{X}$  is *nonsingular at  $x$*  (or equivalently, that  $x$  is a *nonsingular point of  $\underline{X}$* ) if the tangent cone to  $\underline{X}$  in  $x$  coincides with the tangent space to  $\underline{X}$  in  $x$ . An immediate consequence is

**PROPOSITION 4.4.** *If  $\underline{X}$  is nonsingular at  $x$ , then  $\mathcal{O}_x(\underline{X})$  is a domain. That is, in a Zariski neighborhood of  $x$ ,  $\underline{X}$  is an irreducible variety.*

**PROOF.** If  $\underline{X}$  is nonsingular at  $x$ , then

$$gr(\mathcal{O}_x) \simeq \mathbb{C}[TC_x(\underline{X})] = \mathbb{C}[T_x(\underline{X})]$$

the latter one being a polynomial algebra whence a domain. Now, let  $0 \neq a, b \in \mathcal{O}_x$  then there exist  $k, l$  such that  $a \in \mathfrak{m}^k - \mathfrak{m}^{k+1}$  and  $b \in \mathfrak{m}^l - \mathfrak{m}^{l+1}$ , that is  $\bar{a}$  is a nonzero homogeneous element of  $gr(\mathcal{O}_x)$  of degree  $-k$  and  $\bar{b}$  one of degree  $-l$ . But then,  $\bar{a}\bar{b} \in \mathfrak{m}^{k+l} - \mathfrak{m}^{k+l-1}$  hence certainly  $a \cdot b \neq 0$  in  $\mathcal{O}_x$ .

Now, consider the natural map  $\phi : \mathbb{C}[\underline{X}] \longrightarrow \mathcal{O}_x$ . Let  $\{P_1, \dots, P_l\}$  be the minimal prime ideals of  $\mathbb{C}[\underline{X}]$ . All but one of them, say  $P_1 = \phi^{-1}(0)$ , extend to the whole ring  $\mathcal{O}_x$ . Taking the product of  $l$  functions  $f_i \in P_i$  nonvanishing in  $x$  for  $2 \leq i \leq l$  gives the Zariski open set  $\mathbb{X}(f)$  containing  $x$  and whose coordinate ring is a domain, whence  $\mathbb{X}(f)$  is an affine irreducible variety.  $\square$

When restricting to nonsingular points we reduce to irreducible affine varieties. From the Jacobian condition it follows that nonsingularity is a Zariski open condition on  $\underline{X}$  and by the implicit function theorem,  $\underline{X}$  is a complex manifold in a neighborhood of a nonsingular point.

Let  $\underline{X} \xrightarrow{\phi} \underline{Y}$  be a morphism of affine varieties corresponding to the algebra morphism  $\mathbb{C}[\underline{Y}] \xrightarrow{\phi^*} \mathbb{C}[\underline{X}]$ . Let  $x$  be a geometric point of  $\underline{X}$  and  $y = \phi(x)$ . As  $\phi^*(m_y) \subset m_x$ ,  $\phi$  induces a linear map  $\frac{m_y}{m_y^2} \longrightarrow \frac{m_x}{m_x^2}$  and taking the dual map gives the *differential of  $\phi$  in  $x$*  which is a linear map

$$d\phi_x : T_x(\underline{X}) \longrightarrow T_{\phi(x)}(\underline{Y}).$$

Assume  $\underline{X}$  a closed subscheme of  $\mathbb{C}^n$  and  $\underline{Y}$  a closed subscheme of  $\mathbb{C}^m$  and let  $\phi$  be determined by the  $m$  polynomials  $\{f_1, \dots, f_m\}$  in  $\mathbb{C}[x_1, \dots, x_n]$ . Then, the Jacobian matrix in  $x$

$$J_x(\phi) = \begin{bmatrix} \frac{\partial f_1}{\partial x_1}(x) & \dots & \frac{\partial f_m}{\partial x_1}(x) \\ \vdots & & \vdots \\ \frac{\partial f_1}{\partial x_n}(x) & \dots & \frac{\partial f_m}{\partial x_n}(x) \end{bmatrix}$$

defines a linear map from  $\mathbb{C}^n$  to  $\mathbb{C}^m$  and the differential  $d\phi_x$  is the induced linear map from  $T_x(\underline{X}) \subset \mathbb{C}^n$  to  $T_{\phi(x)}(\underline{Y}) \subset \mathbb{C}^m$ . Let  $D \in T_x(\underline{X}) = \text{Der}_x(\mathbb{C}[\underline{X}])$  and  $x_D$  the corresponding element of  $\underline{X}(\mathbb{C}[\varepsilon])$  defined by  $x_D(f) = f(x) + D(f)\varepsilon$ , then  $x_D \circ \phi^* \in \underline{Y}(\mathbb{C}[\varepsilon])$  is defined by

$$x_D \circ \phi^*(g) = g(\phi(x)) + (D \circ \phi^*)\varepsilon = g(\phi(x)) + d\phi_x(D)\varepsilon$$

giving us the  $\varepsilon$ -interpretation of the differential

$$\phi(x + v\varepsilon) = \phi(x) + d\phi_x(v)\varepsilon$$

for all  $v \in T_x(\underline{X})$ .

**PROPOSITION 4.5.** *Let  $X \xrightarrow{\phi} Y$  be a dominant morphism between irreducible affine varieties. There is a Zariski open dense subset  $U \hookrightarrow X$  such that  $d\phi_x$  is surjective for all  $x \in U$ .*

**PROOF.** We may assume that  $\phi$  factorizes into

$$\begin{array}{ccc} X & \xrightarrow{\rho} & Y \times \mathbb{C}^d \\ & \searrow \phi & \downarrow \text{pr}_Y \\ & & Y \end{array}$$

with  $\phi$  a finite and surjective morphism. Because the tangent space of a product is the sum of the tangent spaces of the components we have that  $d(\text{pr}_W)_z$  is surjective for all  $z \in Y \times \mathbb{C}^d$ , hence it suffices to verify the claim for a *finite* morphism  $\phi$ . That is, we may assume that  $S = \mathbb{C}[Y]$  is a finite module over  $R = \mathbb{C}[X]$  and let  $L/K$  be the corresponding extension of the function fields. By the *principal element theorem* we know that  $L = K[s]$  for an element  $s \in L$  which is integral over  $R$  with minimal polynomial

$$F = t^n + g_{n-1}t^{n-1} + \dots + g_1t + g_0 \quad \text{with } g_i \in R$$

Consider the ring  $S' = R[t]/(F)$  then there is an element  $r \in R$  such that the localizations  $S'_r$  and  $S_r$  are isomorphic. By restricting we may assume that  $X = \mathbb{V}(F) \hookrightarrow Y \times \mathbb{C}$  and that

$$\begin{array}{ccc} X = \mathbb{V}(F) & \hookrightarrow & Y \times \mathbb{C} \\ & \searrow \varnothing & \downarrow pr_Y \\ & & Y \end{array}$$

Let  $x = (y, c) \in X$  then we have (again using the identification of the tangent space of a product with the sum of the tangent spaces of the components) that

$$T_x(X) = \{(v, a) \in T_y(Y) \oplus \mathbb{C} \mid c \frac{\partial F}{\partial t}(x) + vg_{n-1}c^{n-1} + \dots + vg_1c + vg_0 = 0\}.$$

But then,  $d\phi_x$  is surjective whenever  $\frac{\partial F}{\partial t}(x) \neq 0$ . This condition determines a *non-empty* open subset of  $X$  as otherwise  $\frac{\partial F}{\partial t}$  would belong to the defining ideal of  $X$  in  $\mathbb{C}[Y \times \mathbb{C}]$  (which is the principal ideal generated by  $F$ ) which is impossible by a degree argument  $\square$

EXAMPLE 4.6. Let  $\underline{X}$  be a closed  $GL_n$ -stable subscheme of a  $GL_n$ -representation  $V$  and  $x$  a geometric point of  $\underline{X}$ . Consider the orbitclosure  $\overline{\mathcal{O}(x)}$  of  $x$  in  $V$ . Because the orbit map

$$\mu : GL_n \longrightarrow GL_n \cdot x \hookrightarrow \overline{\mathcal{O}(x)}$$

is dominant we have that  $\mathbb{C}[\overline{\mathcal{O}(x)}] \hookrightarrow \mathbb{C}[GL_n]$  and therefore a domain, so  $\overline{\mathcal{O}(x)}$  is an irreducible affine variety. We define the *stabilizer subgroup*  $Stab(x)$  to be the fiber  $\mu^{-1}(x)$ , then  $Stab(x)$  is a closed subgroup of  $GL_n$ . We claim that the differential of the orbit map in the identity matrix  $e = \mathbb{1}_n$

$$d\mu_e : \mathfrak{gl}_n \longrightarrow T_x(\underline{X})$$

satisfies the following properties

$$Ker d\mu_e = \mathfrak{stab}(x) \quad \text{and} \quad Im d\mu_e = T_x(\overline{\mathcal{O}(x)}).$$

By the proposition we know that there is a dense open subset  $U$  of  $GL_n$  such that  $d\mu_g$  is surjective for all  $g \in U$ . By  $GL_n$ -equivariance of  $\mu$  it follows that  $d\mu_g$  is surjective for all  $g \in GL_n$ , in particular  $d\mu_e : \mathfrak{gl}_n \longrightarrow T_x(\overline{\mathcal{O}(x)})$  is surjective. Further, all fibers of  $\mu$  over  $\mathcal{O}(x)$  have the same dimension. But then it follows from the *dimension formula* of proposition that

$$dim GL_n = dim Stab(x) + dim \overline{\mathcal{O}(x)}$$

(which, incidentally gives us an algorithm to compute the dimensions of orbitclosures). Combining this with the above surjectivity, a dimension count proves that  $Ker d\mu_e = \mathfrak{stab}(x)$ , the Lie algebra of  $Stab(x)$ .

EXAMPLE 4.7. Let  $A$  be an affine  $\mathbb{C}$ -algebra generated by  $\{a_1, \dots, a_m\}$  and  $\rho : A \longrightarrow M_n(\mathbb{C})$  an algebra morphism, that is,  $\rho \in rep_n A$  determines an  $n$ -dimensional  $A$ -representation  $M$ . We claim to have the following description of the *normal space* to the orbitclosure  $C_\rho = \overline{\mathcal{O}(\rho)}$  of  $\rho$

$$N_\rho(rep_n A) \stackrel{def}{=} \frac{T_\rho(rep_n A)}{T_\rho(C_\rho)} = Ext_A^1(M, M).$$

We have already seen that the space of cycles  $Z(M, M)$  is the space of  $\rho$ -derivations of  $A$  in  $M_n(\mathbb{C})$ ,  $Der_\rho(A)$ , which we know to be the tangent space  $T_\rho(rep_n A)$ . Moreover, we know that the differential  $d\mu_e$  of the orbit map  $GL_n \xrightarrow{\mu} C_\rho \hookrightarrow M_n^m$

$$d\mu_e : \mathfrak{gl}_n = M_n \longrightarrow T_\rho(C_\rho)$$



is surjective. Now,  $\rho = (\rho(a_1), \dots, \rho(a_m)) \in M_n^m$  and the action of action of  $GL_n$  is given by simultaneous conjugation. But then we have for any  $A \in \mathfrak{gl}_n = M_n$  that

$$(I_n + A\varepsilon) \cdot \rho(a_i) \cdot (I_n - A\varepsilon) = \rho(a_i) + (A\rho(a_i) - \rho(a_i)A)\varepsilon.$$

Therefore, by definition of the differential we have that

$$d\mu_\varepsilon(A)(a) = A\rho(a) - \rho(a)A \quad \text{for all } a \in A.$$

that is,  $d\mu_\varepsilon(A) \in B(M, M)$  and as the differential map is surjective we have  $T_\rho(C_\rho) = B(M, M)$  from which the claim follows.

**EXAMPLE 4.8.** Let  $A$  be a Cayley-Hamilton algebra with trace map  $tr_A$  and trace generated by  $\{a_1, \dots, a_m\}$ . Let  $\rho \in \text{rep}_n^{tr} A$ , that is,  $\rho : A \rightarrow M_n(\mathbb{C})$  is a trace preserving algebra morphism. Any cycle  $\lambda : A \rightarrow M_n(\mathbb{C})$  in  $Z(M, M) = \text{Der}_\rho(A)$  determines an algebra morphism

$$\rho + \lambda\varepsilon : A \rightarrow M_n(\mathbb{C}[\varepsilon])$$

We know that the tangent space  $T_\rho(\text{rep}_n^{tr} A)$  is the subspace  $\text{Der}_\rho^{tr}(A)$  of trace preserving  $\rho$ -derivations, that is, those satisfying

$$\lambda(tr_A(a)) = tr(\lambda(a)) \quad \text{for all } a \in A$$

Observe that for all boundaries  $\delta \in B(M, M)$ , that is, such that there is an  $m \in M_n(\mathbb{C})$  with  $\delta(a) = \rho(a) \cdot m - m \cdot \rho(a)$  are trace preserving as

$$\begin{aligned} \delta(tr_A(a)) &= \rho(tr_A(a)) \cdot m - m \cdot \rho(tr_A(a)) = tr(\rho(a)) \cdot m - m \cdot tr(\rho(a)) \\ &= 0 = tr(m \cdot \rho(a) - \rho(a) \cdot m) = tr(\delta(a)) \end{aligned}$$

Hence, we can define the space of *trace preserving self-extensions*

$$\text{Ext}_A^{tr}(M, M) = \frac{\text{Der}_\rho^{tr}(A)}{B(M, M)}$$

and obtain as before that the *normal space* to the orbit closure  $C_\rho = \overline{\mathcal{O}(\rho)}$  is equal to

$$N_\rho(\text{rep}_n^{tr} A) \stackrel{\text{def}}{=} \frac{T_\rho(\text{rep}_n^{tr} A)}{T_\rho(C_\rho)} = \text{Ext}_A^{tr}(M, M)$$

### 4.3. Knop-Luna slices.

The results of this section hold for any reductive algebraic group  $G$ . We will apply them only in the case  $G = GL_n$  or  $GL(\alpha) = GL_{a_1} \times \dots \times GL_{a_k}$ , so restrict to the case of  $GL_n$ . We fix the setting :  $\underline{X}$  and  $\underline{Y}$  are (not necessarily reduced) affine  $GL_n$ -varieties,  $\psi$  is a  $GL_n$ -equivariant map

$$\begin{array}{ccc} x = \psi(y) \vdash & \underline{X} & \xleftarrow{\psi} & \underline{Y} & \dashv & y \\ & \downarrow \pi_X & & \downarrow \pi_Y & & \\ & \underline{X}/GL_n & & \underline{Y}/GL_n & & \end{array}$$

and we assume the following restrictions :

- $\psi$  is étale in  $y$ ,
- the  $GL_n$ -orbits  $\mathcal{O}(y)$  in  $\underline{Y}$  and  $\mathcal{O}(x)$  in  $\underline{X}$  are closed. For example, in representation varieties, we restrict to semi-simple representations,
- the stabilizer subgroups are equal  $\text{Stab}(x) = \text{Stab}(y)$ . In the case of representation varieties, for a semi-simple  $n$ -dimensional representation with decomposition

$$M = S_1^{\oplus e_1} \oplus \dots \oplus S_k^{\oplus e_k}$$

into distinct simple components, this stabilizer subgroup is

$$GL(\alpha) = \begin{bmatrix} GL_{e_1}(\mathbb{C} \otimes \mathbb{1}_{d_1}) & & \\ & \ddots & \\ & & GL_{e_k}(\mathbb{C} \otimes \mathbb{1}_{d_k}) \end{bmatrix} \hookrightarrow GL_n$$

where  $d_i = \dim S_i$ . In particular, the stabilizer subgroup is again reductive.

In algebraic terms : consider the coordinate rings  $R = \mathbb{C}[\underline{X}]$  and  $S = \mathbb{C}[\underline{Y}]$  and the dual morphism  $R \xrightarrow{\psi^*} S$ . Let  $I \triangleleft R$  be the ideal describing  $\mathcal{O}(x)$  and  $J \triangleleft S$  the ideal describing  $\mathcal{O}(y)$ . With  $\widehat{R}$  we will denote the  $I$ -adic completion  $\varprojlim \frac{R}{I^n}$  of  $R$  and with  $\widehat{S}$  the  $J$ -adic completion of  $S$ .

LEMMA 4.9. *The morphism  $\psi^*$  induces for all  $n$  an isomorphism*

$$\frac{R}{I^n} \xrightarrow{\psi^*} \frac{S}{J^n}$$

In particular,  $\widehat{R} \simeq \widehat{S}$ .

PROOF. Let  $\underline{Z}$  be the closed  $GL_n$ -stable subvariety of  $\underline{Y}$  where  $\psi$  is not étale. By the separation property, there is an invariant function  $f \in S^{GL_n}$  vanishing on  $\underline{Z}$  such that  $f(y) = 1$  because the two closed  $GL_n$ -subschemes  $\underline{Z}$  and  $\mathcal{O}(y)$  are disjoint. Replacing  $S$  by  $S_f$  we may assume that  $\psi^*$  is an étale morphism. Because  $\mathcal{O}(x)$  is smooth,  $\psi^{-1} \mathcal{O}(x)$  is the disjoint union of its irreducible components and restricting  $\underline{Y}$  if necessary we may assume that  $\psi^{-1} \mathcal{O}(x) = \mathcal{O}(y)$ . But then  $J = \psi^*(I)S$  and as  $\mathcal{O}(y) \xrightarrow{\simeq} \mathcal{O}(x)$  we have  $\frac{R}{I} \simeq \frac{S}{J}$  so the result holds for  $n = 1$ .

Because étale maps are flat, we have  $\psi^*(I^n)S = I^n \otimes_R S = J^n$  and an exact sequence

$$0 \longrightarrow I^{n+1} \otimes_R S \longrightarrow I^n \otimes_R S \longrightarrow \frac{I^n}{I^{n+1}} \otimes_R S \longrightarrow 0$$

But then we have

$$\frac{I^n}{I^{n+1}} = \frac{I^n}{I^{n+1}} \otimes_{R/I} \frac{S}{J} = \frac{I^n}{I^{n+1}} \otimes_R S \simeq \frac{J^n}{J^{n+1}}$$

and the result follows from induction on  $n$  and the commuting diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \frac{I^n}{I^{n+1}} & \longrightarrow & \frac{R}{I^{n+1}} & \longrightarrow & \frac{R}{I^n} \longrightarrow 0 \\ & & \downarrow \simeq & & \vdots & & \downarrow \simeq \\ 0 & \longrightarrow & \frac{J^n}{J^{n+1}} & \longrightarrow & \frac{S}{J^{n+1}} & \longrightarrow & \frac{S}{J^n} \longrightarrow 0 \end{array}$$

□

For an irreducible  $GL_n$ -representation  $s$  and a locally finite  $GL_n$ -module  $X$  we denote its  $s$ -isotypical component by  $X_{(s)}$ .

LEMMA 4.10. *Let  $s$  be an irreducible  $GL_n$ -representation. There are natural numbers  $m \geq 1$  (independent of  $s$ ) and  $n \geq 0$  such that for all  $k \in \mathbb{N}$  we have*

$$I^{mk+n} \cap R_{(s)} \hookrightarrow (I^{GL_n})^k R_{(s)} \hookrightarrow I^k \cap R_{(s)}$$

PROOF. Consider  $A = \bigoplus_{i=0}^{\infty} I^n t^i \hookrightarrow R[t]$ , then  $A^{GL_n}$  is affine so certainly finitely generated as  $R^{GL_n}$ -algebra say by

$$\{r_1 t^{m_1}, \dots, r_z t^{m_z}\} \quad \text{with } r_i \in R \text{ and } m_i \geq 1.$$

Further,  $A_{(s)}$  is a finitely generated  $A^{GL_n}$ -module, say generated by

$$\{s_1 t^{n_1}, \dots, s_y t^{n_y}\} \quad \text{with } s_i \in R_{(s)} \text{ and } n_i \geq 0.$$

Take  $m = \max m_i$  and  $n = \max n_i$  and  $r \in I^{m+k+n} \cap R_{(s)}$ , then  $rt^{m+k+n} \in A_{(s)}$  and

$$rt^{m+k+n} = \sum_j p_j(r_1 t^{m_1}, \dots, r_z t^{m_z}) s_j t^{n_j}$$

with  $p_j$  a homogeneous polynomial of  $t$ -degree  $mk + n - n_j \geq mk$ . But then each monomial in  $p_j$  occurs at least with ordinary degree  $\frac{mk}{m} = k$  and therefore is contained in  $(I^{GL_n})^k R_{(s)} t^{m+k+n}$ .  $\square$

Let  $\widehat{R^{GL_n}}$  be the  $I^{GL_n}$ -adic completion of the invariant ring  $R^{GL_n}$  and let  $\widehat{S^{GL_n}}$  be the  $J^{GL_n}$ -adic completion of  $S^{GL_n}$ .

LEMMA 4.11. *The morphism  $\psi^*$  induces an isomorphism*

$$R \otimes_{R^{GL_n}} \widehat{R^{GL_n}} \xrightarrow{\simeq} S \otimes_{S^{GL_n}} \widehat{S^{GL_n}}$$

PROOF. Let  $s$  be an irreducible  $GL_n$ -module, then the  $I^{GL_n}$ -adic completion of  $R_{(s)}$  is equal to  $\widehat{R_{(s)}} = R_{(s)} \otimes_{R^{GL_n}} \widehat{R^{GL_n}}$ . Moreover,

$$\widehat{R_{(s)}} = \varprojlim \left( \frac{R}{I^k} \right)_{(s)} = \varprojlim \frac{R_{(s)}}{I^k \cap R_{(s)}}$$

which is the  $I$ -adic completion of  $R_{(s)}$ . By the foregoing lemma both topologies coincide on  $R_{(s)}$  and therefore

$$\widehat{R_{(s)}} = \widehat{R_{(s)}} \quad \text{and similarly} \quad \widehat{S_{(s)}} = \widehat{S_{(s)}}$$

Because  $\widehat{R} \simeq \widehat{S}$  it follows that  $\widehat{R_{(s)}} \simeq \widehat{S_{(s)}}$  from which the result follows as the foregoing holds for all  $s$ .  $\square$

THEOREM 4.12. *Consider a  $GL_n$ -equivariant map  $\underline{Y} \xrightarrow{\psi} \underline{X}$ ,  $y \in \underline{Y}$ ,  $x = \psi(y)$  and  $\psi$  étale in  $y$ . Assume that the orbits  $\mathcal{O}(x)$  and  $\mathcal{O}(y)$  are closed and that  $\psi$  is injective on  $\mathcal{O}(y)$ . Then, there is an affine open subset  $U \hookrightarrow \underline{Y}$  containing  $y$  such that*

- (1)  $U = \pi_Y^{-1}(\pi_Y(U))$  and  $\pi_Y(U) = U/GL_n$ .
- (2)  $\psi$  is étale on  $U$  with affine image.
- (3) The induced morphism  $U/GL_n \xrightarrow{\bar{\psi}} \underline{X}/GL_n$  is étale.
- (4) The diagram below is commutative

$$\begin{array}{ccc} U & \xrightarrow{\psi} & \underline{X} \\ \pi_U \downarrow & & \downarrow \pi_X \\ U/GL_n & \xrightarrow{\bar{\psi}} & \underline{X}/GL_n \end{array}$$

$$\begin{array}{ccccc}
GL_n \times^H N_x & \xleftarrow{GL_n \times^H \phi} & GL_n \times^H \underline{S} & \xrightarrow{\psi} & \underline{X} \\
\downarrow & & \downarrow & & \downarrow \pi \\
N_x/H & \xleftarrow{\phi/H} & \underline{S}/H & \xrightarrow{\psi/GL_n} & \underline{X}/GL_n
\end{array}$$

FIGURE 1. Étale slice diagram

PROOF. By the foregoing lemma we have  $\widehat{R^{GL_n}} \simeq \widehat{S^{GL_n}}$  which means that  $\bar{\psi}$  is étale in  $\pi_Y(y)$ . As étaleness is an open condition, there is an open affine neighborhood  $V$  of  $\pi_Y(y)$  on which  $\bar{\psi}$  is étale. If  $\bar{R} = R \otimes_{R^{GL_n}} S^{GL_n}$  then the above lemma implies that

$$\bar{R} \otimes_{S^{GL_n}} \widehat{S^{GL_n}} \simeq S \otimes_{S^{GL_n}} \widehat{S^{GL_n}}$$

Let  $S_{loc}^{GL_n}$  be the local ring of  $S^{GL_n}$  in  $J^{GL_n}$ , then as the morphism  $S_{loc}^{GL_n} \longrightarrow \widehat{S^{GL_n}}$  is faithfully flat we deduce that

$$\bar{R} \otimes_{S^{GL_n}} S_{loc}^{GL_n} \simeq S \otimes_{S^{GL_n}} S_{loc}^{GL_n}$$

but then there is an  $f \in S^{GL_n} - J^{GL_n}$  such that  $\bar{R}_f \simeq S_f$ . Now, intersect  $V$  with the open affine subset where  $f \neq 0$  and let  $U'$  be the inverse image under  $\pi_Y$  of this set. Remains to prove that the image of  $\psi$  is affine. As  $U' \xrightarrow{\psi} X$  is étale, its image is open and  $GL_n$ -stable. By the separation property we can find an invariant  $h \in R^{GL_n}$  such that  $h$  is zero on the complement of the image and  $h(x) = 1$ . But then we take  $U$  to be the subset of  $U'$  of points  $u$  such that  $h(u) \neq 0$ .  $\square$

**THEOREM 4.13 (Slice theorem).** *Let  $\underline{X}$  be an affine  $GL_n$ -variety with quotient map  $\underline{X} \xrightarrow{\pi} \underline{X}/GL_n$ . Let  $x \in \underline{X}$  be such that its orbit  $\mathcal{O}(x)$  is closed and its stabilizer subgroup  $Stab(x) = H$  is reductive. Then, there is a locally closed affine subscheme  $\underline{S} \hookrightarrow \underline{X}$  containing  $x$  with the following properties*

- (1)  $\underline{S}$  is an affine  $H$ -variety,
- (2) the action map  $GL_n \times \underline{S} \longrightarrow \underline{X}$  induces an étale  $GL_n$ -equivariant morphism  $GL_n \times^H \underline{S} \xrightarrow{\psi} \underline{X}$  with affine image,
- (3) the induced quotient map  $\psi/GL_n$  is étale

$$(GL_n \times^H \underline{S})/GL_n \simeq \underline{S}/H \xrightarrow{\psi/GL_n} \underline{X}/GL_n$$

and the right hand side of figure 1 is commutative.

If we assume moreover that  $\underline{X}$  is smooth in  $x$ , then we can choose the slice  $\underline{S}$  such that also the following properties are satisfied

- (1)  $\underline{S}$  is smooth,
- (2) there is an  $H$ -equivariant morphism  $\underline{S} \xrightarrow{\phi} T_x \underline{S} = N_x$  with  $\phi(x) = 0$  having an affine image,
- (3) the induced morphism is étale

$$\underline{S}/H \xrightarrow{\phi/H} N_x/H$$

and the left hand side of figure 1 is commutative.

PROOF. Choose a finite dimensional  $GL_n$ -subrepresentation  $V$  of  $\mathbb{C}[\underline{X}]$  that generates the coordinate ring as algebra. This gives a  $GL_n$ -equivariant embedding

$$\underline{X} \xrightarrow{i} W = V^*$$

Choose in the vectorspace  $W$  an  $H$ -stable complement  $S_0$  of  $\mathfrak{gl}_n \cdot i(x) = T_{i(x)} \mathcal{O}(x)$  and denote  $S_1 = i(x) + S_0$  and  $\underline{S}_2 = i^{-1}(S_1)$ . Then, the diagram below is commutative

$$\begin{array}{ccc} GL_n \times^H \underline{S}_2 & \xrightarrow{\quad} & GL_n \times^H S_1 \\ \downarrow \psi & & \downarrow \psi_0 \\ \underline{X} & \xrightarrow{i} & W \end{array}$$

By construction we have that  $\psi_0$  induces an isomorphism between the tangent spaces in  $(1, i(x)) \in GL_n \times^H S_0$  and  $i(x) \in W$  which means that  $\psi_0$  is étale in  $i(x)$ , whence  $\psi$  is étale in  $(1, x) \in GL_n \times^H \underline{S}_2$ . By the fundamental lemma we get an affine neighborhood  $U$  which must be of the form  $U = GL_n \times^H \underline{S}$  giving a slice  $\underline{S}$  with the required properties.

Assume that  $\underline{X}$  is smooth in  $x$ , then  $S_1$  is transversal to  $\underline{X}$  in  $i(x)$  as

$$T_{i(x)} i(\underline{X}) + S_0 = W$$

Therefore,  $\underline{S}$  is smooth in  $x$ . Again using the separation property we can find an invariant  $f \in \mathbb{C}[\underline{S}]^H$  such that  $f$  is zero on the singularities of  $\underline{S}$  (which is a  $H$ -stable closed subscheme) and  $f(x) = 1$ . Then replace  $\underline{S}$  with its affine reduced subvariety of points  $s$  such that  $f(s) \neq 0$ . Let  $\mathfrak{m}$  be the maximal ideal of  $\mathbb{C}[\underline{S}]$  in  $x$ , then we have an exact sequence of  $H$ -modules

$$0 \longrightarrow \mathfrak{m}^2 \longrightarrow \mathfrak{m} \xrightarrow{\alpha} N_x^* \longrightarrow 0$$

Choose a  $H$ -equivariant section  $\phi^* : N_x^* \longrightarrow \mathfrak{m} \hookrightarrow \mathbb{C}[\underline{S}]$  of  $\alpha$  then this gives an  $H$ -equivariant morphism  $\underline{S} \xrightarrow{\phi} N_x$  which is étale in  $x$ . Applying again the fundamental lemma to this setting finishes the proof.  $\square$

#### 4.4. Smoothness.

DEFINITION 4.14. Let  $\mathbf{cat}$  be a category of  $\mathbb{C}$ -algebras. An object  $A \in Ob(\mathbf{cat})$  is said to be  $\mathbf{cat}$ -smooth if it satisfies the following lifting property. For  $B \in Ob(\mathbf{cat})$ , a nilpotent ideal  $I \triangleleft B$  such that  $B/I \in Ob(\mathbf{cat})$  and a  $\mathbb{C}$ -algebra morphism  $A \xrightarrow{\kappa} B/I$  in  $Mor(\mathbf{cat})$ , there exist a lifting

$$\begin{array}{ccc} & A & \\ & \swarrow \exists \lambda & \downarrow \kappa \\ B & & B \\ & \longleftarrow & I \end{array}$$

with  $\lambda \in Mor(\mathbf{cat})$  making the diagram commutative.  $\mathbf{alg}$ -smooth is called *Quillen-smooth*,  $\mathbf{comm}$ -smooth is called *Grothendieck-smooth* and  $\mathbf{alg} @_{\mathbf{n}}$ -smooth is called *Cayley-smooth*.

We will show that the categorical notion of **comm**-smoothness coincides with geometric smoothness. Let  $\underline{X}$  be a possibly non-reduced affine variety and  $x$  a geometric point of  $\underline{X}$ . As we are interested in local properties of  $\underline{X}$  near  $x$ , we may assume (after translation) that  $x = o$  in  $\mathbb{C}^n$  and that we have a presentation

$$\mathbb{C}[\underline{X}] = \mathbb{C}[x_1, \dots, x_n]/I \quad \text{with } I = (f_1, \dots, f_m) \text{ and } m_x = (x_1, \dots, x_n)/I.$$

Denote the polynomial algebra  $P = \mathbb{C}[x_1, \dots, x_n]$  and consider the map

$$d : I \longrightarrow (Pdx_1 \oplus \dots \oplus Pdx_n) \otimes_P \mathbb{C}[\underline{X}] = \mathbb{C}[\underline{X}]dx_1 \oplus \dots \oplus \mathbb{C}[\underline{X}]dx_n$$

where the  $dx_i$  are a formal basis of the free module of rank  $n$  and the map is defined by

$$d(f) = \left( \frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_n} \right) \text{ mod } I.$$


This gives a  $\mathbb{C}[\underline{X}]$ -linear mapping  $\frac{I}{I^2} \xrightarrow{d} \mathbb{C}[\underline{X}]dx_1 \oplus \dots \oplus \mathbb{C}[\underline{X}]dx_n$ . Extending to the local algebra  $\mathcal{O}_x$  at  $x$  and then quotient out the maximal ideal  $\mathfrak{m}_x$  we get a  $\mathbb{C} = \mathcal{O}_x/\mathfrak{m}_x$ -linear map  $\frac{\mathfrak{J}}{\mathfrak{J}^2} \xrightarrow{d(x)} \mathbb{C}dx_1 \oplus \dots \oplus \mathbb{C}dx_n$ . Clearly,  $x$  is a nonsingular point of  $\underline{X}$  if and only if the  $\mathbb{C}$ -linear map  $d(x)$  is injective. This is equivalent to the existence of a  $\mathbb{C}$ -section and by the Nakayama lemma also to the existence of a  $\mathcal{O}_x$ -linear splitting  $s_x$  of the induced  $\mathcal{O}_x$ -linear map  $d_x$

$$\frac{\mathfrak{J}}{\mathfrak{J}^2} \begin{array}{c} \xleftarrow{d_x} \\ \xrightarrow{s_x} \end{array} \mathcal{O}_x dx_1 \oplus \dots \oplus \mathcal{O}_x dx_n$$


satisfying  $s_x \circ d_x = id_{\frac{\mathfrak{J}}{\mathfrak{J}^2}}$

A  $\mathbb{C}$ -algebra epimorphism (between commutative algebras)  $R \xrightarrow{\pi} S$  with square zero kernel is called an *infinitesimal extension of  $S$* . It is called a *trivial infinitesimal extension* if  $\pi$  has an algebra section  $\sigma : S \hookrightarrow R$  satisfying  $\pi \circ \sigma = id_S$ .

An infinitesimal extension  $R \xrightarrow{\pi} S$  of  $S$  is said to be *versal* if for any other infinitesimal extension  $R' \xrightarrow{\pi'} S$  of  $S$  there is a  $\mathbb{C}$ -algebra morphism



making the diagram commute. From this universal property it is clear that versal infinitesimal extensions are uniquely determined up to isomorphism. Moreover, if a versal infinitesimal extension is trivial, then so is any infinitesimal extension. By iterating,  $S$  is Grothendieck-smooth if and only if it has the lifting property with respect to nilpotent ideals  $I$  with square zero. Therefore, assume we have a *test object*  $(T, I)$  with  $I^2 = 0$ , then we have a commuting diagram



where we define the *pull-back algebra*  $S \times_{T/I} T = \{(s, t) \in S \times T \mid \kappa(s) = p(t)\}$ . Observe that  $pr_1 : S \times_{T/I} T \rightarrow S$  is a  $\mathbb{C}$ -algebra epimorphism with kernel  $0 \times_{T/I} I$  having square zero, that is, it is an infinitesimal extension of  $S$ . Moreover, the existence of a lifting  $\lambda$  of  $\kappa$  is equivalent to the existence of a  $\mathbb{C}$ -algebra section

$$\sigma : S \longrightarrow S \times_{T/I} T \quad \text{defined by } s \mapsto (s, \lambda(s)).$$

Hence,  $S$  is Grothendieck-smooth if and only if a versal infinitesimal extension of  $S$  is trivial.

Returning to the situation of interest to us, we claim that the algebra epimorphism  $\mathcal{O}_x(\mathbb{C}^n)/I_x^2 \xrightarrow{c_x} \mathcal{O}_x$  is a versal infinitesimal extension of  $\mathcal{O}_x$ . Indeed, consider any other infinitesimal extension  $R \xrightarrow{\pi} \mathcal{O}_x$  then we define a  $\mathbb{C}$ -algebra morphism  $\mathcal{O}_x(\mathbb{C}^n)/I_x^2 \rightarrow R$  as follows: let  $r_i \in R$  such that  $\pi(r_i) = c_x(x_i)$  and define an algebra morphism  $\mathbb{C}[x_1, \dots, x_n] \rightarrow R$  by sending the variable  $x_i$  to  $r_i$ . As the image of any polynomial non-vanishing in  $x$  is a unit in  $R$ , this algebra map extends to one from the local algebra  $\mathcal{O}_x(\mathbb{C}^n)$  and it factors over  $\mathcal{O}_x(\mathbb{C}^n)/I_x^2$  as the image of  $I_x$  lies in the kernel of  $\pi$  which has square zero, proving the claim. Hence,  $\mathcal{O}_x$  is Grothendieck-smooth if and only if there is a  $\mathbb{C}$ -algebra section

$$\mathcal{O}_x(\mathbb{C}^n)/I_x^2 \begin{array}{c} \xrightarrow{c_x} \\ \xleftarrow{r_x} \end{array} \mathcal{O}_x$$

satisfying  $c_x \circ r_x = id_{\mathcal{O}_x}$ .

**PROPOSITION 4.15.** *The affine scheme  $\underline{X}$  is non-singular at the geometric point  $x$  if and only if the local algebra  $\mathcal{O}_x(\underline{X})$  is Grothendieck-smooth.*

**PROOF.** The result will follow once we prove that there is a natural one-to-one correspondence between  $\mathcal{O}_x$ -module splittings  $s_x$  of  $d_x$  and  $\mathbb{C}$ -algebra sections  $r_x$  of  $c_x$ . This correspondence is given by assigning to an algebra section  $r_x$  the map  $s_x$  defined by

$$s_x(dx_i) = (x_i - r_x \circ c_x(x_i)) \text{ mod } I_x^2$$

□

If  $\underline{X}$  is an affine scheme which is smooth in *all* of its geometric points, then we have seen before that  $\underline{X} = X$  must be reduced, that is, an affine variety. Restricting to its disjoint irreducible components we may assume that

$$\mathbb{C}[\underline{X}] = \bigcap_{x \in X} \mathcal{O}_x.$$

Clearly, if  $\mathbb{C}[\underline{X}]$  is Grothendieck-smooth, so is any of the local algebras  $\mathcal{O}_x$ . Conversely, if all  $\mathcal{O}_x$  are Grothendieck-smooth and  $\mathbb{C}[\underline{X}] = \mathbb{C}[x_1, \dots, x_n]/I$  one knows that the algebra epimorphism

$$\mathbb{C}[x_1, \dots, x_n]/I^2 \xrightarrow{c} \mathbb{C}[\underline{X}]$$

has local sections in every  $x$ , but then there is an algebra section. Because  $c$  is clearly a versal infinitesimal deformation of  $\mathbb{C}[\underline{X}]$ , it follows that  $\mathbb{C}[\underline{X}]$  is Grothendieck-smooth.

**PROPOSITION 4.16.** *Let  $\underline{X}$  be an affine scheme. Then,  $\mathbb{C}[\underline{X}]$  is Grothendieck-smooth if and only if  $\underline{X}$  is non-singular in all of its geometric points. In this case,  $\underline{X}$  is a reduced affine variety.*

As  $\text{comm} = \text{alg } @_1$ , we will generalize the foregoing to Cayley-smooth algebras. Let  $B$  be a Cayley-Hamilton algebra of degree  $n$  with trace map  $tr_B$  and trace generated by  $m$  elements say  $\{b_1, \dots, b_m\}$ . Then, we can write

$$B = \mathbb{T}_n^m / T_B \quad \text{with } T_B \text{ closed under traces.}$$

Now, consider the extended ideal

$$E_B = M_n(\mathbb{C}[M_n^m]) \cdot T_B \cdot M_n(\mathbb{C}[M_n^m]) = M_n(N_B)$$

and we have seen that  $\mathbb{C}[\underline{rep}^{tr} B] = \mathbb{C}[M_n^m] / N_B$ . We need the following technical result.

LEMMA 4.17. *With notations as above, we have for all  $k$  that*

$$E_B^{kn^2} \cap \mathbb{T}_n^m \subset T_B^k.$$

PROOF. Let  $\mathbb{T}_n^m$  be the trace algebra on the generic  $n \times n$  matrices  $\{X_1, \dots, X_m\}$  and  $\mathbb{T}_n^{l+m}$  the trace algebra on the generic matrices  $\{Y_1, \dots, Y_l, X_1, \dots, X_m\}$ . Let  $\{U_1, \dots, U_l\}$  be elements of  $\mathbb{T}_n^m$  and consider the trace preserving map  $\mathbb{T}_n^{l+m} \xrightarrow{u} \mathbb{T}_n^m$  induced by the map defined by sending  $Y_i$  to  $U_i$ . Then, by the universal property we have a commutative diagram of Reynold operators

$$\begin{array}{ccc} M_n(\mathbb{C}[M_n^{l+m}]) & \xrightarrow{\tilde{u}} & M_n(\mathbb{C}[M_n^m]) \\ \downarrow R & & \downarrow R \\ \mathbb{T}_n^{l+m} & \xrightarrow{u} & \mathbb{T}_n^m \end{array} .$$

Now, let  $A_1, \dots, A_{l+1}$  be elements from  $M_n(\mathbb{C}[M_n^m])$ , then we can calculate  $R(A_1 U_1 A_2 U_2 A_3 \dots A_l U_l A_{l+1})$  by first computing

$$r = R(A_1 Y_1 A_2 Y_2 A_3 \dots A_l Y_l A_{l+1})$$

and then substituting the  $Y_i$  with  $U_i$ . The Reynolds operator preserves the degree in each of the generic matrices, therefore  $r$  will be linear in each of the  $Y_i$  and is a sum of trace algebra elements. By our knowledge of the generators of necklaces and the trace algebra we can write each term of the sum as an expression

$$tr(M_1)tr(M_2) \dots tr(M_z)M_{z+1}$$

where each of the  $M_i$  is a monomial of degree  $\leq n^2$  in the generic matrices  $\{Y_1, \dots, Y_l, X_1, \dots, X_m\}$ . Now, look at how the generic matrices  $Y_i$  are distributed among the monomials  $M_j$ . Each  $M_j$  contains *at most*  $n^2$  of the  $Y_i$ 's, hence the monomial  $M_{z+1}$  contains *at least*  $l - vn^2$  of the  $Y_i$  where  $v \leq z$  is the number of  $M_i$  with  $i \leq z$  containing at least one  $Y_j$ .

Now, assume all the  $U_i$  are taken from the ideal  $T_B \triangleleft \mathbb{T}_n^m$  which is closed under taking traces, then it follows that

$$R(A_1 U_1 A_2 U_2 A_3 \dots A_l U_l A_{l+1}) \in T_B^{v+(l-vn^2)} \subset T_B^k$$

if we take  $l = kn^2$  as  $v + (k-v)n^2 \geq k$ . But this finishes the proof of the required inclusion.  $\square$



Let  $B$  be a Cayley-Hamilton algebra of degree  $n$  with trace map  $tr_B$  and  $I$  a twosided ideal of  $B$  which is closed under taking traces. We will denote by  $E(I)$  the extended ideal with respect to the universal embedding, that is,

$$E(I) = M_n(\mathbb{C}[\underline{rep}_n^{tr} B])IM_n(\mathbb{C}[\underline{rep}_n^{tr} B]).$$

Then, for all powers  $k$  we have the inclusion  $E(I)^{kn^2} \cap B \subset I^k$ .

**THEOREM 4.18.** *Let  $A$  be a Cayley-Hamilton algebra of degree  $n$  with trace map  $tr_A$ . Then,  $A$  is Cayley-smooth if and only if the trace preserving representation variety  $\underline{rep}_n^{tr} A$  is non-singular in all points (in particular,  $\underline{rep}_n^{tr} A$  is reduced).*

**PROOF.** Let  $A$  be Cayley-smooth, then we have to show that  $\mathbb{C}[\underline{rep}_n^{tr} A]$  is Grothendieck-smooth. Take a commutative test-object  $(T, I)$  with  $I$  nilpotent and an algebra map  $\kappa : \mathbb{C}[\underline{rep}_n^{tr} A] \longrightarrow T/I$ . Composing with the universal embedding  $i_A$  we obtain a trace preserving morphism  $\mu_0$

$$\begin{array}{ccc} A & \xrightarrow{\mu_1} & M_n(T) \\ \downarrow i_A & \searrow \mu_0 & \downarrow \\ M_n(\mathbb{C}[\underline{rep}_n^{tr} A]) & \xrightarrow{M_n(\kappa)} & M_n(T/I) \end{array}$$

Because  $M_n(T)$  with the usual trace is a Cayley-Hamilton algebra of degree  $n$  and  $M_n(I)$  a trace stable ideal and  $A$  is Cayley-smooth there is a trace preserving algebra map  $\mu_1$ . But then, by the universal property of the embedding  $i_A$  there exists a  $\mathbb{C}$ -algebra morphism

$$\lambda : \mathbb{C}[\underline{rep}_n^{tr} A] \longrightarrow T$$

such that  $M_n(\lambda)$  completes the diagram. The morphism  $\lambda$  is the required lift.

Conversely, assume that  $\mathbb{C}[\underline{rep}_n^{tr} A]$  is Grothendieck-smooth. Assume we have a Cayley-Hamilton algebra of degree  $n$  with trace map  $tr_T$  and a trace-stable nilpotent ideal  $I$  of  $T$  and a trace preserving  $\mathbb{C}$ -algebra map  $\kappa : A \longrightarrow T/I$ . If we combine this test-data with the universal embeddings we obtain a diagram

$$\begin{array}{ccccc} & & T \hookrightarrow & i_T & M_n(\mathbb{C}[\underline{rep}_n^{tr} T]) \\ & \nearrow \exists \lambda & \downarrow & & \downarrow \\ A & \xrightarrow{\kappa} & T/I \hookrightarrow & i_{T/I} & M_n(\mathbb{C}[\underline{rep}_n^{tr} T/I]) = M_n(\mathbb{C}[\underline{rep}_n^{tr} T]/J) \\ \downarrow i_A & & \nearrow M_n(\alpha) & & \\ M_n(\mathbb{C}[\underline{rep}_n^{tr} A]) & & & & \end{array}$$


Here,  $J = M_n(\mathbb{C}[\underline{rep}_n^{tr} T])IM_n(\mathbb{C}[\underline{rep}_n^{tr} T])$  and we know already that  $J \cap T = I$ . By the universal property of the embedding  $i_A$  we obtain a  $\mathbb{C}$ -algebra map

$$\mathbb{C}[\underline{rep}_n^{tr} A] \xrightarrow{\alpha} \mathbb{C}[\underline{rep}_n^{tr} T]/J$$

which we would like to lift to  $\mathbb{C}[\underline{rep}_n^{tr} T]$ . This does *not* follow from Grothendieck-smoothness of  $\mathbb{C}[\underline{rep}_n^{tr} A]$  as  $J$  is usually not nilpotent. However, as  $I$  is a nilpotent ideal of  $T$  there is some  $h$  such that  $I^h = 0$ . As  $I$  is closed under taking traces we know by the remark preceding the theorem that

$$E(I)^{hn^2} \cap T \subset I^h = 0.$$

Now, by definition  $E(I) = M_n(\mathbb{C}[\underline{rep}_n^{tr} T])IM_n(\mathbb{C}[\underline{rep}_n^{tr} T])$  which is equal to  $M_n(J)$ . That is, the inclusion can be rephrased as  $M_n(J)^{hn^2} \cap T = 0$ , whence there is a trace preserving embedding  $T \hookrightarrow M_n(\mathbb{C}[\underline{rep}_n^{tr} T]/J^{hn^2})$ . Now, we have the following situation



This time we *can* lift  $\alpha$  to a  $\mathbb{C}$ -algebra morphism

$$\mathbb{C}[\underline{rep}_n^{tr} A] \longrightarrow \mathbb{C}[\underline{rep}_n^{tr} T]/J^{hn^2}.$$

This in turn gives us a trace preserving morphism


$$A \xrightarrow{\lambda} M_n(\mathbb{C}[\underline{rep}_n^{tr} T]/J^{hn^2})$$

the image of which is contained in the algebra of  $GL_n$ -invariants. Because  $T \hookrightarrow M_n(\mathbb{C}[\underline{rep}_n^{tr} T]/J^{hn^2})$  and by surjectivity of invariants under surjective maps, the  $GL_n$ -equivariants are equal to  $T$ , giving the required lift  $\lambda$ .  $\square$

For an affine  $\mathbb{C}$ -algebra  $A$  recall the construction of its *level  $n$  approximation*

$$A@_n = \frac{A^{tr}}{(tr(1) - n, \chi_a^{(n)}(a) \forall a \in A)}$$

In general, it may happen that  $A@_n = 0$  for example if  $A$  has no  $n$ -dimensional representations. The characteristic feature of  $A@_n$  is that any  $\mathbb{C}$ -algebra map  $A \longrightarrow B$  with  $B$  a Cayley-Hamilton algebra of degree  $n$  factors through  $A@_n$



with  $\phi_n$  a trace preserving algebra morphism. From this universal property we deduce

PROPOSITION 4.19. *If  $A$  is Quillen-smooth, then for every integer  $n$ , the Cayley-Hamilton algebra of degree  $n$ ,  $A@_n$ , is Cayley-smooth. Moreover,*

$$\underline{rep}_n A \simeq \underline{rep}_n^{tr} A@_n$$

*is a smooth affine  $GL_n$ -variety.*

This result allows us to study a Quillen-smooth algebra locally in the étale topology. We know that the algebra  $A@_n$  is given by the  $GL_n$ -equivariant maps from  $rep_n A = rep_n^{tr} A@_n$  to  $M_n(\mathbb{C})$ . As this representation variety is smooth we can apply the full strength of the slice theorem to determine the local structure of the  $GL_n$ -variety  $rep_n^{tr} A@_n$  and hence of  $A@_n$ . In the next section we will prove that this local structure is fully determined by a quiver setting. That is, the étale local study of Quillen-smooth algebras can be reduced to that of the subclass of path algebras of quivers.

#### 4.5. Local structure.

In this section we give some applications of the slice theorem to the local structure of quotient varieties of representation spaces. We will first handle the case of an affine  $\mathbb{C}$ -algebra  $A$  leading to a local description of  $A@_n$ . Next, we will refine this slightly to prove similar results for an arbitrary affine  $\mathbb{C}$ -algebra  $B$  in  $\mathfrak{alg} @_n$ .

When  $A$  is an affine  $\mathbb{C}$ -algebra generated by  $m$  elements  $\{a_1, \dots, a_m\}$ , its level  $n$  approximation  $A@_n$  is trace generated by  $m$  determining a trace preserving epimorphism  $\mathbb{T}_n^m \twoheadrightarrow A@_n$ . Thus we have a  $GL_n$ -equivariant closed embedding of affine schemes

$$\underline{rep}_n A = \underline{rep}_n^{tr} A@_n \xrightarrow{\subset^\psi} \underline{rep}_n^{tr} \mathbb{T}_n^m = M_n^m$$

Take a point  $\xi$  of the quotient scheme  $iss_n A = \underline{rep}_n^{tr} A@_n/GL_n$ . We know that  $\xi$  determines the isomorphism class of a semi-simple  $n$ -dimensional representation of  $A$ , say

$$M_\xi = S_1^{\oplus e_1} \oplus \dots \oplus S_k^{\oplus e_k}$$

where the  $S_i$  are distinct simple  $A$ -representations, say of dimension  $d_i$  and occurring in  $M_\xi$  with multiplicity  $e_i$ . These numbers determine the *representation type*  $\tau(\xi)$  of  $\xi$  (or of the semi-simple representation  $M_\xi$ ), that is

$$\tau(\xi) = (e_1, d_1; e_2, d_2; \dots; e_k, d_k)$$

Choosing a basis of  $M_\xi$  adapted to this decomposition gives us a point  $x = (X_1, \dots, X_m)$  in the orbit  $\mathcal{O}(M_\xi)$  such that each  $n \times n$  matrix  $X_i$  is of the form

$$X_i = \begin{bmatrix} m_1^{(i)} \otimes \mathbb{1}_{e_1} & 0 & \dots & 0 \\ 0 & m_2^{(i)} \otimes \mathbb{1}_{e_2} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & m_k^{(i)} \otimes \mathbb{1}_{e_k} \end{bmatrix}$$

where each  $m_j^{(i)} \in M_{d_j}(\mathbb{C})$ . Using this description we can compute the stabilizer subgroup  $Stab(x)$  of  $GL_n$  consisting of those invertible matrices  $g \in GL_n$  commuting with every  $X_i$ . That is,  $Stab(x)$  is the multiplicative group of units of the




FIGURE 2. Big and small normal spaces to the orbit.

centralizer of the algebra generated by the  $X_i$ . It is easy to verify that this group is isomorphic to

$$\text{Stab}(x) \simeq GL_{e_1} \times GL_{e_2} \times \dots \times GL_{e_k} = GL(\alpha_\xi)$$

for the dimension vector  $\alpha_\xi = (e_1, \dots, e_k)$  determined by the multiplicities and with embedding  $\text{Stab}(x) \hookrightarrow GL_n$  given by

$$\begin{bmatrix} GL_{e_1}(\mathbb{C} \otimes \mathbb{1}_{d_1}) & 0 & \dots & 0 \\ 0 & GL_{e_2}(\mathbb{C} \otimes \mathbb{1}_{d_2}) & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & GL_{e_k}(\mathbb{C} \otimes \mathbb{1}_{d_k}) \end{bmatrix}$$

A different choice of point in the orbit  $\mathcal{O}(M_\xi)$  gives a subgroup of  $GL_n$  conjugated to  $\text{Stab}(x)$ .

We know that the normal space  $N_x^{sm}$  can be identified with the self-extensions  $\text{Ext}_A^1(M, M)$  and we will give a quiver-description of this space. The idea is to describe first the  $GL(\alpha)$ -module structure of  $N_x^{big}$ , the normal space to the orbit  $\mathcal{O}(M_\xi)$  in  $M_n^m$  (see figure 2) and then to identify the direct summand  $N_x^{sm}$ . The description of  $N_x^{big}$  follow from a book-keeping operation involving  $GL(\alpha)$ -representations. For  $x = (X_1, \dots, X_m)$ , the tangent space  $T_x \mathcal{O}(M_{x_i})$  in  $M_n^m$  to the orbit is equal to the image of the linear map

$$\begin{array}{ccc} \mathfrak{gl}_n = M_n & \longrightarrow & M_n \oplus \dots \oplus M_n = T_x M_n^m \\ A & \longmapsto & ([A, X_1], \dots, [A, X_m]) \end{array}$$


Observe that the kernel of this map is the centralizer of the subalgebra generated by the  $X_i$ , so we have an exact sequence of  $Stab(x) = GL(\alpha)$ -modules

$$0 \longrightarrow \mathfrak{gl}(\alpha) = Lie GL(\alpha) \longrightarrow \mathfrak{gl}_n = M_n \longrightarrow T_x \mathcal{O}(x) \longrightarrow 0$$

Because  $GL(\alpha)$  is a reductive group every  $GL(\alpha)$ -module is completely reducible and so the sequence splits. But then, the normal space in  $M_n^m = T_x M_n^m$  to the orbit is isomorphic as  $GL(\alpha)$ -module to

$$N_x^{big} = M_n \oplus \underbrace{\dots \oplus M_n}_{m-1} \oplus \mathfrak{gl}(\alpha)$$

with the action of  $GL(\alpha)$  (embedded as above in  $GL_n$ ) is given by simultaneous conjugation. If we consider the  $GL(\alpha)$ -action on  $M_n$



we see that it decomposes into a direct sum of subrepresentations


- for each  $1 \leq i \leq k$  we have  $d_i^2$  copies of the  $GL(\alpha)$ -module  $M_{e_i}$  on which  $GL_{e_i}$  acts by conjugation and the other factors of  $GL(\alpha)$  act trivially,
- for all  $1 \leq i, j \leq k$  we have  $d_i d_j$  copies of the  $GL(\alpha)$ -module  $M_{e_i \times e_j}$  on which  $GL_{e_i} \times GL_{e_j}$  acts via  $g.m = g_i m g_j^{-1}$  and the other factors of  $GL(\alpha)$  act trivially.

These  $GL(\alpha)$  components are precisely the modules appearing in representation spaces of quivers.

**THEOREM 4.20.** *Let  $\xi$  be of representation type  $\tau = (e_1, d_1; \dots; e_k, d_k)$  and let  $\alpha = (e_1, \dots, e_k)$ . Then, the  $GL(\alpha)$ -module structure of the normal space  $N_x^{big}$  in  $M_n^m$  to the orbit of the semi-simple  $n$ -dimensional representation  $\mathcal{O}(M_\xi)$  is isomorphic to*


$$rep_\alpha Q_\xi^{big}$$

where the quiver  $Q_\xi^{big}$  has  $k$  vertices (the number of distinct simple summands of  $M_\xi$ ) and the subquiver on any two vertices  $v_i, v_j$  for  $1 \leq i \neq j \leq k$  has the following shape



That is, in each vertex  $v_i$  there are  $(m-1)d_i^2 + 1$ -loops and there are  $(m-1)d_i d_j$  arrows from vertex  $v_i$  to vertex  $v_j$  for all  $1 \leq i \neq j \leq k$ .

EXAMPLE 4.21. If  $m = 2$  and  $n = 3$  and the representation type is  $\tau = (1, 1; 1, 1; 1, 1)$  (that is,  $M_\xi$  is the direct sum of three distinct one-dimensional simple representations) then the quiver  $Q_\xi$  is



We have  $GL_n$ -equivariant embeddings  $\mathcal{O}(M_\xi) \hookrightarrow \underline{rep}_n^{tr} A @ \mathbf{n} \hookrightarrow M_n^m$  and corresponding embeddings of the tangent spaces in  $x$

$$T_x \mathcal{O}(M_\xi) \hookrightarrow T_x \underline{rep}_n^{tr} A @ \mathbf{n} \hookrightarrow T_x M_n^m$$

Because  $GL(\alpha)$  is reductive we then obtain that the normal spaces to the orbit is a direct summand of  $GL(\alpha)$ -modules.

$$N_x^{sm} = \frac{T_x \underline{rep}_n^{tr} A @ \mathbf{n}}{T_x \mathcal{O}(M_\xi)} \triangleleft N_x^{big} = \frac{T_x M_n^m}{T_x \mathcal{O}(M_\xi)}$$

As we know the isotypical decomposition of  $N_x^{big}$  as the  $GL(\alpha)$ -module  $rep_\alpha Q_\xi$  this allows us to control  $N_x^{sm}$ . We only have to observe that arrows in  $Q_\xi$  correspond to simple  $GL(\alpha)$ -modules, whereas a loop at vertex  $v_i$  decomposes as  $GL(\alpha)$ -module into the simples


$$M_{e_i} = M_{e_i}^0 \oplus \mathbb{C}_{triv}$$

where  $\mathbb{C}_{triv}$  is the one-dimensional simple with trivial  $GL(\alpha)$ -action and  $M_{e_i}^0$  is the space of trace zero matrices in  $M_{e_i}$ . Any  $GL(\alpha)$ -submodule of  $N_x^{big}$  can be represented by a *marked quiver* using the dictionary

- a loop at vertex  $v_i$  corresponds to the  $GL(\alpha)$ -module  $M_{e_i}$  on which  $GL_{e_i}$  acts by conjugation and the other factors act trivially,
- a marked loop at vertex  $v_i$  corresponds to the simple  $GL(\alpha)$ -module  $M_{e_i}^0$  on which  $GL_{e_i}$  acts by conjugation and the other factors act trivially,
- an arrow from vertex  $v_i$  to vertex  $v_j$  corresponds to the simple  $GL(\alpha)$ -module  $M_{e_i \times e_j}$  on which  $GL_{e_i} \times GL_{e_j}$  acts via  $g.m = g_i m g_j^{-1}$  and the other factors act trivially,

Combining this with the calculation that the normalspace is the space of self-extensions  $Ext_A^1(M_\xi, M_\xi)$  or the trace preserving self-extensions  $Ext_B^{tr}(M_\xi, M_{x_i})$  (in case  $B \in Ob(\mathbf{alg} @ \mathbf{n})$ ) we have.

THEOREM 4.22. Consider the marked quiver on  $k$  vertices such that the full marked subquiver on any two vertices  $v_i \neq v_j$  has the form



where these numbers satisfy  $a_{ij} \leq (m-1)d_i d_j$  and  $a_{ii} + m_{ii} \leq (m-1)d_i^2 + 1$ . Then,

$$\begin{array}{ccccc}
GL_n \times^{GL(\alpha)} N_x^{sm} & \xleftarrow{GL_n \times^{GL(\alpha)} \phi} & GL_n \times^{GL(\alpha)} S_x & \xrightarrow{\psi} & \underline{rep}_n A \\
\downarrow & & \downarrow & & \downarrow \\
N_x^{sm}/GL(\alpha) & \xleftarrow{\phi/GL(\alpha)} & S_x/GL(\alpha) & \xrightarrow{\psi/GL(\alpha)} & \underline{iss}_n A
\end{array}$$

FIGURE 3. Slice diagram for representation space.

- (1) Let  $A$  be an affine  $\mathbb{C}$ -algebra generated by  $m$  elements, let  $M_\xi$  be an  $n$ -dimensional semisimple  $A$ -module of representation-type  $\tau = (e_1, d_1; \dots; e_k, d_k)$  and let  $\alpha = (e_1, \dots, e_k)$ . Then, the normal space  $N_x^{sm}$  in a point  $x \in \mathcal{O}(M_\xi)$  to the orbit with respect to the representation space  $\underline{rep}_n A$  is isomorphic to the  $GL(\alpha)$ -module of quiver-representations  $rep_\alpha Q_\xi$  of above type with
- $a_{ii} = \dim_{\mathbb{C}} Ext_A^1(S_i, S_i)$  and  $m_{ii} = 0$  for all  $1 \leq i \leq k$ .
  - $a_{ij} = \dim_{\mathbb{C}} Ext_A^1(S_i, S_j)$  for all  $1 \leq i \neq j \leq n$ .
- (2) Let  $B$  be a Cayley-Hamilton algebra of degree  $n$ , trace generated by  $m$  elements, let  $M_\xi$  be a trace preserving  $n$ -dimensional semisimple  $B$ -module of representation type  $\tau = (e_1, d_1; \dots; e_k, d_k)$  and let  $\alpha = (e_1, \dots, e_k)$ . Then, the normal space  $N_x^{tr}$  in a point  $x \in \mathcal{O}(M_\xi)$  to the orbit with respect to the trace preserving representation space  $\underline{rep}_n^{tr} B$  is isomorphic to the  $GL(\alpha)$ -module of marked quiver-representations  $rep_\alpha Q_\xi^\bullet$  of above type with
- $a_{ij} = \dim_{\mathbb{C}} Ext_B^1(S_i, S_j)$  for all  $1 \leq i \neq j \leq k$ .
- and the (marked) vertex loops further determine the structure of  $Ext_B^{tr}(M_\xi, M_\xi)$ .

By a *marked quiver-representation* we mean a representation of the underlying quiver (that is, forgetting the marks) subject to the condition that the matrices corresponding to marked loops have trace zero.

Consider the slice diagram of figure 3 for the representation space  $\underline{rep}_n A$ . The left hand side exists when  $x$  is a smooth point of  $\underline{rep}_n A$ , the right hand side exists always. The horizontal maps are étale and the upper ones  $GL_n$ -equivariant.

DEFINITION 4.23. A point  $\xi \in \underline{iss}_n A$  is said to belong to the  $n$ -smooth locus of  $A$  iff the representation space  $\underline{rep}_n A$  is smooth in  $x \in \mathcal{O}(M_\xi)$ . The  $n$ -smooth locus of  $A$  will be denoted by  $Sm_n(A)$ .

The quiver necklace algebra  $\mathbb{N}_\alpha Q_\xi$  is the coordinate ring of  $N_x/GL(\alpha)$ . Recall that  $\mathbb{N}_\alpha Q_\xi$  is a graded algebra and is generated by all traces along oriented cycles in the quiver  $Q_\xi$ . Let  $\mathfrak{m}_0$  be the graded maximal ideal of  $\mathbb{N}_\alpha Q_\xi$ , that is corresponding to the closed orbit of the trivial representation. With  $\widehat{\mathbb{T}}_\xi$  (respectively  $\widehat{\mathbb{N}}_\alpha$ ) we will denote the  $\mathfrak{m}_0$ -adic filtration of the quiver-order  $\mathbb{T}_\alpha Q_\xi$  (respectively of the quiver necklace algebra  $\mathbb{N}_\alpha Q_\xi$ ). Recall that the quiver-order  $\mathbb{T}_\alpha Q_\xi$  has a block-decomposition determined by oriented paths in the quiver  $Q_\xi$ . A consequence of the slice theorem and the description of Cayley-Hamilton algebras and their algebra of traces by geometric data we deduce.

THEOREM 4.24. Let  $\xi \in Sm_n(A)$ . Let  $N = tr A \widehat{\mathbb{O}}_{\mathbf{n}}$ , let  $\mathfrak{m}$  be the maximal ideal of  $N$  corresponding to  $\xi$  and denote  $T = A \widehat{\mathbb{O}}_{\mathbf{n}}$ , then we have the isomorphism and Morita equivalence

$$\widehat{N}_{\mathfrak{m}} \simeq \widehat{N}_{\alpha} \quad \text{and} \quad \widehat{T}_{\mathfrak{m}} \underset{\text{Morita}}{\sim} \widehat{T}_{\alpha}$$

We have an explicit description of the algebras on the right in terms of the quiver setting  $(Q_{\xi}, \alpha)$  and the Morita equivalence is determined by the embedding  $GL(\alpha) \hookrightarrow GL_n$ .

Let  $Q^{\bullet}$  be a marked quiver with underlying quiver  $Q$  and let  $\alpha = (d_1, \dots, d_k)$  be a dimension vector. We define the *marked quiver-necklace algebra*  $\mathbb{N}_{\alpha} Q^{\bullet}$  to be the ring of  $GL(\alpha)$ -polynomial invariants on the representation space  $rep_{\alpha} Q^{\bullet}$ , that is,  $\mathbb{N}_{\alpha} Q^{\bullet}$  is the coordinate ring of the quotient variety  $rep_{\alpha} Q^{\bullet}/GL(\alpha)$ . The *marked quiver-order*  $\mathbb{T}_{\alpha} Q^{\bullet}$  is defined to be the algebra of  $GL(\alpha)$ -equivariant polynomial maps from  $rep_{\alpha} Q^{\bullet}$  to  $M_d(\mathbb{C})$  where  $d = \sum_i d_i$ . Because we can separate traces, it follows that

$$\mathbb{N}_{\alpha} Q^{\bullet} = \frac{\mathbb{N}_{\alpha} Q}{(tr(m_1), \dots, tr(m_l))} \quad \text{and} \quad \mathbb{T}_{\alpha} Q^{\bullet} = \frac{\mathbb{T}_{\alpha} Q}{(tr(m_1), \dots, tr(m_l))}$$

where  $\{m_1, \dots, m_l\}$  is the set of all marked loops in  $Q^{\bullet}$ .

Let  $B$  be a Cayley-Hamilton algebra of degree  $n$  and let  $M_{\xi}$  be a trace preserving semi-simple  $B$ -representation of type  $\tau = (e_1, d_1; \dots; e_k, d_k)$  corresponding to the point  $\xi$  in the quotient variety  $iss_n^{tr} B$ .

DEFINITION 4.25. A point  $\xi \in iss_n^{tr} B$  is said to belong to the *smooth locus* of  $B$  iff the trace preserving representation space  $rep_n^{tr} B$  is smooth in  $x \in \mathcal{O}(M_{\xi})$ . The smooth locus of the Cayley-Hamilton algebra  $B$  of degree  $n$  will be denoted by  $Sm_{tr}(B)$ .

Applying the slice theorem to the trace preserving representation space, we obtain with the obvious modifications in notation.

THEOREM 4.26. Let  $\xi \in Sm_{tr}(B)$  and  $N = tr B$ . Let  $\mathfrak{m}$  be the maximal ideal of  $N$  corresponding to  $\xi$ , then we have the isomorphism and Morita equivalence

$$\widehat{N}_{\mathfrak{m}} \simeq \widehat{N}_{\alpha}^{\bullet} \quad \text{and} \quad \widehat{B}_{\mathfrak{m}} \underset{\text{Morita}}{\sim} \widehat{T}_{\alpha}^{\bullet}$$

where we have an explicit description of the algebras on the right in terms of the quiver setting  $(Q_{\xi}, \alpha)$  and where the Morita equivalence is determined by the embedding  $GL(\alpha) \hookrightarrow GL(n)$ .

Even if the left hand sides of the slice diagrams are not defined when  $\xi$  is not contained in the smooth locus, the dimension of the normal spaces (that is, the (trace preserving) self-extensions of  $M_{\xi}$ ) allow us to have a numerical measure of the 'badness' of this noncommutative singularity.

DEFINITION 4.27. Let  $A$  be an affine  $\mathbb{C}$ -algebra and  $\xi \in iss_n A$  of type  $\tau = (e_1, d_1; \dots; e_k, d_k)$ . The *measure of singularity* in  $\xi$  is given by the non-negative number

$$ms(\xi) = n^2 + \dim_{\mathbb{C}} Ext_A^1(M_{\xi}, M_{\xi}) - e_1^2 - \dots - e_k^2 - \dim_{M_{\xi}} rep_n A$$

Let  $B$  be a Cayley-Hamilton algebra of degree  $n$  and  $\xi \in iss_n^{tr} B$  of type  $\tau = (e_1, d_1; \dots; e_k, d_k)$ . The measure of singularity in  $\xi$  is given by the non-negative



number

$$ms(\xi) = n^2 + \dim_{\mathbb{C}} Ext_B^{tr}(M_\xi, M_\xi) - e_1^2 - \dots - e_k^2 - \dim_{M_\xi} \underline{rep}_n^{tr} A$$

Clearly,  $\xi \in Sm_n(A)$  (respectively,  $\xi \in Sm_{tr}(B)$ ) if and only if  $ms(\xi) = 0$ .

As an application to the slice theorem, let us prove the connection between Azumaya algebras and principal fibrations. The *Azumaya locus* of an algebra  $A$  will be the open subset  $U_{Az}$  of  $\underline{iss}_n A$  consisting of the points  $\xi$  of type  $(1, n)$ . Let  $\underline{rep}_n A \xrightarrow{\pi} \underline{iss}_n A$  be the quotient map.

PROPOSITION 4.28. *The quotient  $\pi^{-1}(U_{Az}) \longrightarrow U_{Az}$  is a principal  $PGL_n$ -fibration in the étale topology, that is determines an element in  $H_{\acute{e}t}^1(U_{Az}, PGL_n)$ .*


PROOF. Let  $\xi \in U_{Az}$  and  $x = M_\xi$  a corresponding simple representation. Let  $S_x$  be the slice in  $x$  for the  $PGL_n$ -action on  $\underline{rep}_n A$ . By taking traces of products of a lifted basis from  $M_n(\mathbb{C})$  we find a  $PGL_n$ -affine open neighborhood  $U_\xi$  of  $\xi$  contained in  $U_{Az}$  and hence by the slice result a commuting diagram

$$\begin{array}{ccc} PGL_n \times S_x & \xrightarrow{\psi} & \pi^{-1}(U_\xi) \\ \downarrow & & \downarrow \pi \\ S_x & \xrightarrow{\psi/PGL_n} & U_\xi \end{array}$$

where  $\psi$  and  $\psi/PGL_n$  are étale maps. That is,  $\psi/PGL_n$  is an étale neighborhood of  $\xi$  over which  $\pi$  is trivialized. As this holds for all points  $\xi \in U_{Az}$  the result follows.  $\square$

#### 4.6. $A_\infty$ -algebras.

The category  $\mathbf{alg}$  has a topological origin. Consider the *tiny interval operad*  $D_1$ , that is, let  $D_1(n)$  be the collection of all configurations



consisting of the unit interval with  $n$  closed intervals removed, each gap given a label  $i_j$  where  $(i_1, i_2, \dots, i_n)$  is a permutation of  $(1, 2, \dots, n)$ . Clearly,  $D_1(n)$  is a real  $2n$ -dimensional  $\mathcal{C}^\infty$ -manifold having  $n!$  connected components, each of which is a contractible space. The operad structure comes from the collection of composition maps

$$D_1(n) \times (D_1(m_1) \times \dots \times D_1(m_n)) \xrightarrow{m(n, m_1, \dots, m_n)} D_1(m_1 + \dots + m_n)$$

defined by resizing the configuration in the  $D_1(m_i)$ -component such that it fits precisely in the  $i$ -th gap of the configuration of the  $D_1(n)$ -component, see figure 4. We obtain a unit interval having  $m_1 + \dots + m_n$  gaps which are labeled in the natural way, that is the first  $m_1$  labels are for the gaps in the  $D_1(m_1)$ -configuration fitted in gap 1, the next  $m_2$  labels are for the gaps in the  $D_1(m_2)$ -configuration fitted in gap 2 and so on. The tiny interval operad  $D_1$  consists of

- a collection of topological spaces  $D_1(n)$  for  $n \geq 0$ ,




FIGURE 4. The tiny interval operad.

- a continuous action of  $S_n$  on  $D_1(n)$  by relabeling, for every  $n$ ,
- an identity element  $id \in D_1(1)$ ,
- the continuous composition maps  $m_{(n,m_1,\dots,m_n)}$  which satisfy associativity and equivariance with respect to the symmetric group actions.

By taking the *homology groups* of these manifolds  $D_1(n)$  we obtain a *linear operad*  $\mathbf{ASSOC}$ . Because  $D_1(n)$  has  $n!$  contractible components we can identify  $\mathbf{ASSOC}(n)$  with the subspace of the free algebra  $\mathbb{C}\langle x_1, \dots, x_n \rangle$  spanned by the *multilinear monomials*.  $\mathbf{ASSOC}(n)$  has dimension  $n!$  with basis  $x_{\sigma(1)} \dots x_{\sigma(n)}$  for  $\sigma \in S_n$ . Each  $\mathbf{ASSOC}(n)$  has a natural action of  $S_n$  and as  $S_n$ -representation it is isomorphic to the regular representation. The composition maps  $m_{(n,m_1,\dots,m_n)}$  induce on the homology level linear composition maps

$$\mathbf{ASSOC}(n) \otimes \mathbf{ASSOC}(m_1) \otimes \dots \otimes \mathbf{ASSOC}(m_n) \xrightarrow{\gamma_{(n,m_1,\dots,m_n)}} \mathbf{ASSOC}(m_1 + \dots + m_n)$$

obtained by substituting the multilinear monomials  $\phi_i \in \mathbf{ASSOC}(m_i)$  in the place of the variable  $x_i$  into the multilinear monomial  $\psi \in \mathbf{ASSOC}(n)$ .

In general, a  $\mathbb{C}$ -linear operad  $\mathbf{P}$  consists of a family of vectorspaces  $\mathbf{P}(n)$  each equipped with an  $S_n$ -action,  $\mathbf{P}(1)$  contains an identity element and there are composition linear morphisms

$$\mathbf{P}(n) \otimes \mathbf{P}(m_1) \otimes \dots \otimes \mathbf{P}(m_n) \xrightarrow{c_{(n,m_1,\dots,m_n)}} \mathbf{P}(m_1 + \dots + m_n)$$

satisfying the same compatibility relations as the maps  $\gamma_{(n,m_1,\dots,m_n)}$  above. An example is the *endomorphism operad*  $\mathbf{end}_V$  for a vectorspace  $V$  defined by taking

$$\mathbf{end}_V(n) = \mathit{Hom}_{\mathbb{C}}(V^{\otimes n}, V)$$

with compositions and  $S_n$ -action defined in the obvious way and unit element  $\mathbb{1}_V \in \mathbf{end}_V(1) = \mathit{End}(V)$ . A *morphism* of linear operads  $\mathbf{P} \xrightarrow{f} \mathbf{P}'$  is a collection of

linear maps which are equivariant with respect to the  $S_n$ -action, commute with the composition maps and take the identity element of  $\mathbb{P}$  to the identity element of  $\mathbb{P}'$ .

DEFINITION 4.29. Let  $\mathbb{P}$  be a  $\mathbb{C}$ -linear operad. A  $\mathbb{P}$ -algebra is a vectorspace  $A$  equipped with a morphism of operads  $\mathbb{P} \xrightarrow{f} \mathbf{end}_A$ .

For example, **assoc**-algebras are just associative  $\mathbb{C}$ -algebras, explaining the topological origin of **alg**. Instead of considering the *homology operad* **assoc** of the tiny intervals  $D_1$  we can consider its *chain operad* **chain**. For a topological space  $X$ , let **chains**( $X$ ) be the complex concentrated in non-positive degrees, whose  $-k$ -component consists of the finite formal additive combinations  $\sum c_i \cdot f_i$  where  $c_i \in \mathbb{C}$  and  $f_i : [0, 1]^k \longrightarrow X$  is a continuous map (a *singular cube* in  $X$ ) modulo the following relations

- For any  $\sigma \in S_k$  acting on  $[0, 1]^k$  by permutation, we have  $f \circ \sigma = sg(\sigma)f$ .
- For  $pr_{k-1}^k : [0, 1]^k \xrightarrow{k-1}$  the projection on the first  $k-1$  coordinates and any continuous map  $[0, 1]^{k-1} \xrightarrow{f'} X$  we have  $f' \circ pr_{k-1}^k = 0$ .

Then, **chain** is the collection of complexes **chains**( $D_1(n)$ ) and is an operad in the category of complexes of vectorspaces with cohomology the homology operad **assoc**. Again, we can consider **chain**-algebras, this time as complexes of vectorspaces. These are the  $A_\infty$ -algebras.

DEFINITION 4.30. An  $A_\infty$ -algebra is a  $\mathbb{Z}$ -graded complex vectorspace

$$B = \bigoplus_{p \in \mathbb{Z}} B_p$$

endowed with homogeneous  $\mathbb{C}$ -linear maps

$$m_n : B^{\otimes n} \longrightarrow B$$

of degree  $2-n$  for all  $n \geq 1$ , satisfying the following relations

- We have  $m_1 \circ m_1 = 0$ , that is  $(B, m_1)$  is a differential complex

$$\dots \xrightarrow{m_1} B_{i-1} \xrightarrow{m_1} B_i \xrightarrow{m_1} B_{i+1} \xrightarrow{m_1} \dots$$

- We have the equality of maps  $B \otimes B \longrightarrow B$

$$m_1 \circ m_2 = m_2 \circ (m_1 \otimes \mathbb{1} + \mathbb{1} \otimes m_1)$$

where  $\mathbb{1}$  is the identity map on the vectorspace  $B$ . That is,  $m_1$  is a derivation with respect to the multiplication  $B \otimes B \xrightarrow{m_2} B$ .

- We have the equality of maps  $B \otimes B \otimes B \longrightarrow B$

$$\begin{aligned} & m_2 \circ (\mathbb{1} \otimes m_2 - m_2 \otimes \mathbb{1}) \\ &= m_1 \circ m_3 + m_3 \circ (m_1 \otimes \mathbb{1} \otimes \mathbb{1} + \mathbb{1} \otimes m_1 \otimes 1 + \mathbb{1} \otimes \mathbb{1} \otimes m_1) \end{aligned}$$

where the right second expression is the associator for the multiplication  $m_2$  and the first is a boundary of  $m_3$ , implying that  $m_2$  is associative up to homology.

- More generally, for  $n \geq 1$  we have the relations

$$\sum (-1)^{i+j+k} m_l \circ (\mathbb{1}^{\otimes i} \otimes m_j \otimes \mathbb{1}^{\otimes k}) = 0$$

where the sum runs over all decompositions  $n = i + j + k$  and where  $l = i + 1 + k$ . These identities are pictorially represented in figure 5.




FIGURE 5.  $A_\infty$ -identities.

Observe that an  $A_\infty$ -algebra  $B$  is in general not associative for the multiplication  $m_2$ , but its homology

$$H^* B = H^*(B, m_2)$$

is an associative graded algebra for the multiplication induced by  $m_2$ . Further, if  $m_n = 0$  for all  $n \geq 3$ , then  $B$  is an associative *differentially graded algebra* and conversely every differentially graded algebra yields an  $A_\infty$ -algebra with  $m_n = 0$  for all  $n \geq 3$ .

Let  $A$  be an associative  $\mathbb{C}$ -algebra and  $M$  a left  $A$ -module. Choose an *injective resolution* of  $M$

$$0 \longrightarrow M \longrightarrow I^0 \longrightarrow I^1 \longrightarrow \dots$$

with the  $I^k$  injective left  $A$ -modules and denote by  $I^\bullet$  the complex

$$I^\bullet : 0 \longrightarrow I^0 \xrightarrow{d} I^1 \xrightarrow{d} \dots$$

Let  $B = \text{HOM}_A^\bullet(I^\bullet, I^\bullet)$  be the morphism complex. That is, its  $n$ -th component are the graded  $A$ -linear maps  $I^\bullet \longrightarrow I^\bullet$  of degree  $n$ . This space can be equipped with a differential

$$d(f) = d \circ f - (-1)^n f \circ d \quad \text{for } f \text{ in the } n\text{-th part}$$

Then,  $B$  is a differentially graded algebra where the multiplication is the natural composition of graded maps. The homology algebra

$$H^* B = \text{Ext}_A^*(M, M)$$

is the *extension algebra* of  $M$ . Generalizing the description of  $\text{Ext}_A^1(M, M)$  given in section 3.3, an element of  $\text{Ext}_A^k(M, M)$  is an equivalence class of exact sequences of  $A$ -modules

$$0 \longrightarrow M \longrightarrow P_1 \longrightarrow P_2 \longrightarrow \dots \longrightarrow P_k \longrightarrow M \longrightarrow 0$$

and the algebra structure on the extension algebra is induced by concatenation of such sequences. This extension algebra has a *canonical* structure of  $A_\infty$ -algebra with  $m_1 = 0$  and  $m_2$  the usual multiplication.

Now, let  $M_1, \dots, M_k$  be  $A$ -modules (for example, finite dimensional representations) and with  $\text{filt}(M_1, \dots, M_k)$  we denote the full subcategory of all  $A$ -modules

whose objects admit finite filtrations with subquotients among the  $M_i$ . We have the following result, for a proof and more details we refer to the excellent notes by B. Keller [35, §6].

**THEOREM 4.31.** *Let  $M = M_1 \oplus \dots \oplus M_k$ . The canonical  $A_\infty$ -structure on the extension algebra  $Ext_A^*(M, M)$  contains enough information to reconstruct the category  $filt(M_1, \dots, M_k)$ .*

If we specialize to the case when  $M$  is a semi-simple  $n$ -dimensional representation of  $A$  of representation type  $\tau = (e_1, d_1; \dots; e_k, d_k)$  say with decomposition

$$M_\xi = S_1^{\oplus e_1} \oplus \dots \oplus S_k^{\oplus e_k}$$

Then, the first two terms of the extension algebra  $Ext_A^*(M_\xi, M_\xi)$  are

- $Ext_A^0(M_\xi, M_\xi) = End_A(M_\xi) = M_{e_1}(\mathbb{C}) \oplus \dots \oplus M_{e_k}(\mathbb{C})$  because by *Schur's lemma*  $Hom_A(S_i, S_j) = \delta_{ij}\mathbb{C}$ . Hence, the 0-th part of  $Ext_A^*(M_\xi, M_\xi)$  determine the dimension vector  $\alpha = (e_1, \dots, e_k)$ .
- $Ext_A^1(M_\xi, M_\xi) = \bigoplus_{i,j=1}^k M_{e_j \times e_i}(Ext_A^1(S_i, S_j))$  and we have seen that  $dim_{\mathbb{C}} Ext_A^1(S_i, S_j)$  is the number of arrows from vertex  $v_i$  to  $v_j$  in the local quiver  $Q_\xi$ .

Summarizing the results of the previous section, we have :

**PROPOSITION 4.32.** *Let  $\xi \in Sm_n(A)$ , then the first two terms of the extension algebra  $Ext_A^*(M_\xi, M_\xi)$  contain enough information to determine the étale local structure of  $\underline{rep}_n A$  and  $\underline{iss}_n A$  near  $M_\xi$ .*

If one wants to extend this result to noncommutative singular points  $\xi \notin Sm_n(A)$ , one will have to consider the canonical  $A_\infty$ -structure on the full extension algebra  $Ext_A^*(M_\xi, M_\xi)$ .

#### 4.7. Indecomposable roots.

Throughout,  $Q$  will be a quiver on  $k$  vertices  $\{v_1, \dots, v_k\}$  with Euler form  $\chi_Q$ . For a dimension vector  $\alpha = (d_1, \dots, d_k)$ , any  $V \in rep_\alpha Q$  decomposes uniquely into

$$V = W_1^{\oplus f_1} \oplus \dots \oplus W_z^{\oplus f_z}$$

where the  $W_i$  are *indecomposable* representations . This follows from the fact that  $End(V)$  is finite dimensional. Recall also that a representation  $W$  of  $Q$  is indecomposable if and only if  $End(W)$  is a *local algebra* , that is, the nilpotent endomorphisms in  $End_{\mathbb{C}Q}(W)$  form an ideal of codimension one. Equivalently, the maximal torus of the stabilizer subgroup  $Stab_{GL(\alpha)}(W) = Aut_{\mathbb{C}Q}(W)$  is one-dimensional, which means that every semisimple element of  $Aut_{\mathbb{C}Q}(W)$  lies in  $\mathbb{C}^*(\mathbb{1}_{d_1}, \dots, \mathbb{1}_{d_k})$ . More generally, decomposing a representation  $V$  into indecomposables corresponds to choosing a maximal torus in the stabilizer subgroup  $Aut_{\mathbb{C}Q}(V)$ . Let  $T$  be such a maximal torus, we define a decomposition of the vertexspaces

$$V_i = \bigoplus_\chi V_i(\chi) \quad \text{where} \quad V_i(\chi) = \{v \in V_i \mid t.v = \chi(t)v \ \forall t \in T\}$$

where  $\chi$  runs over all characters of  $T$ . One verifies that each  $V(\chi) = \bigoplus_i V_i(\chi)$  is a subrepresentation of  $V$  giving a decomposition  $V = \bigoplus_\chi V(\chi)$ . Because  $T$  acts by scalar multiplication on each component  $V(\chi)$ , we have that  $\mathbb{C}^*$  is the maximal torus of  $Aut_{\mathbb{C}Q}(V(\chi))$ , whence  $V(\chi)$  is indecomposable. Conversely, if  $V = W_1 \oplus \dots \oplus W_r$  is a decomposition with the  $W_i$  indecomposable, then the product of all the one-dimensional maximal tori in  $Aut_{\mathbb{C}Q}(W_i)$  is a maximal torus of  $Aut_{\mathbb{C}Q}(V)$ .

In this section we will give a classification of the *indecomposable roots*, that is, the dimension vectors of indecomposable representations. As the name suggests, these dimension vectors will form a *root system*.

The *Tits form* of a quiver  $Q$  is the symmetrization of its Euler form, that is,

$$T_Q(\alpha, \beta) = \chi_Q(\alpha, \beta) + \chi_Q(\beta, \alpha)$$

This symmetric bilinear form is described by the *Cartan matrix*

$$C_Q = \begin{bmatrix} c_{11} & \dots & c_{1k} \\ \vdots & & \vdots \\ c_{k1} & \dots & c_{kk} \end{bmatrix} \quad \text{with } c_{ij} = 2\delta_{ij} - \# \{ \textcircled{i} \text{---} \textcircled{j} \}$$

where we count all arrows connecting  $v_i$  with  $v_j$  forgetting the orientation. The corresponding quadratic form  $q_Q(\alpha) = \frac{1}{2}\chi_Q(\alpha, \alpha)$  on  $\mathbb{Q}^k$  is defined to be

$$q_Q(x_1, \dots, x_k) = \sum_{i=1}^k x_i^2 - \sum_{a \in Q_a} x_{t(a)}x_{h(a)}$$

Hence,  $q_Q(\alpha) = \dim GL(\alpha) - \dim \text{rep}_\alpha Q$ . With  $\Gamma_Q$  we denote the underlying graph of  $Q$ , that is, forgetting the orientation of the arrows. The following classification result is classical, see for example [9]. A quadratic form  $q$  on  $\mathbb{Z}^k$  is said to be *positive definite* if  $0 \neq \alpha \in \mathbb{Z}^k$  implies  $q(\alpha) > 0$ . It is called *positive semi-definite* if  $q(\alpha) \geq 0$  for all  $\alpha \in \mathbb{Z}^k$ . The *radical* of  $q$  is  $\text{rad}(q) = \{ \alpha \in \mathbb{Z}^k \mid T(\alpha, -) = 0 \}$ . Recall that when  $Q$  is a connected and  $\alpha \geq \underline{0}$  is a non-zero radical vector, then  $\alpha$  is *sincere* (that is, all components of  $\alpha$  are non-zero) and  $q_Q$  is positive semi-definite. There exist a minimal  $\delta_Q \geq \underline{0}$  with the property that  $q_Q(\alpha) = 0$  if and only if  $\alpha \in \mathbb{Q}\delta_Q$  if and only if  $\alpha \in \text{rad}(q_Q)$ . If the quadratic form  $q$  is neither positive definite nor semi-definite, it is called *indefinite*.

**THEOREM 4.33.** *Let  $Q$  be a connected quiver with Tits form  $q_Q$ , Cartan matrix  $C_Q$  and underlying graph  $\Gamma_Q$ . Then,*

- (1)  $q_Q$  is positive definite if and only if  $\Gamma_Q$  is a Dynkin diagram, that is one of the graphs of figure 6. The number of vertices is  $m$ .
- (2)  $q_Q$  is semidefinite if and only if  $\Gamma_Q$  is an extended Dynkin diagram, that is one of the graphs of figure 7 and  $\delta_Q$  is the indicated dimension vector. The number of vertices is  $m + 1$ .

Let  $V \in \text{rep}_\alpha Q$  be decomposed into indecomposables

$$V = W_1^{\oplus f_1} \oplus \dots \oplus W_z^{\oplus f_z}$$

If  $\dim(W_i) = \gamma_i$  we say that  $V$  is of type  $(f_1, \gamma_1; \dots; f_z, \gamma_z)$ .

**PROPOSITION 4.34.** *For any dimension vector  $\alpha$ , there exists a unique type  $\tau_{\text{can}} = (e_1, \beta_1; \dots; e_l, \beta_l)$  with  $\alpha = \sum_i e_i \beta_i$  such that the set  $\text{rep}_\alpha(\tau_{\text{can}}) =$*

$\{ V \in \text{rep}_\alpha Q \mid V \simeq W_1^{\oplus e_1} \oplus \dots \oplus W_l^{\oplus e_l}, \dim(W_i) = \beta_i, W_i \text{ is indecomposable} \}$  contains a dense open set of  $\text{rep}_\alpha Q$ .

**PROOF.** Recall from example 2.7 that for any dimension vector  $\beta$  the subset  $\text{rep}_\beta^{\text{ind}} Q$  of *indecomposable* representations of dimension  $\beta$  is constructible. Consider for a type  $\tau = (f_1, \gamma_1; \dots; f_z, \gamma_z)$  the subset  $\text{rep}_\alpha(\tau) =$

$$\{ V \in \text{rep}_\alpha Q \mid V \simeq W_1^{\oplus f_1} \oplus \dots \oplus W_z^{\oplus f_z}, \dim(W_i) = \gamma_i, W_i \text{ indecomposable} \}$$




FIGURE 6. The Dynkin diagrams.




FIGURE 7. The extended Dynkin diagrams.

then  $rep_\alpha(\tau)$  is a constructible subset of  $rep_\alpha Q$  as it is the image of the constructible set

$$GL(\alpha) \times rep_{\gamma_1}^{ind} Q \times \dots \times rep_{\gamma_z}^{ind} Q$$

under the map sending  $(g, W_1, \dots, W_z)$  to  $g \cdot (W_1^{\oplus f_1} \oplus \dots \oplus W_z^{\oplus f_z})$ . Because of the uniqueness of the decomposition into indecomposables we have a finite disjoint decomposition

$$rep_\alpha Q = \bigsqcup_{\tau} rep_\alpha(\tau)$$

and by irreducibility of  $rep_\alpha Q$  precisely one of the  $rep_\alpha(\tau)$  contains a dense open set of  $rep_\alpha Q$ .  $\square$

We call  $\tau_{can}$  the *canonical decomposition* of  $\alpha$ . In the next section we will give an algorithm to compute the canonical decomposition. Consider the action morphisms  $GL(\alpha) \times rep_\alpha Q \xrightarrow{\phi} rep_\alpha Q$ . By Chevalley's theorem 2.4 we know that the function

$$V \mapsto \dim Stab_{GL(\alpha)}(V)$$

is upper semi-continuous. Because  $\dim GL(\alpha) = \dim Stab_{GL(\alpha)}(V) + \dim \mathcal{O}(V)$  we conclude that for all  $m$ , the subset

$$rep_\alpha(m) = \{V \in rep_\alpha Q \mid \dim \mathcal{O}(V) \geq m\}$$

is Zariski open. In particular,  $rep_\alpha(max)$  the union of all orbits of maximal dimension is open and dense in  $rep_\alpha Q$ . A representation  $V \in rep_\alpha Q$  lying in the intersection

$$rep_\alpha(\tau_{can}) \cap rep_\alpha(max)$$

is called a *generic representation* of dimension  $\alpha$ .

Assume that  $Q$  is a connected quiver of *finite representation type*, that is, there are only a finite number of isomorphism classes of indecomposable representations. Let  $\alpha$  be an arbitrary dimension vector. Since any representation of  $Q$  can be decomposed into a direct sum of indecomposables,  $rep_\alpha Q$  contains only finitely many orbits. Hence, one orbit  $\mathcal{O}(V)$  must be dense and have the same dimension as  $rep_\alpha Q$ , but then

$$\dim rep_\alpha Q = \dim \mathcal{O}(V) \leq \dim GL(\alpha) - 1$$

as any representation has  $\mathbb{C}^*(\mathbb{1}_{a_1}, \dots, \mathbb{1}_{a_k})$  in its stabilizer subgroup. That is, for every  $\alpha \in \mathbb{N}^k$  we have  $q_Q(\alpha) \geq 1$ . Because all off-diagonal entries of the Cartan matrix  $C_Q$  are non-positive, it follows that  $q_Q$  is positive definite on  $\mathbb{Z}^k$  whence  $\Gamma_Q$  must be a Dynkin diagram. It is well known that to a Dynkin diagram one associates a simple Lie algebra and a corresponding *root system*. We will generalize the notion of a root system to an arbitrary quiver  $Q$ .

Let  $\epsilon_i = (\delta_{1i}, \dots, \delta_{ki})$  be the standard basis of  $\mathbb{Q}^k$ . The *fundamental set of roots* is defined to be the following set of dimension vectors

$$F_Q = \{\alpha \in \mathbb{N}^k - \underline{0} \mid T_Q(\alpha, \epsilon_i) \leq 0 \text{ and } \text{supp}(\alpha) \text{ is connected}\}$$

Recall that it follows from the description of dimension vectors of simple representations given in section 3.4 that any simple root lies in the fundamental set.

**LEMMA 4.35.** *Let  $\alpha = \beta_1 + \dots + \beta_s \in F_Q$  with  $\beta_i \in \mathbb{N}^k - \underline{0}$  for  $1 \leq i \leq s \leq 2$ . If  $q_Q(\alpha) \geq q_Q(\beta_1) + \dots + q_Q(\beta_s)$ , then  $\text{supp}(\alpha)$  is a tame quiver (that is, its underlying graph is an extended Dynkin diagram) and  $\alpha \in \mathbb{N}\delta_{\text{supp}(\alpha)}$ .*

**PROOF.** Let  $s = 2$ ,  $\beta_1 = (c_1, \dots, c_k)$  and  $\beta_2 = (d_1, \dots, d_k)$  and we may assume that  $\text{supp}(\alpha) = Q$ . By assumption  $T_Q(\beta_1, \beta_2) = q_Q(\alpha) - q_Q(\beta_1) - q_Q(\beta_2) \geq 0$ . Using that  $C_Q$  is symmetric and  $\alpha = \beta_1 + \beta_2$  we have

$$\begin{aligned} 0 \leq T_Q(\beta_1, \beta_2) &= \sum_{i,j} c_{ij} c_i d_j \\ &= \sum_j \frac{c_j d_j}{a_j} \sum_i c_{ij} a_i + \frac{1}{2} \sum_{i \neq j} c_{ij} \left( \frac{c_i}{a_i} - \frac{c_j}{a_j} \right)^2 a_i a_j \end{aligned}$$



and because  $T_Q(\alpha, \epsilon_i) \leq 0$  and  $c_{ij} \leq 0$  for all  $i \neq j$ , we deduce that

$$\frac{c_i}{a_i} = \frac{c_j}{a_j} \quad \text{for all } i \neq j \text{ such that } c_{ij} \neq 0$$

Because  $Q$  is connected,  $\alpha$  and  $\beta_1$  are proportional. But then,  $T_Q(\alpha, \epsilon_i) = 0$  and hence  $C_Q\alpha = \underline{0}$ . By the classification result,  $q_Q$  is semidefinite whence  $\Gamma_Q$  is an extended Dynkin diagram and  $\alpha \in \mathbb{N}\delta_Q$ . Finally, if  $s > 2$ , then

$$T_Q(\alpha, \alpha) = \sum_i T_Q(\alpha, \beta_i) \geq \sum_i T_Q(\beta_i, \beta_i)$$

whence  $T_Q(\alpha - \beta_i, \beta_i) \geq 0$  for some  $i$  and then we can apply the foregoing argument to  $\beta_i$  and  $\alpha - \beta_i$ .  $\square$

**DEFINITION 4.36.** If  $G$  is an algebraic group acting on a variety  $Y$  and if  $X \hookrightarrow Y$  is a  $G$ -stable subset, then we can decompose  $X = \bigcup_d X_{(d)}$  where  $X_{(d)}$  is the union of all orbits  $\mathcal{O}(x)$  of dimension  $d$ . The *number of parameters* of  $X$  is

$$\mu(X) = \max_d (\dim X_{(d)} - d)$$

where  $\dim X_{(d)}$  denotes the dimension of the Zariski closure of  $X_{(d)}$ .

In the special case of  $GL(\alpha)$  acting on  $\text{rep}_\alpha Q$ , we denote  $\mu(\text{rep}_\alpha(\text{max})) = p_Q(\alpha)$  and call it the *number of parameters* of  $\alpha$ . For example, if  $\alpha$  is a Schur root, then  $p(\alpha) = \dim \text{rep}_\alpha Q - (\dim GL(\alpha) - 1) = 1 - q_Q(\alpha)$ .

Recall that a matrix  $m \in M_n(\mathbb{C})$  is *unipotent* if some power  $m^k = \mathbb{1}_n$ . It follows from the Jordan normal form that  $GL(\alpha)$  and  $PGL(\alpha) = GL(\alpha)/\mathbb{C}^*$  contain only finitely many conjugacy classes of unipotent matrices.

**THEOREM 4.37.** *If  $\alpha$  lies in the fundamental set and  $\text{supp}(\alpha)$  is not tame, then*

$$p_Q(\alpha) = \mu(\text{rep}_\alpha(\text{max})) = \mu(\text{rep}_\alpha^{\text{ind}} Q) = 1 - q_Q(\alpha) > \mu(\text{rep}_\alpha^{\text{ind}}(d))$$

for all  $d > 1$  where  $\text{rep}_\alpha^{\text{ind}}(d)$  is the union of all indecomposable orbits of dimension  $d$ .

**PROOF.** A representation  $V \in \text{rep}_\alpha Q$  is indecomposable if and only if its stabilizer subgroup  $\text{Stab}_{GL(\alpha)}(V)$  is a *unipotent group*, that is all its elements are unipotent elements. By proposition 4.47 we know that  $\text{rep}_\alpha(\text{max}) \hookrightarrow \text{rep}_\alpha^{\text{ind}} Q$  and that  $p_Q(\alpha) = \mu(\text{rep}_\alpha(\text{max})) = 1 - q_Q(\alpha)$ . Denote  $\text{rep}_\alpha(\text{sub}) = \text{rep}_\alpha Q - \text{rep}_\alpha(\text{max})$ . We claim that for any unipotent element  $u \neq \mathbb{1}$  we have that

$$\dim \text{rep}_\alpha(\text{sub})(u) - \dim \text{cen}_{GL(\alpha)}(u) + 1 < 1 - q_Q(\alpha)$$

where  $\text{rep}_\alpha(\text{sub})(g)$  denotes the representations in  $\text{rep}_\alpha(\text{sub})$  having  $g$  in their stabilizer subgroup. In fact, for any  $g \in GL(\alpha) - \mathbb{C}^*$  we have

$$\dim \text{cen}_{GL(\alpha)}(g) - \dim \text{rep}_\alpha(g) > q_Q(\alpha)$$

Indeed, we may reduce to  $g$  being a semisimple element, see [47, lemma 3.4]. then, if  $\alpha = \alpha_1 + \dots + \alpha_s$  is the decomposition of  $\alpha$  obtained from the eigenspace decompositions of  $g$  (we have  $s \geq 2$  as  $g \notin \mathbb{C}^*$ ), then

$$\text{cen}_{GL(\alpha)}(g) = \prod_i GL(\alpha_i) \quad \text{and} \quad \text{rep}_\alpha(g) = \prod_i \text{rep}_{\alpha_i}(g)$$

whence  $\dim \text{cen}_{PGL(\alpha)}(g) - \dim \text{rep}_\alpha(g) = \sum_i q_Q(\alpha_i) > q_Q(\alpha)$ , proving the claim. Further, we claim that

$$\mu(\text{rep}_\alpha(\text{sub})) \leq \max_u (\dim \text{rep}_\alpha(\text{sub})(u) - \dim \text{cen}_{PGL(\alpha)}(u) + 1)$$

Let  $Z = \text{rep}_\alpha(\text{sub})$  and consider the closed subvariety of  $PGL(\alpha) \times Z$

$$L = \{(g, z) \mid g.z = z\}$$

For  $z \in Z$  we have  $pr_1^{-1}(z) = \text{Stab}_{PGL(\alpha)}(z) \times \{z\}$  and if  $z$  is indecomposable with orbit dimension  $d$  then  $\dim \text{Stab}_{PGL(\alpha)}(z) = \dim PGL(\alpha) - d$ , whence

$$\dim pr_1^{-1}(\text{rep}_\alpha^{\text{ind}})_{(d)} = \dim (\text{rep}_\alpha^{\text{ind}})_{(d)} + \dim PGL(\alpha) - d$$

But then,

$$\begin{aligned} p_Q(\alpha) &= \max_d (\dim (\text{rep}_\alpha^{\text{ind}})_{(d)} - d) \\ &= -\dim PGL(\alpha) + \max_d \dim pr_1^{-1}((\text{rep}_\alpha^{\text{ind}})_{(d)}) \\ &= -\dim PGL(\alpha) + \dim pr_1^{-1}(\text{rep}_\alpha^{\text{ind}} Q) \end{aligned}$$

By the characterization of indecomposables, we have  $pr_1^{-1}(\text{rep}_\alpha^{\text{ind}} Q) \subset pr_2^{-1}(U)$  where  $U$  consists of the (finitely many) conjugacy classes  $C_u$  of conjugacy classes of unipotent  $u \in PGL(\alpha)$ . But then,

$$\begin{aligned} p_Q(\alpha) &\leq -\dim PGL(\alpha) + \max_u \dim pr_2^{-1}(C_u) \\ &= -\dim PGL(\alpha) + \max_u \dim \text{rep}_\alpha(\text{sub})(u) + \dim PGL(\alpha) - \dim \text{cen}_{PGL(\alpha)}(u) \end{aligned}$$

proving the claim. Finally, as  $\dim \text{rep}_\alpha(\text{sub}) - \dim PGL(\alpha) < \dim \text{rep}_\alpha Q - \dim GL(\alpha) + 1 < 1 - q_Q(\alpha)$ , we are done.  $\square$

We will now extend this result to arbitrary roots using *reflection functors*. Let  $v_i$  be a *source vertex* of  $Q$  and let  $\alpha = (a_1, \dots, a_k)$  be a dimension vector such that  $\sum_{t(a)=v_i} a_{h(a)} \geq a_i$ , then we can consider the subset

$$\text{rep}_\alpha^{\text{mono}}(i) = \{V \in \text{rep}_\alpha Q \mid \oplus V_a : V_i \longrightarrow \oplus_{t(a)=v_i} V_{s(a)} \text{ is injective}\}$$

Clearly, all indecomposable representations are contained in  $\text{rep}_\alpha^{\text{mono}}(i)$ . Construct the *reflected quiver*  $R_i Q$  obtained from  $Q$  by reversing the direction of all arrows with tail  $v_i$ . The *reflected dimension vector*  $R_i \alpha = (r_1, \dots, r_k)$  is defined to be

$$r_j = \begin{cases} a_j & \text{if } j \neq i \\ \sum_{t(a)=i} a_{s(a)} - a_i & \text{if } j = i \end{cases}$$


then clearly we have in the reflected quiver  $R_i Q$  that  $\sum_{h(a)=i} r_{t(a)} \geq r_i$  and we define the subset

$$\text{rep}_{R_i \alpha}^{\text{epi}}(i) = \{V \in \text{rep}_{R_i \alpha} R_i Q \mid \oplus V_a : \oplus_{s(a)=i} V_{t(a)} \longrightarrow V_i \text{ is surjective}\}$$

Before stating the main result on reflection functors, we need to recall the definition of the Grassmann manifolds.


Let  $k \leq l$  be integers, then the points of the Grassmannian  $Grass_k(l)$  are in one-to-one correspondence with  $k$ -dimensional subspaces of  $\mathbb{C}^l$ . For example, if  $k = 1$  then  $Grass_1(l) = \mathbb{P}^{l-1}$ . We know that projective space can be covered by affine spaces defining a manifold structure on it. Also Grassmannians admit a cover by affine spaces.

Let  $W$  be a  $k$ -dimensional subspace of  $\mathbb{C}^l$  then fixing a basis  $\{w_1, \dots, w_k\}$  of  $W$  determines an  $k \times l$  matrix  $M$  having as  $i$ -th row the coordinates of  $w_i$  with respect to the standard basis of  $\mathbb{C}^l$ . Linear independence of the vectors  $w_i$  means that there is a barcode design  $I$  on  $M$



where  $I = 1 \leq i_1 < i_2 < \dots < i_k \leq l$  such that the corresponding  $k \times k$  minor  $M_I$  of  $M$  is invertible. Observe that  $M$  can have several such designs.

Conversely, given a  $k \times l$  matrix  $M$  of rank  $k$  determines a  $k$ -dimensional subspace of  $l$  spanned by the transposed rows. Two  $k \times l$   $M$  and  $M'$  matrices of rank  $k$  determine the same subspace provided there is a basechange matrix  $g \in GL_k$  such that  $gM = M'$ . That is, we can identify  $Grass_k(l)$  with the orbit space of the linear action of  $GL_k$  by left multiplication on the open set  $M_{k \times l}^{max}(\mathbb{C})$  of matrices of maximal rank. Let  $I$  be a barcode design and consider the subset of  $Grass_k(l)(I)$  of subspaces having a matrix representation  $M$  having  $I$  as barcode design. Multiplying on the left with  $M_I^{-1}$  the  $GL_k$ -orbit  $\mathcal{O}_M$  has a unique representant  $N$  with  $N_I = \mathbb{1}_k$ . Conversely, any matrix  $N$  with  $N_I = \mathbb{1}_k$  determines a point in  $Grass_k(l)(I)$ . Thus,  $Grass_k(l)(I)$  depends on  $k(l - k)$  free parameters (the entries of the negative of the barcode)



and we have an identification  $Grass_k(l)(I) \xrightarrow{\pi_I} \mathbb{C}^{k(l-k)}$ . For a different barcode design  $I'$  the image  $\pi_{I'}(Grass_k(l)(I) \cap Grass_k(l)(I'))$  is an open subset of  $\mathbb{C}^{k(l-k)}$  (one extra nonsingular minor condition) and  $\pi_{I'} \circ \pi_I^{-1}$  is a diffeomorphism on this set. That is, the maps  $\pi_I$  provide us with an atlas and determine a manifold structure on  $Grass_k(l)$ .

**THEOREM 4.38.** *For the quotient Zariski topology, we have an homeomorphism*

$$rep_{\alpha}^{mono}(i)/GL(\alpha) \xrightarrow{\cong} rep_{R_i\alpha}^{epi}(i)/GL(R_i\alpha)$$

such that corresponding representations have isomorphic endomorphism rings.

In particular, the number of parameters as well as the number of irreducible components of maximal dimension coincide for  $(rep_{\alpha}^{ind} Q)_{(d)}$  and  $(rep_{R_i\alpha}^{ind} R_iQ)_{(d)}$  for all dimensions  $d$ .

**PROOF.** Let  $m = \sum_{t(a)=i} a_i$ ,  $\overline{rep} = \bigoplus_{t(a) \neq i} M_{a_s(a) \times a_t(a)}(\mathbb{C})$  and  $\overline{GL} = \prod_{j \neq i} GL_{a_j}$ . We have the following isomorphisms

$$rep_{\alpha}^{mono}(i)/GL_{a_i} \xrightarrow{\cong} \overline{rep} \times Grass_{a_i}(m)$$

defined by sending a representation  $V$  to its restriction to  $\overline{rep}$  and  $im \bigoplus_{t(a)=i} V_a$ . In a similar way, sending a representation  $V$  to its restriction and  $ker \bigoplus_{s(a)=i} V_a$




FIGURE 8. Reflection functor diagram.

we have

$$rep_{R_i \alpha}^{epi}(i)/GL_{r_i} \xrightarrow{\cong} \overline{rep} \times Grass_{a_i}(m)$$

But then, the first claim follows from the diagram of figure 8. If  $V \in rep_{\alpha} Q$  and  $V' \in rep_{R_i \alpha} R_i Q$  with images respectively  $v$  and  $v'$  in  $\overline{rep} \times Grass_{a_i}(m)$ , we have isomorphisms

$$\begin{cases} Stab_{\overline{GL} \times GL_{a_i}}(V) & \xrightarrow{\cong} Stab_{\overline{GL}}(v) \\ Stab_{\overline{GL} \times GL_{r_i}}(V') & \xrightarrow{\cong} Stab_{\overline{GL}}(v') \end{cases}$$

from which the claim about endomorphisms follows. □

A similar results holds for *sink vertices*, hence we can apply these *Bernstein-Gelfand- Ponomarev reflection functors* iteratively using a sequence of *admissible vertices* (that is, either a source or a sink).

To a vertex  $v_i$  in which  $Q$  has no loop, we define a *reflection*  $\mathbb{Z}^k \xrightarrow{r_i} \mathbb{Z}^k$  by

$$r_i(\alpha) = \alpha - T_Q(\alpha, \epsilon_i)$$

The *Weyl group of the quiver*  $Q$   $Weyl_Q$  is the subgroup of  $GL_k(\mathbb{Z})$  generated by all reflections  $r_i$ .

A *root* of the quiver  $Q$  is a dimension vector  $\alpha \in \mathbb{N}^k$  such that  $rep_{\alpha} Q$  contains indecomposable representations. All roots have connected support. A root is said to be

$$\begin{cases} real & \text{if } \mu(rep_{\alpha}^{ind} Q) = 0 \\ imaginary & \text{if } \mu(rep_{\alpha}^{ind} Q) \geq 1 \end{cases}$$

For a fixed quiver  $Q$  we will denote the set of all roots, real roots and imaginary roots respectively by  $\Delta, \Delta_{re}$  and  $\Delta_{im}$ . With  $\Pi$  we denote the set  $\{\epsilon_i \mid v_i \text{ has no loops}\}$ . the main result on indecomposable representations is due to V. Kac .

**THEOREM 4.39.** *With notations as before, we have*

- (1)  $\Delta_{re} = \text{Weyl}_Q \cdot \Pi \cap \mathbb{N}^k$  and if  $\alpha \in \Delta_{re}$ , then  $\text{rep}_\alpha^{\text{ind}} Q$  is one orbit.  
(2)  $\Delta_{im} = \text{Weyl}_Q \cdot F_Q \cap \mathbb{N}^k$  and if  $\alpha \in \Delta_{im}$  then

$$p_Q(\alpha) = \mu(\text{rep}_\alpha^{\text{ind}} Q) = 1 - q_Q(\alpha)$$

For a sketch of the proof we refer to [23, §7], full details can be found in the lecture notes [47].

#### 4.8. Canonical decomposition.

In this section we will determine the canonical decomposition. We need a technical result.

LEMMA 4.40. *Let  $W \twoheadrightarrow W'$  be an epimorphism of  $\mathbb{C}Q$ -representations. Then, for any  $\mathbb{C}Q$ -representation  $V$  we have that the canonical map*

$$\text{Ext}_{\mathbb{C}Q}^1(V, W) \twoheadrightarrow \text{Ext}_{\mathbb{C}Q}^1(V, W')$$

*is surjective. If  $W \hookrightarrow W'$  is a monomorphism of  $\mathbb{C}Q$ -representations, then the canonical map*

$$\text{Ext}_{\mathbb{C}Q}^1(W', V) \twoheadrightarrow \text{Ext}_{\mathbb{C}Q}^1(W, V)$$

*is surjective.*

PROOF. From the proof of theorem 3.29 we have the exact diagram

$$\begin{array}{ccccccc} \bigoplus_{v_i \in Q_v} \text{Hom}_{\mathbb{C}}(V_i, W_i) & \xrightarrow{d_W^V} & \bigoplus_{a \in Q_a} \text{Hom}_{\mathbb{C}}(V_{s(a)}, W_{t(a)}) & \rightarrow & \text{Ext}_{\mathbb{C}Q}^1(V, W) & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow & & \\ \bigoplus_{v_i \in Q_v} \text{Hom}_{\mathbb{C}}(V_i, W'_i) & \xrightarrow{d_{W'}^V} & \bigoplus_{a \in Q_a} \text{Hom}_{\mathbb{C}}(V_{s(a)}, W'_{t(a)}) & \rightarrow & \text{Ext}_{\mathbb{C}Q}^1(V, W') & \longrightarrow & 0 \end{array}$$

and applying the *snake lemma* gives the result. The second part is proved similarly.  $\square$

LEMMA 4.41. *If  $V = V' \oplus V'' \in \text{rep}_\alpha(\text{max})$ , then  $\text{Ext}_{\mathbb{C}Q}^1(V', V'') = 0$ .*

PROOF. Assume  $\text{Ext}_{\mathbb{C}Q}^1(V', V'') \neq 0$ , that is, there is a non-split exact sequence

$$0 \longrightarrow V'' \longrightarrow W \longrightarrow V' \longrightarrow 0$$

then it follows from section 2.3 that  $\mathcal{O}(V) \subset \overline{\mathcal{O}(W)} - \mathcal{O}(W)$ , whence  $\dim \mathcal{O}(W) > \dim \mathcal{O}(V)$  contradicting the assumption that  $V \in \text{rep}_\alpha(\text{max})$ .  $\square$

LEMMA 4.42. *If  $W, W'$  are indecomposable representation with  $\text{Ext}_{\mathbb{C}Q}^1(W, W') = 0$ , then any non-zero map  $W' \xrightarrow{\phi} W$  is an epimorphism or a monomorphism. In particular, if  $W$  is indecomposable with  $\text{Ext}_{\mathbb{C}Q}^1(W, W) = 0$ , then  $\text{End}_{\mathbb{C}Q}(W) \simeq \mathbb{C}$ .*

PROOF. Assume  $\phi$  is neither mono- nor epimorphism then decompose  $\phi$  into

$$W' \xrightarrow{\epsilon} U \xrightarrow{\mu} W$$

As  $\epsilon$  is epi, we get a surjection from lemma 4.40

$$\text{Ext}_{\mathbb{C}Q}^1(W/U, W') \twoheadrightarrow \text{Ext}_{\mathbb{C}Q}^1(W/U, U)$$

giving a representation  $V$  fitting into the exact diagram of extensions

$$\begin{array}{ccccccccc} 0 & \longrightarrow & W' & \xrightarrow{\mu'} & V & \longrightarrow & W'/U & \longrightarrow & 0 \\ & & \downarrow \epsilon & & \downarrow \epsilon' & & \downarrow id & & \\ 0 & \longrightarrow & U & \xrightarrow{\mu} & W & \longrightarrow & W'/U & \longrightarrow & 0 \end{array}$$

from which we construct an exact sequence of representations

$$0 \longrightarrow W' \xrightarrow{\begin{bmatrix} \epsilon \\ -\mu' \end{bmatrix}} U \oplus V \xrightarrow{\begin{bmatrix} \mu & \epsilon' \end{bmatrix}} W \longrightarrow 0$$

This sequence cannot split as otherwise we would have  $W \oplus W' \simeq U \oplus V$  contradicting uniqueness of decompositions, whence  $Ext_{\mathbb{C}Q}^1(W, W') \neq 0$ , a contradiction.

For the second part, as  $W$  is finite dimensional it follows that  $End_{\mathbb{C}Q}(W)$  is a (finite dimensional) division algebra whence it must be  $\mathbb{C}$ .  $\square$

**DEFINITION 4.43.** A representation  $V \in rep_{\alpha} Q$  is said to be a *Schur representation* if  $End_{\mathbb{C}Q}(V) = \mathbb{C}$ . The dimension vector  $\alpha$  of a Schur representation is said to be a *Schur root*.

**THEOREM 4.44.**  $\alpha$  is a Schur root if and only if there is a Zariski open subset of  $rep_{\alpha} Q$  consisting of indecomposable representations.

**PROOF.** If  $V \in rep_{\alpha} Q$  is a Schur representation,  $V \in rep_{\alpha}(max)$  and therefore all representations in the dense open subset  $rep_{\alpha}(max)$  have endomorphism ring  $\mathbb{C}$  and are therefore indecomposable. Conversely, let  $Ind \hookrightarrow rep_{\alpha} Q$  be an open subset of indecomposable representations and assume that for  $V \in Ind$  we have  $Stab_{GL(\alpha)}(V) \neq \mathbb{C}^*$  and consider  $\phi_0 \in Stab_{GL(\alpha)}(V) - \mathbb{C}^*$ . For any  $g \in GL(\alpha)$  we define the *set of fixed elements*

$$rep_{\alpha}(g) = \{W \in rep_{\alpha} Q \mid g.W = W\}$$

Define the subset of  $GL(\alpha)$

$$S = \{g \in GL(\alpha) \mid dim rep_{\alpha}(g) = dim rep_{\alpha}(\phi_0)\}$$

which has no intersection with  $\mathbb{C}^*(\mathbb{1}_{d_1}, \dots, \mathbb{1}_{d_k})$  as  $\phi_0 \notin \mathbb{C}^*$ . Consider the subbundle of the trivial vectorbundle over  $S$

$$\mathcal{B} = \{(s, W) \in S \times rep_{\alpha} Q \mid s.W = W\} \hookrightarrow S \times rep_{\alpha} Q \xrightarrow{p} S$$

As all fibers have equal dimension, the restriction of  $p$  to  $\mathcal{B}$  is a *flat morphism* whence *open*. In particular, the image of the open subset  $\mathcal{B} \cap S \times Ind$

$$S' = \{g \in S \mid \exists W \in Ind : g.W = W\}$$

is an open subset of  $S$ . Now,  $S$  contains a dense set of semisimple elements, see for example [47, (2.5)], whence so does  $S' = \cup_{W \in Ind} End_{\mathbb{C}Q}(W) \cap S$ . But then one of the  $W \in Ind$  must have a torus of rank greater than one in its stabilizer subgroup contradicting indecomposability.  $\square$

Schur roots give rise to principal  $PGL(\alpha) = GL(\alpha)/C^*$ -fibrations, and hence to quiver orders and division algebras.

PROPOSITION 4.45. *If  $\alpha = (a_1, \dots, a_k)$  is a Schur root, then there is a  $GL(\alpha)$ -stable affine open subvariety  $U_\alpha$  of  $rep_\alpha Q$  such that generic orbits are closed in  $U$ .*

PROOF. Let  $T_k = \mathbb{C}^* \times \dots \times \mathbb{C}^*$  the  $k$ -dimensional torus in  $GL(\alpha)$ . Consider the semisimple subgroup  $SL(\alpha) = SL_{a_1} \times \dots \times SL_{a_k}$  and consider the corresponding quotient map

$$rep_\alpha Q \xrightarrow{\pi_s} rep_\alpha Q / SL(\alpha)$$

As  $GL(\alpha) = T_k SL(\alpha)$ ,  $T_k$  acts on  $rep_\alpha Q / SL(\alpha)$  and the generic stabilizer subgroup is trivial by the Schurian condition. Hence, there is a  $T_k$ -invariant open subset  $U_1$  of  $rep_\alpha Q / SL(\alpha)$  such that  $T_k$ -orbits are closed. But then, according to [36, §2, Thm.5] there is a  $T_k$ -invariant affine open  $U_2$  in  $U_1$ . Because the quotient map  $\psi_s$  is an affine map,  $U = \psi_s^{-1}(U_2)$  is an affine  $GL(\alpha)$ -stable open subvariety of  $rep_\alpha Q$ . Let  $x$  be a generic point in  $U$ , then its orbit

$$\mathcal{O}(x) = GL(\alpha).x = T_k SL(\alpha).x = T_k(\psi_s^{-1}(\psi_s(x))) = \psi_s^{-1}(T_k.\psi_s(x))$$

is the inverse image under the quotient map of a closed set, hence is itself closed.  $\square$

If we define  $\mathbb{T}_\alpha^s Q$  to be the ring of  $GL(\alpha)$ -equivariant maps from  $U_\alpha$  to  $M_n(\mathbb{C})$ , then this Schurian quiver order has simple  $\alpha$ -dimensional representations. Then, extending the argument of proposition 4.28 we have that the quotient map  $rep_\alpha Q \twoheadrightarrow iss_\alpha Q$  is a principal  $PGL(\alpha)$ -fibration in the étale topology over the Azumaya locus of the Schurian quiver order  $\mathbb{T}_\alpha^s Q$ . Recall that  $H_{\text{ét}}^1(X, PGL(\alpha))$  classifies twisted forms of  $M_n(\mathbb{C})$  (where  $n = \sum_a a_i$ ) as  $C_k$ -algebra. That is, Azumaya algebras over  $X$  with a distinguished embedding of  $C_k$  that are split by an étale cover on which this embedding is conjugate to the standard  $\alpha$ -embedding of  $C_k$  in  $M_n(\mathbb{C})$ . The class in the Brauer group of the function field of  $iss_\alpha \mathbb{T}_\alpha^s Q$  determined by the quiver order  $\mathbb{T}_\alpha^s Q$  is rather special.

PROPOSITION 4.46. *If  $\alpha = (a_1, \dots, a_k)$  is a Schur root of  $Q$  such that  $\gcd(a_1, \dots, a_k) = 1$ , then  $\mathbb{T}_\alpha^s Q$  determines the trivial class in the Brauer group.*

PROOF. Let  $A$  be an Azumaya localization of  $\mathbb{T}_\alpha^s Q$ . By assumption, the natural map between the  $K$ -groups  $K_0(C_k) \rightarrow K_0(M_n(\mathbb{C}))$  is surjective, whence the same is true for  $A$  proving that the class of  $A$  is split by a Zariski cover, that is  $rep_\alpha Q \simeq X \times PGL(\alpha)$  where  $X = iss_\alpha A$ .  $\square$

PROPOSITION 4.47. *If  $\alpha$  lies in the fundamental region  $F_Q$  and  $\text{supp}(\alpha)$  is not a tame quiver, then,  $\alpha$  is a Schur root.*

PROOF. Let  $\alpha = \beta_1 + \dots + \beta_s$  be the canonical decomposition of  $\alpha$  (some  $\beta_i$  may occur with higher multiplicity) and assume that  $s \geq 2$ . By definition, the image of

$$GL(\alpha) \times (rep_{\beta_1} Q \times \dots \times rep_{\beta_s} Q) \xrightarrow{\phi} rep_\alpha Q$$

is dense and  $\phi$  is constant on orbits of the free action of  $GL(\alpha)$  on the left hand side given by  $h.(g, V) = (gh^{-1}, h.V)$ . But then,

$$\dim GL(\alpha) + \sum_i \dim rep_{\beta_i} Q - \sum_i \dim GL(\beta_i) \geq \dim rep_\alpha Q$$

whence  $q_Q(\alpha) \geq \sum_i q_Q(\beta_i)$  and lemma 4.35 finishes the proof.  $\square$

Next, we want to describe morphisms between quiver-representations. Let  $\alpha = (a_1, \dots, a_k)$  and  $\beta = (b_1, \dots, b_k)$  and  $V \in \text{rep}_\alpha Q$ ,  $W \in \text{rep}_\beta Q$ . Consider the closed subvariety

$$\text{Hom}_Q(\alpha, \beta) \hookrightarrow M_{a_1 \times b_1} \oplus \dots \oplus M_{a_k \times b_k} \oplus \text{rep}_\alpha Q \oplus \text{rep}_\beta Q$$

consisting of the triples  $(\phi, V, W)$  where  $\phi = (\phi_1, \dots, \phi_k)$  is a morphism of quiver-representations  $V \rightarrow W$ . Projecting to the two last components we have an onto morphism between affine varieties

$$\text{Hom}_Q(\alpha, \beta) \xrightarrow{h} \text{rep}_\alpha Q \oplus \text{rep}_\beta Q$$

In theorem 2.4 we have proved that the dimension of fibers is an upper-semicontinuous function. That is, for every natural number  $d$ , the set

$$\{\Phi \in \text{Hom}_Q(\alpha, \beta) \mid \dim_\Phi h^{-1}(h(\Phi)) \leq d\}$$

is a Zariski open subset of  $\text{Hom}_Q(\alpha, \beta)$ . As the target space  $\text{rep}_\alpha Q \oplus \text{rep}_\beta Q$  is irreducible, it contains a non-empty open subset  $\text{hom}_{min}$  where the dimension of the fibers attains a minimal value. This minimal fiber dimension will be denoted by  $\text{hom}(\alpha, \beta)$ .

Similarly, we could have defined an affine variety  $\text{Ext}_Q(\alpha, \beta)$  where the fiber over a point  $(V, W) \in \text{rep}_\alpha Q \oplus \text{rep}_\beta Q$  is given by the extensions  $\text{Ext}_{\mathbb{C}Q}^1(V, W)$ . If  $\chi_Q$  is the Euler-form of  $Q$  we recall that for all  $V \in \text{rep}_\alpha Q$  and  $W \in \text{rep}_\beta Q$  we have

$$\dim_{\mathbb{C}} \text{Hom}_{\mathbb{C}Q}(V, W) - \dim_{\mathbb{C}} \text{Ext}_{\mathbb{C}Q}^1(V, W) = \chi_Q(\alpha, \beta)$$

Hence, there is also an open set  $\text{ext}_{min}$  of  $\text{rep}_\alpha Q \oplus \text{rep}_\beta Q$  where the dimension of  $\text{Ext}^1(V, W)$  attains a minimum. This minimal value we denote by  $\text{ext}(\alpha, \beta)$ . As  $\text{hom}_{min} \cap \text{ext}_{min}$  is a non-empty open subset we have the numerical equality

$$\text{hom}(\alpha, \beta) - \text{ext}(\alpha, \beta) = \chi_Q(\alpha, \beta).$$

In particular, if  $\text{hom}(\alpha, \alpha + \beta) > 0$ , there will be an open subset where the morphism  $V \xrightarrow{\phi} W$  is a monomorphism. Hence, there will be an open subset of  $\text{rep}_{\alpha + \beta} Q$  consisting of representations containing a subrepresentation of dimension vector  $\alpha$ . We say that  $\alpha$  is a general subrepresentation of  $\alpha + \beta$  and denote this with  $\alpha \hookrightarrow \alpha + \beta$ . We want to characterize this property. To do this, we introduce the quiver-Grassmannians

$$\text{Grass}_\alpha(\alpha + \beta) = \prod_{i=1}^k \text{Grass}_{a_i}(a_i + b_i)$$

which is a projective manifold.



Consider the following diagram of morphisms of reduced varieties

$$\begin{array}{ccc}
 & & rep_{\alpha+\beta} Q \\
 & \nearrow s & \uparrow pr_1 \\
 rep_{\alpha}^{\alpha+\beta} Q & \hookrightarrow & rep_{\alpha+\beta} Q \times Grass_{\alpha}(\alpha+\beta) \\
 & \searrow p & \downarrow pr_2 \\
 & & Grass_{\alpha}(\alpha+\beta)
 \end{array}$$

with the following properties

- $rep_{\alpha+\beta} Q \times Grass_{\alpha}(\alpha+\beta)$  is the trivial vectorbundle with fiber  $rep_{\alpha+\beta} Q$  over the projective smooth variety  $Grass_{\alpha}(\alpha+\beta)$  with structural morphism  $pr_2$ .
- $rep_{\alpha}^{\alpha+\beta} Q$  is the subvariety of  $rep_{\alpha+\beta} Q \times Grass_{\alpha}(\alpha+\beta)$  consisting of couples  $(W, V)$  where  $V$  is a subrepresentation of  $W$  (observe that this is for fixed  $W$  a linear condition). Because  $GL(\alpha+\beta)$  acts transitively on the Grassmannian  $Grass_{\alpha}(\alpha+\beta)$  (by multiplication on the right) we see that  $rep_{\alpha}^{\alpha+\beta} Q$  is a sub-vectorbundle over  $Grass_{\alpha}(\alpha+\beta)$  with structural morphism  $p$ . In particular,  $rep_{\alpha}^{\alpha+\beta} Q$  is a reduced variety.
- The morphism  $s$  is a projective morphism, that is, can be factored via the natural projection

$$\begin{array}{ccc}
 & & rep_{\alpha+\beta} Q \times \mathbb{P}^N \\
 & \nearrow f & \downarrow \pi_2 \\
 rep_{\alpha}^{\alpha+\beta} Q & \xrightarrow{s} & rep_{\alpha+\beta} Q
 \end{array}$$

where  $f$  is the composition of the inclusion  $rep_{\alpha}^{\alpha+\beta} Q \hookrightarrow rep_{\alpha+\beta} Q \times Grass_{\alpha}(\alpha+\beta)$  with the natural inclusion of Grassmannians in projective spaces recalled in the previous section  $Grass_{\alpha}(\alpha+\beta) \hookrightarrow \prod_{i=1}^k \mathbb{P}^{n_i}$  with the Segre embedding  $\prod_{i=1}^k \mathbb{P}^{n_i} \hookrightarrow \mathbb{P}^N$ . In particular,  $s$  is proper by [28, Thm. II.4.9], that is, maps closed subsets to closed subsets.

We are interested in the scheme-theoretic fibers of  $s$ . If  $W \in rep_{\alpha+\beta} Q$  lies in the image of  $s$ , we denote the fiber  $s^{-1}(W)$  by  $\underline{Grass}_{\alpha}(W)$ . Its geometric points are couples  $(W, V)$  where  $V$  is an  $\alpha$ -dimensional subrepresentation of  $W$ . Whereas  $\underline{Grass}_{\alpha}(W)$  is a projective scheme, it is in general neither smooth, nor irreducible nor even reduced. Therefore, in order to compute the tangent space in a point  $(W, V)$  of  $\underline{Grass}_{\alpha}(W)$  we have to clarify the functor it represents on the category  $commalg$  of commutative  $\mathbb{C}$ -algebras.

Let  $C$  be a commutative  $\mathbb{C}$ -algebra, a representation  $\mathcal{R}$  of the quiver  $Q$  over  $C$  consists of a collection  $\mathcal{R}_i = P_i$  of projective  $C$ -modules of finite rank and a collection of  $C$ -module morphisms for every arrow  $a$  in  $Q$

$$\textcircled{i} \xleftarrow{a} \textcircled{j} \quad \mathcal{R}_j = P_j \xleftarrow{\mathcal{R}_a} P_i = \mathcal{R}_i$$

The dimension vector of the representation  $\mathcal{R}$  is given by the  $k$ -tuple  $(rk_C \mathcal{R}_1, \dots, rk_C \mathcal{R}_k)$ . A subrepresentation  $\mathcal{S}$  of  $\mathcal{R}$  is determined by a collection of projective sub-summands (and not merely sub-modules)  $\mathcal{S}_i \triangleleft \mathcal{R}_i$ . In particular, for  $W \in rep_{\alpha+\beta} Q$  we define the representation  $\mathcal{W}_C$  of  $Q$  over the commutative ring  $C$  by

$$\begin{cases} (\mathcal{W}_C)_i &= C \otimes_C W_i \\ (\mathcal{W}_C)_a &= id_C \otimes_C W_a \end{cases}$$

With these definitions, we can now define the functor represented by  $\underline{Grass}_\alpha(W)$  as the functor assigning to a commutative  $\mathbb{C}$ -algebra  $C$  the set of all subrepresentations of dimension vector  $\alpha$  of the representation  $\mathcal{W}_C$ .

LEMMA 4.48. *Let  $x = (W, V)$  be a geometric point of  $\underline{Grass}_\alpha(W)$ , then*

$$T_x \underline{Grass}_\alpha(W) = Hom_{\mathbb{C}Q}(V, \frac{W}{V})$$

PROOF. The tangent space in  $x = (W, V)$  are the  $\mathbb{C}[\epsilon]$ -points of  $\underline{Grass}_\alpha(W)$  lying over  $(W, V)$ . To start, let  $V \xrightarrow{\psi} \frac{W}{V}$  be a homomorphism of representations of  $Q$  and consider a  $\mathbb{C}$ -linear lift of this map  $\tilde{\psi} : V \rightarrow W$ . Consider the  $\mathbb{C}$ -linear subspace of  $\mathcal{W}_{\mathbb{C}[\epsilon]} = \mathbb{C}[\epsilon] \otimes W$  spanned by the sets

$$\{v + \epsilon \otimes \tilde{\psi}(v) \mid v \in V\} \quad \text{and} \quad \epsilon \otimes V$$

This determines a  $\mathbb{C}[\epsilon]$ -subrepresentation of dimension vector  $\alpha$  of  $\mathcal{W}_{\mathbb{C}[\epsilon]}$  lying over  $(W, V)$  and is independent of the chosen linear lift  $\tilde{\psi}$ .

Conversely, if  $\mathcal{S}$  is a  $\mathbb{C}[\epsilon]$ -subrepresentation of  $\mathcal{W}_{\mathbb{C}[\epsilon]}$  lying over  $(W, V)$ , then  $\frac{\mathcal{S}}{\epsilon \mathcal{S}} = V \hookrightarrow W$ . But then, a  $\mathbb{C}$ -linear complement of  $\epsilon \mathcal{S}$  is spanned by elements of the form  $v + \epsilon \psi(v)$  where  $\psi(v) \in W$  and  $\epsilon \otimes \psi$  is determined modulo an element of  $\epsilon \otimes V$ . But then, we have a  $\mathbb{C}$ -linear map  $\tilde{\psi} : V \rightarrow \frac{W}{V}$  and as  $\mathcal{S}$  is a  $\mathbb{C}[\epsilon]$ -subrepresentation,  $\tilde{\psi}$  must be a homomorphism of representations of  $Q$ .  $\square$

THEOREM 4.49. *The following are equivalent*

- (1)  $\alpha \hookrightarrow \alpha + \beta$ .
- (2) *Every representation  $W \in rep_{\alpha+\beta} Q$  has a subrepresentation  $V$  of dimension  $\alpha$ .*
- (3)  $ext(\alpha, \beta) = 0$ .

PROOF. Assume 1. , then the image of the proper map  $s : rep_{\alpha+\beta}^{\alpha+\beta} Q \rightarrow rep_{\alpha+\beta} Q$  contains a Zariski open subset. As properness implies that the image of  $s$  must also be a closed subset of  $rep_{\alpha+\beta} Q$  it follows that  $Im s = rep_{\alpha+\beta} Q$ , that is 2. holds. Conversely, 2. clearly implies 1. so they are equivalent.

We compute the dimension of the vectorbundle  $rep_{\alpha+\beta}^{\alpha+\beta} Q$  over  $Grass_\alpha(\alpha + \beta)$ . Using that the dimension of a Grassmannians  $Grass_k(l)$  is  $k(l - k)$  we know that the base has dimension  $\sum_{i=1}^k a_i b_i$ . Now, fix a point  $V \hookrightarrow W$  in  $Grass_\alpha(\alpha + \beta)$ , then the fiber over it determines all possible ways in which this inclusion is a subrepresentation of quivers. That is, for every arrow in  $Q$  of the form  $\textcircled{j} \xleftarrow{a} \textcircled{i}$

we need to have a commuting diagram

$$\begin{array}{ccc} V_i & \longrightarrow & V_j \\ \downarrow & & \downarrow \\ W_i & \longrightarrow & W_j \end{array}$$

Here, the vertical maps are fixed. If we turn  $V \in \text{rep}_\alpha Q$ , this gives us the  $a_i a_j$  entries of the upper horizontal map as degrees of freedom, leaving only freedom for the lower horizontal map determined by a linear map  $\frac{W_i}{V_i} \longrightarrow W_j$ , that is, having  $b_i(a_j + b_j)$  degrees of freedom. Hence, the dimension of the vectorspace-fibers is

$$\sum_{\textcircled{j} \xleftarrow{a} \textcircled{i}} (a_i a_j + b_i(a_j + b_j))$$

giving the total dimension of the reduced variety  $\text{rep}_\alpha^{\alpha+\beta} Q$ . But then,

$$\begin{aligned} \dim \text{rep}_\alpha^{\alpha+\beta} Q - \dim \text{rep}_{\alpha+\beta} Q &= \sum_{i=1}^k a_i b_i + \sum_{\textcircled{j} \xleftarrow{a} \textcircled{i}} (a_i a_j + b_i(a_j + b_j)) \\ &\quad - \sum_{\textcircled{j} \xleftarrow{a} \textcircled{i}} (a_i + b_i)(a_j + b_j) \\ &= \sum_{i=1}^k a_i b_i - \sum_{\textcircled{j} \xleftarrow{a} \textcircled{i}} a_i b_j = \chi_Q(\alpha, \beta) \end{aligned}$$

Assume that 2. holds, then the proper map  $\text{rep}_\alpha^{\alpha+\beta} \xrightarrow{s} \text{rep}_{\alpha+\beta} Q$  is onto and as both varieties are reduced, the general fiber is a reduced variety of dimension  $\chi_Q(\alpha, \beta)$ , whence the general fiber contains points such that their tangentspaces have dimension  $\chi_Q(\alpha, \beta)$ . By the foregoing lemma we can compute the dimension of this tangentspace as  $\dim \text{Hom}_{\mathbb{C}Q}(V, \frac{W}{V})$ . But then, as

$$\chi_Q(\alpha, \beta) = \dim_{\mathbb{C}} \text{Hom}_{\mathbb{C}Q}(V, \frac{W}{V}) - \dim_{\mathbb{C}} \text{Ext}_{\mathbb{C}Q}^1(V, \frac{W}{V})$$

it follows that  $\text{Ext}^1(V, \frac{W}{V}) = 0$  for some representation  $V$  of dimension vector  $\alpha$  and  $\frac{W}{V}$  of dimension vector  $\beta$ . But then,  $\text{ext}(\alpha, \beta) = 0$ , that is, 3. holds.

Conversely, assume that  $\text{ext}(\alpha, \beta) = 0$ . Then, for a general point  $W \in \text{rep}_{\alpha+\beta} Q$  in the image of  $s$  and for a general point in its fiber  $(W, V) \in \text{rep}_\alpha^{\alpha+\beta} Q$  we have  $\dim_{\mathbb{C}} \text{Ext}_{\mathbb{C}Q}^1(V, \frac{W}{V}) = 0$  whence  $\dim_{\mathbb{C}} \text{Hom}_{\mathbb{C}Q}(V, \frac{W}{V}) = \chi_Q(\alpha, \beta)$ . But then, the general fiber of  $s$  has dimension  $\chi_Q(\alpha, \beta)$  and as this is the difference in dimension between the two irreducible varieties, the map is generically onto. Finally, properness of  $s$  then implies that it is onto, giving 2. and finishing the proof.  $\square$

**PROPOSITION 4.50.** *Let  $\alpha$  be a Schur root such that  $\chi_Q(\alpha, \alpha) < 0$ , then for any integer  $n$  we have that  $n\alpha$  is a Schur root.*

PROOF. There are infinitely many non-isomorphic Schur representations of dimension vector  $\alpha$ . Pick  $n$  of them  $\{W_1, \dots, W_n\}$  and from  $\chi_Q(\alpha, \alpha) < 0$  we deduce

$$\text{Hom}_{\mathbb{C}Q}(W_i, W_j) = \delta_{ij}\mathbb{C} \quad \text{and} \quad \text{Ext}_{\mathbb{C}Q}^1(W_i, W_j) \neq 0$$

By lemma 4.40 we can construct a representation  $V_n$  having a filtration

$$0 = V_0 \subset V_1 \subset \dots \subset V_n \quad \text{with} \quad \frac{V_j}{V_{j-1}} \simeq W_j$$

and such that the short exact sequences  $0 \longrightarrow V_{j-1} \longrightarrow V_j \longrightarrow W_j \longrightarrow 0$  do not split. By induction on  $n$  we may assume that  $\text{End}_{\mathbb{C}Q}(V_{n-1}) = \mathbb{C}$  and we have that  $\text{Hom}_{\mathbb{C}Q}(V_{n-1}, W_n) = 0$ . But then, the restriction of any endomorphism  $\phi$  of  $V_n$  to  $V_{n-1}$  must be an endomorphism of  $V_{n-1}$  and therefore a scalar  $\lambda\mathbb{1}$ . Hence,  $\phi - \lambda\mathbb{1} \in \text{End}_{\mathbb{C}Q}(V_n)$  is trivial on  $V_{n-1}$ . As  $\text{Hom}_{\mathbb{C}Q}(W_n, V_{n-1}) = 0$ ,  $\text{End}_{\mathbb{C}Q}(W_n) = \mathbb{C}$  and non-splitness of the sequence  $0 \longrightarrow V_{n-1} \longrightarrow V_n \longrightarrow W_n \longrightarrow 0$  we must have  $\phi - \lambda\mathbb{1} = 0$  whence  $\text{End}_{\mathbb{C}Q}(V_n) = \mathbb{C}$ , that is,  $n\alpha$  is a Schur root.  $\square$

We say that a dimension vector  $\alpha$  is *left orthogonal* to  $\beta$  if  $\text{hom}(\alpha, \beta) = 0$  and  $\text{ext}(\alpha, \beta) = 0$ .

DEFINITION 4.51. An ordered sequence  $C = (\beta_1, \dots, \beta_s)$  of dimension vectors is said to be a *compartment* for  $Q$  if and only if

- (1) for all  $i$ ,  $\beta_i$  is a Schur root,
- (2) for all  $i < j$ ,  $\beta_i$  is left orthogonal to  $\beta_j$ ,
- (3) for all  $i < j$  we have  $\chi_Q(\beta_j, \beta_i) \geq 0$ .

THEOREM 4.52. *Suppose that  $C = (\beta_1, \dots, \beta_s)$  is a compartment for  $Q$  and that there are non-negative integers  $e_1, \dots, e_s$  such that  $\alpha = e_1\beta_1 + \dots + e_s\beta_s$ . Assume that  $e_i = 1$  whenever  $\chi_Q(\beta_i, \beta_i) < 0$ . Then,*

$$\tau_{can} = (e_1, \beta_1; \dots; e_s, \beta_s)$$

*is the canonical decomposition of the dimension vector  $\alpha$ .*

PROOF. Let  $V$  be a generic representation of dimension vector  $\alpha$  with decomposition into indecomposables

$$V = W_1^{\oplus e_1} \oplus \dots \oplus W_s^{\oplus e_s} \quad \text{with} \quad \dim(W_i) = \beta_i$$

we will show that (after possibly renumbering the factors  $(\beta_1, \dots, \beta_s)$  is a compartment for  $Q$ . To start, it follows from lemma 4.41 that for all  $i \neq j$  we have  $\text{Ext}_{\mathbb{C}Q}^1(W_i, W_j) = 0$ . From lemma 4.42 we deduce a partial ordering  $i \rightarrow j$  on the indices whenever  $\text{Hom}_{\mathbb{C}Q}(W_i, W_j) \neq 0$ . Indeed, any non-zero morphism  $W_i \longrightarrow W_j$  is either a mono- or an epimorphism, assume  $W_i \twoheadrightarrow W_j$  then there can be no monomorphism  $W_j \hookrightarrow W_i$  as the composition  $W_i \twoheadrightarrow W_j \hookrightarrow W_i$  would be neither mono nor epi. That is, all non-zero morphisms from  $W_i$  to factors must be (proper) epi and we cannot obtain cycles in this way by counting dimensions. If  $W_i \hookrightarrow W_j$ , a similar argument proves the claim. From now on we assume that the chosen index-ordering of the factors is (reverse) compatible with the partial ordering  $i \rightarrow j$ , that is  $\text{Hom}(W_i, W_j) = 0$  whenever  $i < j$ , that is,  $\beta_i$  is left orthogonal to  $\beta_j$  whenever  $i < j$ . As  $\text{Ext}_{\mathbb{C}Q}^1(W_j, W_i) = 0$ , it follows that  $\chi_Q(\beta_j, \beta_i) \geq 0$ . As generic representations are open it follows that all  $\text{rep}_{\beta_i} Q$  have an open subset of indecomposables, proving that the  $\beta_i$  are Schur roots. Finally, it follows from proposition 4.50 that a Schur root  $\beta_i$  with  $\chi_Q(\beta_i, \beta_i) < 0$  can occur only with multiplicity one in any canonical decomposition.

Conversely, assume that  $(\beta_1, \dots, \beta_s)$  is a compartment for  $Q$ ,  $\alpha = \sum_i e_i \beta_i$  satisfying the requirements on multiplicities. Choose Schur representations  $W_i \in \text{rep}_{\beta_i} Q$ , then we have to prove that

$$V = W_1^{\oplus e_1} \oplus \dots \oplus W_s^{\oplus e_s}$$

is a generic representation of dimension vector  $\alpha$ . In view of the properties of the compartment we already know that  $\text{Ext}_{\mathbb{C}Q}^1(W_i, W_j) = 0$  for all  $i < j$  and we need to show that  $\text{Ext}_{\mathbb{C}Q}^1(W_j, W_i) = 0$ . Indeed, if this condition is satisfied we have

$$\begin{aligned} \dim \text{rep}_{\alpha} Q - \dim \mathcal{O}(V) &= \dim_{\mathbb{C}} \text{Ext}^1(V, V) \\ &= \sum_i e_i^2 \dim_{\mathbb{C}} \text{Ext}^1(W_i, W_i) = \sum_i e_i^2 (1 - q_Q(\beta_i)) \end{aligned}$$

We know that the Schur representations of dimension vector  $\beta_i$  depend on  $1 - q_Q(\beta_i)$  parameters by Kac's theorem 4.39 and  $e_i = 1$  unless  $q_Q(\beta_i) = 1$ . Therefore, the union of all orbits of representations with the same Schur-decomposition type as  $V$  contain a dense open set of  $\text{rep}_{\alpha} Q$  and so this must be the canonical decomposition.

If this extension space is nonzero,  $\text{Hom}_{\mathbb{C}Q}(W_j, W_i) \neq 0$  as  $\chi_Q(\beta_j, \beta_i) \geq 0$ . But then by lemma 4.42 any non-zero homomorphism from  $W_j$  to  $W_i$  must be either a mono or an epi. Assume it is a mono, so  $\beta_j < \beta_i$ , so in particular a general representation of dimension  $\beta_i$  contains a subrepresentation of dimension  $\beta_j$  and hence by theorem 4.49 we have  $\text{ext}(\beta_j, \beta_i - \beta_j) = 0$ . Suppose that  $\beta_j$  is a real Schur root, then  $\text{Ext}_{\mathbb{C}Q}^1(W_j, W_j) = 0$  and therefore also  $\text{ext}(\beta_j, \beta_i) = 0$  as  $\text{Ext}_{\mathbb{C}Q}^1(W_j, W_j \oplus (W_j/W_i)) = 0$ . If  $\beta_j$  is not a real root, then for a general representation  $S \in \text{rep}_{\beta_j} Q$  take a representation  $R \in \text{rep}_{\beta_i} Q$  in the open set where  $\text{Ext}_{\mathbb{C}Q}^1(S, R) = 0$ , then there is a monomorphism  $S \hookrightarrow R$ . Because  $\text{Ext}_{\mathbb{C}Q}^1(S, S) \neq 0$  we deduce from lemma 4.40 that  $\text{Ext}_{\mathbb{C}Q}^1(R, S) \neq 0$  contradicting the fact that  $\text{ext}(\beta_i, \beta_j) = 0$ . If the nonzero morphism  $W_j \rightarrow W_i$  is epi one has a similar argument.  $\square$


This result can be used to obtain a fairly efficient algorithm to compute the canonical decomposition in case the quiver  $Q$  has *no oriented cycles*. Fortunately, one can reduce the general problem to that of quiver without oriented cycles using the *bipartite double*  $Q^b$  of  $Q$ . We double the vertex-set of  $Q$  in a left and right set of vertices, that is

$$Q_v^b = \{v_1^l, \dots, v_k^l, v_1^r, \dots, v_k^r\}$$

To every arrow  $a \in Q_a$  from  $v_i$  to  $v_j$  we assign an arrow  $\tilde{a} \in Q_a^b$  from  $v_i^l$  to  $v_j^r$ . In addition, we have for each  $1 \leq i \leq k$  one extra arrow  $\tilde{i}$  in  $Q_a^b$  from  $v_i^l$  to  $v_i^r$ . If  $\alpha = (a_1, \dots, a_k)$  is a dimension vector for  $Q$ , the associated dimension vector  $\tilde{\alpha}$  for  $Q^b$  has components

$$\tilde{\alpha} = (a_1, \dots, a_k, a_1, \dots, a_k).$$

EXAMPLE 4.53. Consider the quiver  $Q$  and dimension vector  $\alpha = (a, b)$  on the left hand side, then



the bipartite quiver situation  $Q^b$  and  $\tilde{\alpha}$  is depicted on the right hand side.

If the canonical decomposition of  $\alpha$  for  $Q$  is  $\tau_{can} = (e_1, \beta_1; \dots; e_s, \beta_s)$ , then the canonical decomposition of  $\tilde{\alpha}$  for  $Q^b$  is  $(e_1, \tilde{\beta}_1; \dots; e_s, \tilde{\beta}_s)$  as for a general representation of  $Q^b$  of dimension vector  $\tilde{\alpha}$  the morphisms corresponding to  $\tilde{i}$  for  $1 \leq i \leq k$  are all invertible matrices and can be used to identify the left and right vertex sets, that is, there is an equivalence of categories between representations of  $Q^b$  where all the maps  $\tilde{i}$  are invertible and representations of the quiver  $Q$ . That is, the algorithm below can be applied to  $(Q^b, \tilde{\alpha})$  to obtain the canonical decomposition of  $\alpha$  for an arbitrary quiver  $Q$ .

Let  $Q$  be a quiver *without oriented cycles* then we can order the vertices  $\{v_1, \dots, v_k\}$  such that there are no oriented paths from  $v_i$  to  $v_j$  whenever  $i < j$  (start with a sink of  $Q$ , drop it and continue recursively). For example, for the bipartite quiver  $Q^b$  we first take all the right vertices and then the left ones.

input : quiver  $Q$ , ordered set of vertices as above, dimension vector  $\alpha = (a_1, \dots, a_k)$  and type  $\tau = (a_1, \vec{v}_1; \dots; a_k, \vec{v}_k)$  where  $\vec{v}_i = (\delta_{ij})_j = \dim v_i$  is the canonical basis. By the assumption on the ordering of vertices we have that  $\tau$  is a *good type* for  $\alpha$ . We say that a type  $(f_1, \gamma_1; \dots; f_s, \gamma_s)$  is a good type for  $\alpha$  if  $\alpha = \sum_i f_i \gamma_i$  and the following properties are satisfied

- (1)  $f_i \geq 0$  for all  $i$ ,
- (2)  $\gamma_i$  is a Schur root,
- (3) for each  $i < j$ ,  $\gamma_i$  is left orthogonal to  $\gamma_j$ ,
- (4)  $f_i = 1$  whenever  $\chi_Q(\gamma_i, \gamma_i) < 0$ .

A type is said to be *excellent* provided that, in addition to the above, we also have that for all  $i < j$ ,  $\chi_Q(\alpha_j, \alpha_i) \geq 0$ . In view of theorem 4.52 the purpose of the algorithm is to transform the good type  $\tau$  into the excellent type  $\tau_{can}$ . We will describe the main loop of the algorithm on a good type  $(f_1, \gamma_1; \dots; f_s, \gamma_s)$ .

step 1 : Omit all couples  $(f_i, \gamma_i)$  with  $f_i = 0$  and verify whether the remaining type is excellent. If it is, **stop and output this type**. If not, proceed.

step 2 : Reorder the type as follows, choose  $i$  and  $j$  such that  $j - i$  is minimal and  $\chi_Q(\beta_j, \beta_i) < 0$ . Partition the intermediate entries  $\{i + 1, \dots, j - 1\}$  into the sets

- $\{k_1, \dots, k_a\}$  such that  $\chi_Q(\gamma_j, \gamma_{k_m}) = 0$ ,
- $\{l_1, \dots, l_b\}$  such that  $\chi_Q(\gamma_j, \gamma_{l_m}) > 0$ .

Reorder the couples in the type in the sequence

$$(1, \dots, i - 1, k_1, \dots, k_a, i, j, l_1, \dots, l_b, j + 1, \dots, s)$$

define  $\mu = \gamma_i$ ,  $\nu = \gamma_j$ ,  $p = f_i$ ,  $q = f_j$ ,  $\zeta = p\mu + q\nu$  and  $t = -\chi_Q(\nu, \mu)$ , then proceed.

**step 3** : Change the part  $(p, \mu; q, \nu)$  of the type according to the following scheme

- If  $\mu$  and  $\nu$  are real Schur roots, consider the subcases
  - (1)  $\chi_Q(\zeta, \zeta) > 0$ , **replace**  $(p, \mu, q, \nu)$  by  $(p', \mu'; q', \nu')$  where  $\nu'$  and  $\mu'$  are non-negative combinations of  $\nu$  and  $\mu$  such that  $\mu'$  is left orthogonal to  $\nu'$ ,  $\chi_Q(\nu', \mu') = t \geq 0$  and  $\zeta = p'\mu' + q'\nu'$  for non-negative integers  $p', q'$ .
  - (2)  $\chi_Q(\zeta, \zeta) = 0$ , **replace**  $(p, \mu; q, \nu)$  by  $(k, \zeta')$  with  $\zeta = k\zeta'$ ,  $k$  positive integer, and  $\zeta'$  an indivisible root.
  - (3)  $\chi_Q(\zeta, \zeta) < 0$ , **replace**  $(p, \mu; q, \nu)$  by  $(1, \zeta)$ .
- If  $\mu$  is a real root and  $\nu$  is imaginary, consider the subcases
  - (1) If  $p + q\chi_Q(\nu, \mu) \geq 0$ , **replace**  $(p, \mu; q, \nu)$  by  $(q, \nu - \chi_Q(\nu, \mu)\mu; p + q\chi_Q(\nu, \mu), \mu)$ .
  - (2) If  $p + q\chi_Q(\nu, \mu) < 0$ , **replace**  $(p, \mu; q, \nu)$  by  $(1, \zeta)$ .
- If  $\mu$  is an imaginary root and  $\nu$  is real, consider the subcases
  - (1) If  $q + p\chi_Q(\nu, \mu) \geq 0$ , **replace**  $(p, \mu; q, \nu)$  by  $(q + p\chi_Q(\nu, \mu), \nu; p, \mu - \chi_Q(\nu, \mu)\nu)$ .
  - (2) If  $q + p\chi_Q(\nu, \mu) < 0$ , **replace**  $(p, \mu; q, \nu)$  by  $(1, \zeta)$ .
- If  $\mu$  and  $\nu$  are imaginary roots, **replace**  $(p, \mu; q, \nu)$  by  $(1, \zeta)$ .

then go to **step 1**.

One can show that in every loop of the algorithm the number  $\sum_i f_i$  decreases, so the algorithm must stop, giving the canonical decomposition of  $\alpha$ . A consequence of this algorithm is that  $r(\alpha) + 2i(\alpha) \leq k$  where  $r(\alpha)$  is the number of real Schur roots occurring in the canonical decomposition of  $\alpha$ ,  $i(\alpha)$  the number of imaginary Schur roots and  $k$  the number of vertices of  $Q$ . For more details we refer to [20].

### References.

The étale slices are due to D. Luna [60] and in the form presented here to F. Knop [40]. For more details we refer to the lecture notes of P. Slodowy [76]. The geometric interpretation of Cayley-smoothness is due to C. Procesi [65]. The local structure description with marked quivers is due to L. Le Bruyn [55]. The description of the division algebras of quiver orders is due to L. Le Bruyn and A. Schofield [59]. The material on  $A_\infty$ -algebras owes to the notes by B. Keller [35] and the work of M. Kontsevich [43] and [42]. The determination of indecomposable dimension vectors is due to V. Kac [32], [33]. The present description owes to the lecture notes of H. Kraft and Ch. Riedtmann [47], W. Crawley-Boevey [13] and the book by P. Gabriel and A.V. Roiter [23]. The main results on the canonical decomposition are due to A. Schofield [70], lemma 4.41 and lemma 4.42 are due to D. Happel and C.M. Ringel [27]. The algorithm is due to H. Derksen and J. Weyman [20] leaning on work of A. Schofield [71].





## loc $@_n$

In this part we study the étale local structure of noncommutative varieties. If  $A @_n$  is an order in a central simple algebra of dimension  $n^2$  over the functionfield of a variety of dimension  $d$ , we show that there are only a finite number of types of étale local Cayley-smooth behavior. In section 5.3 we classify them for small dimensions  $d$ . This result is a noncommutative analogon of the classical fact that manifolds of dimension  $d$  have only one type of local analytic behavior.

Combining the classification (giving in particular information about the local splitting behavior of  $A @_n$ ) with the description of Brauer classes using étale cohomology one can then characterize the central simple algebras admitting a Cayley-smooth model. In the case of surfaces, we prove in section 5.5 that every Brauer class has a model with at most finitely many noncommutative singularities, all of which analytically quantum plane singularities. One cannot resolve these remaining singularities while staying in the category of representation spaces of algebras. The Brauer classes admitting a smooth model are those where all the branch-data are trivial, that is, where one can construct a maximal order such that all irreducible components of the ramification divisor are smooth and disjoint.

When  $A$  is not Quillen-smooth, the Cayley smooth locus of  $A @_n$  will often be as small as possible. We illustrate this in the case of quantum groups and deformed preprojective algebras. In the latter case the description of the exceptional  $\alpha$ -smooth cases is important as they will reappear in the final chapter as coadjoint orbits under the necklace Lie algebra.

When  $A @_n$  is Cayley-smooth in a point  $\xi \in iss_n A$ , we describe the fiber of the quotient map  $rep_n A \twoheadrightarrow iss_n A$  in  $\xi$ , that is, we describe the isomorphism classes of  $n$ -dimensional  $A$ -representations having a fixed Jordan-Hölder decomposition. This description depends on the Hesselink stratification of the nullcone. The usually hard problem to characterize the non-empty strata is solved by representation-theoretic means using  $\theta$ -semistable quiver representations.



## Classification.

### 5.1. Type stratification.

We fix a quiver  $Q$  and dimension vector  $\alpha$ . Closed  $GL(\alpha)$ -orbits in  $rep_\alpha Q$  correspond to isomorphism classes of semi-simple representations of  $Q$  of dimension vector  $\alpha$ . We have a quotient map

$$rep_\alpha Q \xrightarrow{\pi} rep_\alpha Q/GL(\alpha) = iss_\alpha Q$$

and we know that the coordinate ring  $\mathbb{C}[iss_\alpha Q]$  is generated by traces along oriented cycles in the quiver  $Q$ . Consider a point  $\xi \in iss_\alpha Q$  and assume that the corresponding semi-simple representation  $V_\xi$  has a decomposition

$$V_\xi = V_1^{\oplus e_1} \oplus \dots \oplus V_z^{\oplus e_z}$$

into distinct simple representations  $V_i$  of dimension vector say  $\alpha_i$  and occurring in  $V_\xi$  with multiplicity  $e_i$ . We then say that  $\xi$  is a point of *representation-type*

$$\tau = t(\xi) = (e_1, \alpha_1; \dots, e_z, \alpha_z) \quad \text{with} \quad \alpha = \sum_{i=1}^z e_i \alpha_i$$

We want to apply the slice theorem to obtain the étale  $GL(\alpha)$ -local structure of the representation space  $rep_\alpha Q$  in a neighborhood of  $V_\xi$  and the étale local structure of the quotient variety  $iss_\alpha Q$  in a neighborhood of  $\xi$ . We have to calculate the normal space  $N_\xi$  to the orbit  $\mathcal{O}(V_\xi)$  as a representation over the stabilizer subgroup  $GL(\alpha)_\xi = Stab_{GL(\alpha)}(V_\xi)$ .

Denote  $a_i = \sum_{j=1}^k a_{ij}$  where  $\alpha_i = (a_{i1}, \dots, a_{ik})$ , that is,  $a_i = \dim V_i$ . We will choose a basis of the underlying vectorspace

$$\bigoplus_{v_i \in Q_v} \mathbb{C}^{\oplus e_i a_i} \quad \text{of} \quad V_\xi = V_1^{\oplus e_1} \oplus \dots \oplus V_z^{\oplus e_z}$$

as follows : the first  $e_1 a_1$  vectors give a basis of the vertex spaces of all simple components of type  $V_1$ , the next  $e_2 a_2$  vectors give a basis of the vertex spaces of all simple components of type  $V_2$ , and so on. If  $n = \sum_{i=1}^k e_i d_i$  is the total dimension of  $V_\xi$ , then with respect to this basis, the subalgebra of  $M_n(\mathbb{C})$  generated by the representation  $V_\xi$  has the following block-decomposition

$$\begin{bmatrix} M_{a_1}(\mathbb{C}) \otimes \mathbb{1}_{e_1} & 0 & \dots & 0 \\ 0 & M_{a_2}(\mathbb{C}) \otimes \mathbb{1}_{e_2} & & 0 \\ \vdots & & \ddots & \vdots \\ 0 & 0 & \dots & M_{a_z}(\mathbb{C}) \otimes \mathbb{1}_{e_z} \end{bmatrix}$$

But then, the stabilizer subgroup

$$Stab_{GL(\alpha)}(V_\xi) \simeq GL_{e_1} \times \dots \times GL_{e_z}$$

embedded in  $GL(\alpha)$  with respect to this particular basis as

$$\begin{bmatrix} GL_{e_1}(\mathbb{C} \otimes \mathbb{1}_{a_1}) & 0 & \dots & 0 \\ 0 & GL_{e_2}(\mathbb{C} \otimes \mathbb{1}_{a_2}) & & 0 \\ \vdots & & \ddots & \vdots \\ 0 & 0 & \dots & GL_{e_z}(\mathbb{C} \otimes \mathbb{1}_{a_z}) \end{bmatrix}$$

The tangentspace to the  $GL(\alpha)$ -orbit in  $V_\xi$  is equal to the image of the natural linear map

$$Lie GL(\alpha) \longrightarrow rep_\alpha Q$$

sending a matrix  $m \in Lie GL(\alpha) \simeq M_{e_1} \oplus \dots \oplus M_{e_k}$  to the representation determined by the commutator  $[m, V_\xi] = mV_\xi - V_\xi m$ . By this we mean that the matrix  $[m, V_\xi]_a$  corresponding to an arrow  $a$  is obtained as the commutator in  $M_n(\mathbb{C})$  using the canonical embedding with respect to the above choice of basis. The kernel of this linear map is the centralizer subalgebra. That is, we have an exact sequence of  $GL(\alpha)_\xi$ -modules

$$0 \longrightarrow C_{M_n(\mathbb{C})}(V_\xi) \longrightarrow Lie GL(\alpha) \longrightarrow T_{V_\xi} \mathcal{O}(V_\xi) \longrightarrow 0$$

where

$$C_{M_n(\mathbb{C})}(V_\xi) = \begin{bmatrix} M_{e_1}(\mathbb{C} \otimes \mathbb{1}_{a_1}) & 0 & \dots & 0 \\ 0 & M_{e_2}(\mathbb{C} \otimes \mathbb{1}_{a_2}) & & 0 \\ \vdots & & \ddots & \vdots \\ 0 & 0 & \dots & M_{e_z}(\mathbb{C} \otimes \mathbb{1}_{a_z}) \end{bmatrix}$$

and the action of  $GL(\alpha)_{V_\xi}$  is given by conjugation on  $M_n(\mathbb{C})$  via the above embedding. We will now engage in a book-keeping operation counting the factors of the relevant  $GL(\alpha)_\xi$ -spaces. We identify the factors by the action of the  $GL_{e_i}$ -components of  $GL(\alpha)_\xi$

- (1) The centralizer  $C_{M_n(\mathbb{C})}(V_\xi)$  decomposes as a  $GL(\alpha)_\xi$ -module into
  - one factor  $M_{e_i}$  on which  $GL_{e_1}$  acts via conjugation and the other factors act trivially,
  - ⋮
  - one factor  $M_{e_z}$  on which  $GL_{e_z}$  acts via conjugation and the other factors act trivially.
- (2) Recall the notation  $\alpha_i = (a_{i1}, \dots, a_{ik})$ , then the Lie algebra  $Lie GL(\alpha)$  decomposes as a  $GL(\alpha)_\xi$ -module into
  - $\sum_{j=1}^k a_{1j}^2$  factors  $M_{e_1}$  on which  $GL_{e_1}$  acts via conjugation and the other factors act trivially,
  - ⋮
  - $\sum_{j=1}^k a_{zj}^2$  factors  $M_{e_z}$  on which  $GL_{e_z}$  acts via conjugation and the other factors act trivially,
  - $\sum_{j=1}^k a_{1j}a_{2j}$  factors  $M_{e_1 \times e_2}$  on which  $GL_{e_1} \times GL_{e_2}$  acts via  $\gamma_1.m.\gamma_2^{-1}$  and the other factors act trivially,
  - ⋮

- $\sum_{j=1}^k a_{zj} a_{z-1j}$  factors  $M_{e_z \times e_{z-1}}$  on which  $GL_{e_z} \times GL_{e_{z-1}}$  acts via  $\gamma_z \cdot m \cdot \gamma_{z-1}^{-1}$  and the other factors act trivially.
- (3) The representation space  $rep_\alpha Q$  decomposes as a  $GL(\alpha)_\xi$ -modulo into the following factors, for every arrow  $\textcircled{j} \xleftarrow{a} \textcircled{i}$  in  $Q$  (or every loop in  $v_i$  by setting  $i = j$  in the expressions below) we have
- $a_{1i} a_{1j}$  factors  $M_{e_1}$  on which  $GL_{e_1}$  acts via conjugation and the other factors act trivially,
  - $a_{1i} a_{2j}$  factors  $M_{e_1 \times e_2}$  on which  $GL_{e_1} \times GL_{e_2}$  acts via  $\gamma_1 \cdot m \cdot \gamma_2^{-1}$  and the other factors act trivially,
- ⋮
- $a_{zi} a_{z-1j}$  factors  $M_{e_z \times e_{z-1}}$  on which  $GL_{e_z} \times GL_{e_{z-1}}$  act via  $\gamma_z \cdot m \cdot \gamma_{z-1}^{-1}$  and the other factors act trivially,
  - $a_{zi} a_{zj}$  factors  $M_{e_z}$  on which  $GL_{e_z}$  acts via conjugation and the other factors act trivially.

Removing the factors of 1. from those of 2. we obtain a description of the tangentspace to the orbit  $T_{V_\xi} \mathcal{O}(V_\xi)$ . But then, removing these factors from those of 3. we obtain the description of the normal space  $N_{V_\xi}$  as a  $GL(\alpha)_\xi$ -module as there is an exact sequence of  $GL(\alpha)_\xi$ -modules

$$0 \longrightarrow T_{V_\xi} \mathcal{O}(V_\xi) \longrightarrow rep_\alpha Q \longrightarrow N_{V_\xi} \longrightarrow 0$$

This proves that the normal space to the orbit in  $V_\xi$  depends only on the representation type  $\tau = t(\xi)$  of the point  $\xi$  and can be identified with the representation space of a *local quiver*  $Q_\tau$ .

**THEOREM 5.1.** *Let  $\xi \in iss_\alpha Q$  be a point of representation type*

$$\tau = t(\xi) = (e_1, \alpha_1; \dots, e_z, \alpha_z)$$

*Then, the normal space  $N_{V_\xi}$  to the orbit, as a module over the stabilizer subgroup, is identical to the representation space of a local quiver situation*

$$N_{V_\xi} \simeq rep_{\alpha_\tau} Q_\tau$$

*where  $Q_\tau$  is the quiver on  $z$  vertices (the number of distinct simple components of  $V_\xi$ ) say  $\{w_1, \dots, w_z\}$  such that in  $Q_\tau$*

$$\# \textcircled{j} \xleftarrow{a} \textcircled{i} = -\chi_Q(\alpha_i, \alpha_j) \quad \text{for } i \neq j, \text{ and}$$

$$\# \textcircled{i} \begin{array}{c} \curvearrowright \\ \downarrow \end{array} = 1 - \chi_Q(\alpha_i, \alpha_i)$$

*and such that the dimension vector  $\alpha_\tau = (e_1, \dots, e_z)$  (the multiplicities of the simple components in  $V_\xi$ ).*

We can repeat this argument in the case of a marked quiver  $Q^\bullet$ . The only difference is the description of the factors of  $rep_\alpha Q^\bullet$  where we need to replace the factors  $M_{e_j}$  in the description of a loop in  $v_i$  by  $M_{e_i}^0$  (trace zero matrices) in case

the loop gets a mark in  $Q^\bullet$ . We define the *Euler form* of the marked quiver  $Q^\bullet$

$$\chi_{Q^\bullet}^1 = \begin{bmatrix} 1 - a_{11} & \chi_{12} & \cdots & \chi_{1k} \\ \chi_{21} & 1 - a_{22} & \cdots & \chi_{2k} \\ \vdots & \vdots & \ddots & \vdots \\ \chi_{k1} & \chi_{k2} & \cdots & 1 - a_{kk} \end{bmatrix} \quad \chi_{Q^\bullet}^2 = \begin{bmatrix} -m_{11} & & & \\ & -m_{22} & & \\ & & \ddots & \\ & & & -m_{kk} \end{bmatrix}$$

such that  $\chi_Q = \chi_{Q^\bullet}^1 + \chi_{Q^\bullet}^2$ , where  $Q$  is the underlying quiver of  $Q^\bullet$ .

**THEOREM 5.2.** *Let  $\xi \in \text{iss}_\alpha Q^\bullet$  be a point of representation type*

$$\tau = t(\xi) = (e_1, \alpha_1; \dots, e_z, \alpha_z)$$

*Then, the normal space  $N_{V_\xi}$  to the orbit, as a module over the stabilizer subgroup, is identical to the representation space of a local marked quiver situation*

$$N_{V_\xi} \simeq \text{rep}_{\alpha_\tau} Q_\tau^\bullet$$

*where  $Q_\tau^\bullet$  is the quiver on  $z$  vertices (the number of distinct simple components of  $V_\xi$ ) say  $\{w_1, \dots, w_z\}$  such that in  $Q_\tau^\bullet$*

$$\# \begin{array}{c} \textcircled{j} \xleftarrow{a} \textcircled{i} \end{array} = -\chi_Q(\alpha_i, \alpha_j) \quad \text{for } i \neq j, \text{ and}$$

$$\# \begin{array}{c} \textcircled{i} \\ \downarrow \\ \textcircled{i} \end{array} = 1 - \chi_{Q^\bullet}^1(\alpha_i, \alpha_i)$$

$$\# \begin{array}{c} \bullet \\ \downarrow \\ \textcircled{i} \end{array} = -\chi_{Q^\bullet}^2(\alpha_i, \alpha_i)$$

*and such that the dimension vector  $\alpha_\tau = (e_1, \dots, e_z)$  (the multiplicities of the simple components in  $V_\xi$ ).*

**PROPOSITION 5.3.** *If  $\alpha = (d_1, \dots, d_k)$  is the dimension vector of a simple representation of  $Q^\bullet$ , then the dimension of the quotient variety  $\text{iss}_\alpha Q^\bullet$  is equal to*

$$1 - \chi_{Q^\bullet}^1(\alpha, \alpha)$$

**PROOF.** There is a Zariski open subset of  $\text{iss}_\alpha Q^\bullet$  consisting of points  $\xi$  such that the corresponding semi-simple module  $V_\xi$  is simple, that is,  $\xi$  has representation type  $\tau = (1, \alpha)$ . But then the local quiver setting  $(Q_\tau, \alpha_\tau)$  is

$$\boxed{a} \begin{array}{c} \curvearrowright \\ \textcircled{1} \\ \curvearrowleft \end{array} \bullet \boxed{b}$$

where  $a = 1 - \chi_{Q^\bullet}^1(\alpha, \alpha)$  and  $b = -\chi_{Q^\bullet}^2(\alpha, \alpha)$ . The corresponding representation space has coordinate ring

$$\mathbb{C}[\text{rep}_{\alpha_\tau} Q_\tau^\bullet] = \mathbb{C}[x_1, \dots, x_a]$$

on which  $GL(\alpha_\tau) = \mathbb{C}^*$  acts trivially. That is, the quotient variety is

$$\text{rep}_{\alpha_\tau} Q_\tau^\bullet / GL(\alpha_\tau) = \text{rep}_{\alpha_\tau} Q_\tau^\bullet \simeq \mathbb{C}^a$$

By the slice theorem,  $\text{iss}_\alpha Q^\bullet$  has the same local structure near  $\xi$  as this quotient space near the origin and the result follows.  $\square$

We can extend the classifications of simple roots of a quiver to the setting of marked quivers. Let  $Q$  be the underlying quiver of a marked quiver  $Q^\bullet$ . If  $\alpha = (a_1, \dots, a_k)$  is a simple root of  $Q$  and if  $l$  is a marked loop in a vertex  $v_i$  with  $a_i > 1$ , then we can replace the matrix  $V_l$  of a simple representation  $V \in \text{rep}_\alpha Q$  by  $V'_l = V_l - \frac{1}{d_i} \mathbb{1}_{d_i}$  and retain the property that  $V'$  is a simple representation. Things are different, however, for a marked loop in a vertex  $v_i$  with  $a_i = 1$  as this  $1 \times 1$ -matrix factor is removed from the representation space. That is, we have the following characterization result.

**THEOREM 5.4.**  $\alpha = (a_1, \dots, a_k)$  is the dimension vector of a simple representation of a marked quiver  $Q^\bullet$  if and only if  $\alpha = (a_1, \dots, a_k)$  is the dimension vector of a simple representation of the quiver  $Q'$  obtained from the underlying quiver  $Q$  of  $Q^\bullet$  after removing the loops in  $Q$  which are marked in  $Q^\bullet$  in all vertices  $v_i$  where  $a_i = 1$ .

We draw some consequences from the description of the local quiver. We state all results in the setting of marked quivers. Often, the quotient varieties  $\text{iss}_\alpha Q^\bullet = \text{rep}_\alpha Q^\bullet / \text{GL}(\alpha)$  classifying isomorphism classes of semi-simple  $\alpha$ -dimensional representations have singularities. Still, we can decompose these quotient varieties in smooth pieces according to representation types.

**PROPOSITION 5.5.** Let  $\text{iss}_\alpha Q^\bullet(\tau)$  be the set of points  $\xi \in \text{iss}_\alpha Q^\bullet$  of representation type

$$\tau = (e_1, \alpha_1; \dots; e_z, \alpha_z)$$

Then,  $\text{iss}_\alpha Q^\bullet(\tau)$  is a locally closed smooth subvariety of  $\text{iss}_\alpha Q^\bullet$  and

$$\text{iss}_\alpha Q^\bullet = \bigsqcup_{\tau} \text{iss}_\alpha Q^\bullet(\tau)$$

is a finite smooth stratification of the quotient variety.

**PROOF.** Let  $Q_\tau^\bullet$  be the local marked quiver in  $\xi$ . Consider a nearby point  $\xi'$ . If some trace of an oriented cycles of length  $> 1$  in  $Q_\tau^\bullet$  is non-zero in  $\xi'$ , then  $\xi'$  cannot be of representation type  $\tau$  as it contains a simple factor composed of vertices of that cycle. That is, locally in  $\xi$  the subvariety  $\text{iss}_\alpha Q^\bullet(\tau)$  is determined by the traces of unmarked loops in vertices of the local quiver  $Q_\tau^\bullet$  and hence is locally in the étale topology an affine space whence smooth. All other statements are direct.  $\square$

Given a stratification of a topological space, one wants to determine which strata make up the boundary of a given stratum. For the above stratification of  $\text{iss}_\alpha Q^\bullet$  we have a combinatorial solution to this problem. Two representation types

$$\tau = (e_1, \alpha_1; \dots; e_z, \alpha_z) \quad \text{and} \quad \tau' = (e'_1, \alpha'_1; \dots; e'_{z'}, \alpha'_{z'})$$

are said to be direct successors  $\tau < \tau'$  if and only if one of the following two cases occurs

- (splitting of one simple) :  $z' = z + 1$  and for all but one  $1 \leq i \leq z$  we have that  $(e_i, \alpha_i) = (e'_j, \alpha'_j)$  for a uniquely determined  $j$  and for the remaining  $i_0$  we have that the remaining couples of  $\tau'$  are

$$(e_i, \alpha'_u; e_i, \alpha'_v) \quad \text{with} \quad \alpha_i = \alpha'_u + \alpha'_v$$

- (combining two simple types) :  $z' = z - 1$  and for all but one  $1 \leq i \leq z'$  we have that  $(e'_i, \alpha'_i) = (e_j, \alpha_j)$  for a uniquely determined  $j$  and for the remaining  $i$  we have that the remaining couples of  $\tau$  are

$$(e_u, \alpha'_i; e_v, \alpha'_i) \quad \text{with} \quad e_u + e_v = e'_i$$

This direct successor relation  $<$  induces an ordering which we will denote with  $<<$ . Observe that  $\tau << \tau'$  if and only if the stabilizer subgroup  $GL(\alpha)_\tau$  is conjugated to a subgroup of  $GL(\alpha)_{\tau'}$ . The following result either follows from general theory, see for example [73, lemma 5.5], or from the description of the local marked quivers.

PROPOSITION 5.6. *The stratum  $iss_\alpha Q^\bullet(\tau')$  lies in the closure of the stratum  $iss_\alpha Q^\bullet$  if and only if  $\tau << \tau'$ .*

Proposition 5.3 gives us the dimensions of the different strata  $iss_\alpha Q^\bullet(\tau)$ .

PROPOSITION 5.7. *Let  $\tau = (e_1, \alpha_1; \dots; e_z, \alpha_z)$  a representation type of  $\alpha$ . Then,*

$$\dim iss_\alpha Q^\bullet(\tau) = \sum_{j=1}^z (1 - \chi_{Q^\bullet}^1(\alpha_j, \alpha_j))$$

Because  $rep_\alpha Q^\bullet$  and hence  $iss_\alpha Q^\bullet$  is an irreducible variety, there is a unique representation type  $\tau_{gen}^{ss}$  such that  $iss_\alpha Q^\bullet(\tau_{gen}^{ss})$  is Zariski open in the quotient variety  $iss_\alpha Q^\bullet$ . We call  $\tau_{gen}^{ss}$  the *generic semi-simple representation type* for  $\alpha$ . The generic semi-simple representation type can be determined by the following algorithm.

**input :** A quiver  $Q$ , a dimension vector  $\alpha = (a_1, \dots, a_k)$  and a semi-simple representation type

$$\tau = (e_1, \alpha_1; \dots; e_l, \alpha_l)$$

with  $\alpha = \sum +i = 1^l e_i \alpha_i$  and all  $\alpha_i$  simple roots for  $Q$ . For example, we can always start with the type  $(a_1, \vec{v}_1; \dots; a_k, \vec{v}_k)$ .

**step 1 :** Compute the local quiver  $Q_\tau$  on  $l$  vertices and the dimension vector  $\alpha_\tau$ . If the only oriented cycles in  $Q_\tau$  are vertex-loops, **stop and output this type**. If not, proceed.

**step 2 :** Take a proper oriented cycle  $C = (j_1, \dots, j_r)$  with  $r \geq 2$  in  $Q_\tau$  where  $j_s$  is the vertex in  $Q_\tau$  determined by the dimension vector  $\alpha_{j_s}$ . Set  $\beta = \alpha_{j_1} + \dots + \alpha_{j_r}$ ,  $e'_i = e_i - \delta_{iC}$  where  $\delta_{iC} = 1$  if  $i \in C$  and is 0 otherwise. **replace**  $\tau$  by the new semi-simple representation type

$$\tau' = (e'_1, \alpha_1; \dots; e'_l, \alpha_l; 1, \beta)$$

**delete** the terms  $(e'_i, \alpha_i)$  with  $e'_i = 0$  and set  $\tau$  to be the resulting type. **goto step 1**.

The same algorithm extends to marked quivers with the modified construction of the local marked quiver  $Q_\tau^\bullet$  in that case. We can give an  $A_\infty$ -interpretation of the characterization of the *canonical decomposition* and the generic semi-simple representation type. Let

$$\tau = (e_1, \alpha_1; \dots; e_z, \alpha_z) \quad \alpha = \sum_{i=1}^z e_i \alpha_i$$



be a decomposition of  $\alpha$  with all the  $\alpha_i$  roots. We define  $\alpha_\tau = (e_1, \dots, e_z)$  and construct two quivers  $Q_\tau^0$  and  $Q_\tau^1$  on  $z$  vertices determined by the rules

$$\text{in } Q_\tau^0 : \quad \# \textcircled{i} \xleftarrow{a} \textcircled{i} = \dim_{\mathbb{C}} \text{Hom}_{\mathbb{C}Q}(V_i, V_j)$$

$$\text{in } Q_\tau^1 : \quad \# \textcircled{i} \xleftarrow{a} \textcircled{i} = \dim_{\mathbb{C}} \text{Ext}_{\mathbb{C}Q}^1(V_i, V_j)$$

where  $V_i$  is a general representation of  $Q$  of dimension vector  $\alpha_i$ .

**THEOREM 5.8.** *With notations as above, we have :*

- (1) *The canonical decomposition  $\tau_{can}$  is the unique type  $\tau = (e_1, \alpha_1; \dots; e_z, \alpha_z)$  such that all  $\alpha_i$  are Schur roots,  $Q_\tau^0$  has no (non-loop) oriented cycles and  $Q_\tau^1$  has no arrows and loops only in vertices where  $e_i = 1$ .*
- (2) *The generic semi-simple representation type  $\tau_{gen}^{ss}$  is the unique type  $\tau = (e_1, \alpha_1; \dots; e_z, \alpha_z)$  such that all  $\alpha_i$  are simple roots,  $Q_\tau^0$  has only loops and  $Q_\tau^1$  has no (non-loop) oriented cycles.*

## 5.2. Cayley-smooth locus.

Let  $A$  be a Cayley-Hamilton algebra of degree  $n$  equipped with a trace map  $A \xrightarrow{tr} A$  and consider the quotient map

$$\underline{rep}_n^{tr} A \xrightarrow{\pi} \underline{iss}_n^{tr} A$$

Let  $\xi$  be a geometric point of the quotient scheme  $\underline{iss}_n^{tr} A$  with corresponding  $n$ -dimensional trace preserving semi-simple representation  $V_\xi$  with decomposition

$$V_\xi = S_1^{\oplus e_1} \oplus \dots \oplus S_k^{\oplus e_k}$$

where the  $S_i$  are distinct simple representations of  $A$  of dimension  $d_i$  such that  $n = \sum_{i=1}^k d_i e_i$ .

**DEFINITION 5.9.** The Cayley-smooth locus of  $A$  is the subset of  $\underline{iss}_n^{tr} A$

$$Sm_{tr} A = \{ \xi \in \underline{iss}_n^{tr} A \mid \underline{iss}_n^{tr} A \text{ is smooth along } \pi^{-1}(\xi) \}$$

As the singular locus of  $\underline{iss}_n^{tr} A$  is a  $GL_n$ -stable closed subscheme of  $\underline{iss}_n^{tr} A$  this is equivalent to

$$Sm_{tr} A = \{ \xi \in \underline{iss}_n^{tr} A \mid \underline{iss}_n^{tr} A \text{ is smooth in } V_\xi \}$$

We will give some numerical conditions on  $\xi$  to be in the smooth locus  $Sm_{tr} A$ . To start,  $\underline{rep}_n^{tr} A$  is smooth in  $V_\xi$  if and only if the dimension of the tangent space in  $V_\xi$  is equal to the local dimension of  $\underline{rep}_n^{tr} A$  in  $V_\xi$ . From example 4.2 we know that the tangent space is the set of trace preserving derivations  $A \xrightarrow{D} M_n(\mathbb{C})$  satisfying

$$D(aa') = D(a)\rho(a') + \rho(a)D(a')$$

where  $A \xrightarrow{\rho} M_n(\mathbb{C})$  is the  $\mathbb{C}$ -algebra morphism determined by the action of  $A$  on  $V_\xi$ . The  $\mathbb{C}$ -vector space of such derivations is denoted by  $Der_\rho^t A$ . Therefore,

$$\xi \in Sm_{tr} A \iff \dim_{\mathbb{C}} Der_\rho^t A = \dim_{V_\xi} \underline{rep}_n^{tr} A$$

Next, if  $\xi \in Sm_{tr} A$ , then we know from the slice theorem that the local  $GL_n$ -structure of  $\underline{rep}_n^{tr} A$  near  $V_\xi$  is determined by a local marked quiver setting  $(Q_\xi^\bullet, \alpha_\xi)$  as defined in theorem 4.22. We have local étale isomorphisms between the varieties

$$GL_n \times^{GL(\alpha_\xi)} \underline{rep}_{\alpha_\xi} Q_\xi^\bullet \xleftarrow{et} \underline{rep}_n^{tr} A \quad \text{and} \quad \underline{rep}_{\alpha_\xi} Q_\xi^\bullet / GL(\alpha_\xi) \xleftarrow{et} \underline{iss}_n^{tr} A$$

Which gives us the following numerical restrictions on  $\xi \in Sm_{tr} A$  :

PROPOSITION 5.10.  $\xi \in Sm_{tr} A$  if and only if the following two equalities hold

$$\begin{cases} \dim_{V_\xi} \underline{rep}_n^{tr} A &= n^2 - (e_1^2 + \dots + e_k^2) + \dim_{\mathbb{C}} Ext_A^{tr}(V_\xi, V_\xi) \\ \dim_{\xi} \underline{iss}_n^{tr} A &= \dim_{\overline{0}} \underline{rep}_{\alpha_\xi} Q_\xi^\bullet / GL(\alpha_\xi) = \dim_{\overline{0}} \underline{iss}_{\alpha_\xi} Q_\xi^\bullet \end{cases}$$

Moreover, if  $\xi \in Sm_{tr} A$ , then  $\underline{rep}_n^{tr} A$  is a normal variety (that is, the coordinate ring is integrally closed) in a neighborhood of  $\xi$

PROOF. The last statement follows from the fact that  $\mathbb{C}[\underline{rep}_{\alpha_\xi} Q_\xi^\bullet]^{GL(\alpha_\xi)}$  is integrally closed and this property is preserved under the étale map.  $\square$

In general, the difference between these numbers gives a measure for the non-commutative singularity of  $A$  in  $\xi$ .

EXAMPLE 5.11. Consider the affine  $\mathbb{C}$ -algebra  $A = \frac{\mathbb{C}\langle x, y \rangle}{(xy+yx)}$  then  $u = x^2$  and  $v = y^2$  are central elements of  $A$  and  $A$  is a free module of rank 4 over  $\mathbb{C}[u, v]$ . In fact,  $A$  is a  $\mathbb{C}[u, v]$ -order in the quaternion division algebra

$$\Delta = \begin{pmatrix} u & & v \\ & \mathbb{C}(u, v) & \\ & & \end{pmatrix}$$

and the reduced trace map on  $\Delta$  makes  $A$  into a Cayley-Hamilton algebra of degree 2. More precisely,  $tr$  is the linear map on  $A$  such that

$$\begin{cases} tr(x^i y^j) = 0 & \text{if either } i \text{ or } j \text{ are odd, and} \\ tr(x^i y^j) = 2x^i y^j & \text{if } i \text{ and } j \text{ are even.} \end{cases}$$

In particular, a trace preserving 2-dimensional representation is determined by a couple of  $2 \times 2$  matrices

$$\rho = \left( \begin{bmatrix} x_1 & x_2 \\ x_3 & -x_1 \end{bmatrix}, \begin{bmatrix} x_4 & x_5 \\ x_6 & -x_4 \end{bmatrix} \right) \quad \text{with } tr \left( \begin{bmatrix} x_1 & x_2 \\ x_3 & -x_1 \end{bmatrix} \cdot \begin{bmatrix} x_4 & x_5 \\ x_6 & -x_4 \end{bmatrix} \right) = 0$$

That is,  $\underline{rep}_2^{tr} A$  is the hypersurface in  $\mathbb{C}^6$  determined by the equation

$$\underline{rep}_2^{tr} A = \mathbb{V}(2x_1x_4 + x_2x_6 + x_3x_5) \hookrightarrow \mathbb{C}^6$$

and is therefore irreducible of dimension 5 with an isolated singularity at  $p = (0, \dots, 0)$ . The image of the trace map is equal to the center of  $A$  which is  $\mathbb{C}[u, v]$  and the quotient map

$$\underline{rep}_2^{tr} A \xrightarrow{\pi} \underline{iss}_2^{tr} A = \mathbb{C}^2 \quad \pi(x_1, \dots, x_6) = (x_1^2 + x_2x_3, x_4^2 + x_5x_6)$$

There are three different representation types to consider. Let  $\xi = (a, b) \in \mathbb{C}^2 = \underline{iss}_2^{tr} A$  with  $ab \neq 0$ , then  $\pi^{-1}(\xi)$  is a closed  $GL_2$ -orbit and a corresponding simple  $A$ -module is given by the matrixcouple

$$\left( \begin{bmatrix} i\sqrt{a} & 0 \\ 0 & -i\sqrt{a} \end{bmatrix}, \begin{bmatrix} 0 & \sqrt{b} \\ -\sqrt{b} & 0 \end{bmatrix} \right)$$

That is,  $\xi$  is of type (1, 2) and the stabilizer subgroup are the scalar matrixes  $\mathbb{C}^* \mathbb{1}_2 \hookrightarrow GL_2$ . So, the action on both the tangentspace to  $\underline{rep}_2^{tr} A$  and the tangent space to the orbit are trivial. As they have respectively dimension 5 and 3, the normalspace corresponds to the quiver setting

$$N_\xi = \begin{array}{c} \circlearrowleft \\ \circlearrowright \\ \textcircled{1} \end{array}$$

which is compatible with the numerical restrictions. Next, consider a point  $\xi = (0, b)$  (or similarly,  $(a, 0)$ ), then  $\xi$  is of type  $(1, 1; 1, 1)$  and the corresponding semi-simple representation is given by the matrices

$$\left( \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} i\sqrt{b} & 0 \\ 0 & -i\sqrt{b} \end{bmatrix} \right)$$


The stabilizer subgroup is in this case the maximal torus of diagonal matrices  $\mathbb{C}^* \times \mathbb{C}^* \hookrightarrow GL_2$ . The tangent space in this point to  $rep_2^{tr} A$  are the 6-tuples  $(a_1, \dots, a_6)$  such that

$$tr \left( \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} + \epsilon \begin{bmatrix} a_1 & a_2 \\ a_3 & -a_1 \end{bmatrix} \right) \cdot \left( \begin{bmatrix} i\sqrt{b} & 0 \\ 0 & -i\sqrt{b} \end{bmatrix} + \epsilon \begin{bmatrix} b_4 & b_5 \\ b_6 & -b_4 \end{bmatrix} \right) = 0 \quad \text{where } \epsilon^2 = 0$$

This leads to the condition  $a_1 = 0$ , so the tangentspace are the matrix couples

$$\left( \begin{bmatrix} 0 & a_2 \\ a_3 & 0 \end{bmatrix}, \begin{bmatrix} a_4 & a_5 \\ a_6 & -a_4 \end{bmatrix} \right) \quad \text{on which the stabilizer } \begin{bmatrix} \lambda & 0 \\ 0 & \mu \end{bmatrix}$$

acts via conjugation. That is, the tangentspace corresponds to the quiver setting



Moreover, the tangentspace to the orbit is the image of the linear map

$$\left( \mathbb{1}_2 + \epsilon \begin{bmatrix} m_1 & m_2 \\ m_3 & m_4 \end{bmatrix} \right) \cdot \left( \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} \sqrt{b} & 0 \\ 0 & -\sqrt{b} \end{bmatrix} \right), \left( \mathbb{1}_2 - \begin{bmatrix} m_1 & m_2 \\ m_3 & m_4 \end{bmatrix} \right)$$


which is equal to

$$\left( \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} \sqrt{b} & 0 \\ 0 & -\sqrt{b} \end{bmatrix} + \epsilon \begin{bmatrix} 0 & -2m_2\sqrt{b} \\ 2m_3\sqrt{b} & 0 \end{bmatrix} \right)$$


on which the stabilizer acts again via conjugation giving the quiver setting



Therefore, the normal space to the orbit corresponds to the quiver setting



which is again compatible with the numerical restrictions. Finally, consider  $\xi = (0, 0)$  which is of type  $(2, 1)$  and whose semi-simple representation corresponds to the zero matrix-couple. The action fixes this point, so the stabilizer is  $GL_2$  and the tangent space to the orbit is the trivial space. Hence, the tangent space to  $rep_2^{tr} A$  coincides with the normalspace to the orbit and both spaces are acted on by  $GL_2$  via simultaneous conjugation leading to the quiver setting



This time, the data is not compatible with the numerical restriction as

$$5 = \dim rep_2^{tr} A \neq n^2 - e^2 + \dim rep_\alpha Q^\bullet = 4 - 4 + 6$$

consistent with the fact that the zero matrix-couple is a (in fact, the only) singularity on  $rep_2^t A$ .

We will put additional conditions on the Cayley-Hamilton algebra  $A$ . Let  $X$  be a normal affine variety with coordinate ring  $\mathbb{C}[X]$  and functionfield  $\mathbb{C}(X)$ . Let  $\Delta$  be a central simple  $\mathbb{C}(X)$ -algebra of dimension  $n^2$  which is a Cayley-Hamilton algebra of degree  $n$  using the reduced trace map  $tr$ . Let  $A$  be a  $\mathbb{C}[X]$ -order in  $\Delta$ , that is, the center of  $A$  is  $\mathbb{C}[X]$  and  $A \otimes_{\mathbb{C}[X]} \mathbb{C}(X) \simeq \Delta$ . Because  $\mathbb{C}[X]$  is integrally closed, the restriction of the reduced trace  $tr$  to  $A$  has its image in  $\mathbb{C}[X]$ , that is,  $A$  is a Cayley-Hamilton algebra of degree  $n$  and

$$tr(A) = \mathbb{C}[X]$$

Consider the quotient morphism for the representation variety

$$\underline{rep}_n^{tr} A \xrightarrow{\pi} \underline{iss}_n^{tr} A$$

then the above argument shows that  $X \simeq \underline{iss}_n^{tr} A$  and in particular the quotient scheme is reduced.

PROPOSITION 5.12. *Let  $A$  be a Cayley-Hamilton order of degree  $n$  over  $\mathbb{C}[X]$ . Then, its smooth locus  $Sm_{tr} A$  is a nonempty Zariski open subset of  $X$ . In particular, the set  $X_{az}$  of Azumaya points, that is, of points  $x \in X = \underline{iss}_n^{tr} A$  of representation type  $(1, n)$  is a non-empty Zariski open subset of  $X$  and its intersection with the Zariski open subset  $X_{reg}$  of smooth points of  $X$  satisfies*

$$X_{az} \cap X_{reg} \hookrightarrow Sm_{tr} A$$

PROOF. Because  $AC(X) = \Delta$ , there is an  $f \in \mathbb{C}[X]$  such that  $A_f = A \otimes_{\mathbb{C}[X]} \mathbb{C}[X]_f$  is a free  $\mathbb{C}[X]_f$ -module of rank  $n^2$  say with basis  $\{a_1, \dots, a_{n^2}\}$ . Consider the  $n^2 \times n^2$  matrix with entries in  $\mathbb{C}[X]_f$

$$R = \begin{bmatrix} tr(a_1 a_1) & \dots & tr(a_1 a_{n^2}) \\ \vdots & & \vdots \\ tr(a_{n^2} a_1) & \dots & tr(a_{n^2} a_{n^2}) \end{bmatrix}$$

The determinant  $d = \det R$  is nonzero in  $\mathbb{C}[X]_f$ . For, let  $\mathbb{K}$  be the algebraic closure of  $\mathbb{C}(X)$  then  $A_f \otimes_{\mathbb{C}[X]_f} \mathbb{K} \simeq M_n(\mathbb{K})$  and for any  $\mathbb{K}$ -basis of  $M_n(\mathbb{K})$  the corresponding matrix is invertible (for example, verify this on the matrixes  $e_{ij}$ ). As  $\{a_1, \dots, a_{n^2}\}$  is such a basis,  $d \neq 0$ . Next, consider the Zariski open subset  $U = \mathbb{X}(f) \cap \mathbb{X}(d) \hookrightarrow X$ . For any  $x \in X$  with maximal ideal  $\mathfrak{m}_x \triangleleft \mathbb{C}[X]$  we claim that

$$\frac{A}{A\mathfrak{m}_x} \simeq M_n(\mathbb{C})$$

Indeed, the images of the  $a_i$  give a  $\mathbb{C}$ -basis in the quotient such that the  $n^2 \times n^2$ -matrix of their product-traces is invertible. This property is equivalent to the quotient being  $M_n(\mathbb{C})$ . The corresponding semi-simple representation of  $A$  is simple, proving that  $X_{az}$  is a non-empty Zariski open subset of  $X$ . But then, over  $U$  the restriction of the quotient map

$$\underline{rep}_n^{tr} A \big|_{\pi^{-1}(U)} \longrightarrow U$$

is a principal  $PGL_n$ -fibration. In fact, this restricted quotient map determines an element in  $H_{et}^1(U, PGL_n)$  determining the class of the central simple  $\mathbb{C}(X)$ -algebra  $\Delta$  in  $H_{et}^1(\mathbb{C}(X), PGL_n)$ . Restrict this quotient map further to  $U \cap X_{reg}$ , then the  $PGL_n$ -fibration

$$\underline{rep}_n^{tr} A \big|_{\pi^{-1}(U \cap X_{reg})} \longrightarrow U \cap X_{reg}$$

has a smooth base and therefore also the total space is smooth. But then,  $U \cap X_{reg}$  is a non-empty Zariski open subset of  $Sm_{tr} A$ .  $\square$

Observe that the normality assumption on  $X$  is no restriction as the quotient scheme is locally normal in a point of  $Sm_{tr} A$ . Our next result limits the local dimension vectors  $\alpha_\xi$ .

PROPOSITION 5.13. *Let  $A$  be a Cayley-Hamilton order and  $\xi \in Sm_{tr} A$  such that the normal space to the orbit of the corresponding semi-simple  $n$ -dimensional representation is*

$$N_\xi = rep_{\alpha_\xi} Q_\xi^\bullet$$

*Then,  $\alpha_\xi$  is the dimension vector of a simple representation of  $Q_\xi^\bullet$ .*

PROOF. Let  $V_\xi$  be the semi-simple representation of  $A$  determined by  $\xi$ . Let  $S_\xi$  be the slice variety in  $V_\xi$  then by the slice theorem we have the following diagram of étale  $GL_n$ -equivariant maps

$$\begin{array}{ccc} & GL_n \times^{GL(\alpha_\xi)} S_\xi & \\ \swarrow^{et} & & \searrow^{et} \\ GL_n \times^{GL(\alpha_\xi)} rep_{\alpha_\xi} Q_\xi^\bullet & & rep_n^{tr} A \end{array}$$

linking a neighborhood of  $V_\xi$  with one of  $(\overline{\mathbb{A}_n}, 0)$ . Because  $A$  is an order, every Zariski neighborhood of  $V_\xi$  in  $rep_n^{tr} A$  contains simple  $n$ -dimensional representations, that is, closed  $GL_n$ -orbits with stabilizer subgroup isomorphic to  $\mathbb{C}^*$ . Transporting this property via the  $GL_n$ -equivariant étale maps, every Zariski neighborhood of  $(\overline{\mathbb{A}_n}, 0)$  contains closed  $GL_n$ -orbits with stabilizer  $\mathbb{C}^*$ . By the correspondence of orbits is associated fiber bundles, every Zariski neighborhood of the trivial representation  $0 \in rep_{\alpha_\xi} Q_\xi^\bullet$  contains closed  $GL(\alpha_\xi)$ -orbits with stabilizer subgroup  $\mathbb{C}^*$ . We have seen that closed  $GL(\alpha_\xi)$ -orbits correspond to semi-simple representations of  $Q_\xi^\bullet$ . However, if the stabilizer subgroup of a semi-simple representation is  $\mathbb{C}^*$  this representation must be simple.  $\square$

THEOREM 5.14. *Let  $A$  be a Cayley-Hamilton order of degree  $n$  with center  $\mathbb{C}[X]$ ,  $X$  a normal variety of dimension  $d$ . For  $\xi \in X = iss_n^{tr} A$  with corresponding semi-simple representation*

$$V_\xi = S_1^{\oplus e_1} \oplus \dots \oplus S_k^{\oplus e_k}$$

*and normal space to the orbit  $\mathcal{O}(V_\xi)$  isomorphic to  $rep_{\alpha_\xi} Q_\xi^\bullet$  as  $GL(\alpha_\xi)$ -modules where  $\alpha_\xi = (e_1, \dots, e_k)$ . Then,  $\xi \in Sm_{tr} A$  if and only if the following two conditions are met*

$$\begin{cases} \alpha_\xi & \text{is the dimension vector of a simple representation of } Q_\xi^\bullet, \text{ and} \\ d & = 1 - \chi_Q(\alpha_\xi, \alpha_\xi) - \sum_{i=1}^k m_{ii} \end{cases}$$

*where  $Q$  is the underlying quiver of  $Q_\xi^\bullet$  and  $m_{ii}$  is the number of marked loops in  $Q_\xi^\bullet$  in vertex  $v_i$ .*

PROOF. By the slice theorem we have étale maps

$$rep_{\alpha_\xi} Q_\xi^\bullet / GL(\alpha_\xi) \xleftarrow{et} S_\xi / GL(\alpha_\xi) \xrightarrow{et} iss_n^{tr} A = X$$

connecting a neighborhood of  $\xi \in X$  with one of the trivial semi-simple representation  $\bar{0}$ . By definition of the Euler-form of  $Q$  we have that

$$\chi_Q(\alpha_\xi, \alpha_\xi) = - \sum_{i \neq j} e_i e_j \chi_{ij} + \sum_i e_i^2 (1 - a_{ii} - m_{ii})$$

On the other hand we have

$$\begin{aligned} \dim \operatorname{rep}_\alpha Q_{\alpha_\xi}^\bullet &= \sum_{i \neq j} e_i e_j \chi_{ij} + \sum_i e_i^2 (a_{ii} + m_{ii}) - \sum_i m_{ii} \\ \dim GL(\alpha_\xi) &= \sum_i e_i^2 \end{aligned}$$

As any Zariski open neighborhood of  $\xi$  contains an open set where the quotient map is a  $PGL(\alpha_\xi) = \frac{GL(\alpha_\xi)}{\mathbb{C}^*}$ -fibration we see that the quotient variety  $\operatorname{rep}_{\alpha_\xi} Q_\xi^\bullet$  has dimension equal to

$$\dim \operatorname{rep}_{\alpha_\xi} Q_\xi^\bullet - \dim GL(\alpha_\xi) + 1$$

and plugging in the above information we see that this is equal to  $1 - \chi_Q(\alpha_\xi, \alpha_\xi) - \sum_i m_{ii}$ .  $\square$

EXAMPLE 5.15. The quantum plane.

We will generalize the discussion of example 5.11 to the algebra

$$A = \frac{\mathbb{C}\langle x, y \rangle}{(yx - qxy)}$$

where  $q$  is a primitive  $n$ -th root of unity. Let  $u = x^n$  and  $v = y^n$  then it is easy to see that  $A$  is a free module of rank  $n^2$  over its center  $\mathbb{C}[u, v]$  and is a Cayley-Hamilton algebra of degree  $n$  with the trace determined on the basis

$$\operatorname{tr}(x^i y^j) = \begin{cases} 0 & \text{when either } i \text{ or } j \text{ is not a multiple of } n, \\ nx^i y^j & \text{when } i \text{ and } j \text{ are multiples of } n, \end{cases}$$


Let  $\xi \in \operatorname{iss}_n A = \mathbb{C}^2$  be a point  $(a^n, b)$  with  $a, b \neq 0$ , then  $\xi$  is of representation type  $(1, n)$  as the corresponding (semi)simple representation  $V_\xi$  is determined by (if  $m$  is odd, for even  $n$  we replace  $a$  by  $ia$  and  $b$  by  $-b$ )

$$\rho(x) = \begin{bmatrix} a & & & & \\ & qa & & & \\ & & \ddots & & \\ & & & \ddots & \\ & & & & q^{n-1}a \end{bmatrix} \quad \text{and} \quad \rho(y) = \begin{bmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & & 0 \\ \vdots & \vdots & & \ddots & \\ 0 & 0 & 0 & \dots & 1 \\ b & 0 & 0 & \dots & 0 \end{bmatrix}$$

One computes that  $\operatorname{Ext}_A^1(V_\xi, V_\xi) = \mathbb{C}^2$  where the algebra map  $A \xrightarrow{\phi} M_n(\mathbb{C}[\varepsilon])$  corresponding to  $(\alpha, \beta)$  is given by

$$\begin{cases} \phi(x) = \rho(x) + \varepsilon \alpha \mathbb{1}_n \\ \phi(y) = \rho(y) + \varepsilon \beta \mathbb{1}_n \end{cases}$$

and all these algebra maps are trace preserving. That is,  $\operatorname{Ext}_A^1(V_\xi, V_\xi) = \operatorname{Ext}_A^{\operatorname{tr}}(V_\xi, V_\xi)$  and as the stabilizer subgroup is  $\mathbb{C}^*$  the marked quiver-setting  $(Q_\xi^\bullet, \alpha_\xi)$  is



and  $d = 1 - \chi_Q(\alpha, \alpha) - \sum_i m_{ii}$  as  $2 = 1 - (-1) + 0$ , compatible with the fact that over these points the quotient map is a principal  $PGL_n$ -fibration.

Next, let  $\xi = (a^n, 0)$  with  $a \neq 0$  (or, by a similar argument  $(0, b^n)$  with  $b \neq 0$ ). Then, the representation type of  $\xi$  is  $(1, 1; \dots; 1, 1)$  because

$$V_\xi = S_1 \oplus \dots \oplus S_n$$


where the simple one-dimensional representation  $S_i$  is given by

$$\begin{cases} \rho(x) = q^i a \\ \rho(y) = 0 \end{cases}$$

One verifies that

$$\operatorname{Ext}_A^1(S_i, S_i) = \mathbb{C} \quad \text{and} \quad \operatorname{Ext}_A^1(S_i, S_j) = \delta_{i+1, j} \mathbb{C}$$


and as the stabilizer subgroup is  $\mathbb{C}^* \times \dots \times \mathbb{C}^*$ , the *Ext*-quiver setting is



The algebra map  $A \xrightarrow{\phi} M_n(\mathbb{C}[\varepsilon])$  corresponding to the extension  $(\alpha_1, \beta_1, \dots, \alpha_n, \beta_n) \in Ext_A^1(V_\xi, V_\xi)$  is given by

$$\left\{ \begin{array}{l} \phi(x) = \begin{bmatrix} a + \varepsilon \alpha_1 & & & & \\ & qa + \varepsilon \alpha_2 & & & \\ & & \ddots & & \\ & & & q^{n-1}a + \varepsilon \alpha_n & \\ & & & & \end{bmatrix} \\ \phi(y) = \varepsilon \begin{bmatrix} 0 & \beta_1 & 0 & \dots & 0 \\ 0 & 0 & \beta_2 & & 0 \\ \vdots & \vdots & & \ddots & \vdots \\ 0 & 0 & 0 & & \beta_{n-1} \\ \beta_n & 0 & 0 & \dots & 0 \end{bmatrix} \end{array} \right.$$

The conditions  $tr(x^j) = 0$  for  $1 \leq i < n$  impose  $n - 1$  linear conditions among the  $\alpha_j$ , whence the space of trace preserving extensions  $Ext_A^{tr}(V_\xi, V_\xi)$  corresponds to the quiver setting



The Euler-form of this quiver  $Q^\bullet$  is given by the  $n \times n$  matrix

$$\begin{bmatrix} 0 & -1 & 0 & \dots & 0 \\ & 1 & -1 & & 0 \\ & & & \ddots & \\ & & & & 1 & -1 \\ -1 & & & & & 1 \end{bmatrix}$$

giving the numerical restriction as  $\alpha_\xi = (1, \dots, 1)$

$$1 - \chi_Q(\alpha, \alpha) - \sum_i m_{ii} = 1 - (-1) - 0 = 2 = \dim \underline{iss}_n^{tr} A$$

so  $\xi \in Sm_{tr} A$ . Finally, the only remaining point is  $\xi = (0, 0)$ . This has representation type  $(n, 1)$  as the corresponding semi-simple representation  $V_\xi$  is the trivial one. The stabilizer subgroup is


$GL_n$  and the (trace preserving) extensions are given by

$$\text{Ext}_A^1(V_\xi, V_\xi) = M_n \oplus M_n \quad \text{and} \quad \text{Ext}_A^{\text{tr}}(V_\xi, V_\xi) = M_n^0 \oplus M_n^0$$

determined by the algebra maps  $A \xrightarrow{\phi} M_n(\mathbb{C}[\varepsilon])$  given by

$$\begin{cases} \phi(x) = \varepsilon m_1 \\ \phi(y) = \varepsilon m_2 \end{cases}$$

That is, the relevant quiver setting  $(Q_\xi^\bullet, \alpha_\xi)$  is in this point



This time,  $\xi \notin \text{Sm}_{\text{tr}} A$  as the numerical condition fails

$$1 - \chi_Q(\alpha, \alpha) - \sum_i m_{ii} = 1 - (-n^2) - 0 \neq 2 = \dim \text{iss}_n^{\text{tr}} A$$

unless  $n = 1$ . That is,  $\text{Sm}_{\text{tr}} A = \mathbb{C}^2 - \{(0, 0)\}$ .


### 5.3. Local classification.

If we want to study the local structure of Cayley-Hamilton orders  $A$  of degree  $n$  over a central normal variety  $X$  of dimension  $d$ , we have to compile a list of admissible marked quiver settings, that is couples  $(Q^\bullet, \alpha)$  satisfying the two properties


$$\begin{cases} \alpha \text{ is the dimension vector of a simple representation of } Q^\bullet, \text{ and} \\ d = 1 - \chi_Q(\alpha, \alpha) - \sum_i m_i \end{cases}$$

In this section, we will give the first steps in such a classification project

The basic idea that we use is to shrink a marked quiver-setting to its simplest form and classify these simplest forms for given  $d$ . By *shrinking* we mean the following process. Assume  $\alpha = (e_1, \dots, e_k)$  is the dimension vector of a simple representation of  $Q^\bullet$  and let  $v_i$  and  $v_j$  be two vertices connected with an arrow such that  $e_i = e_j = e$ . That is, locally we have the following situation



We will use one of the arrows connecting  $v_i$  with  $v_j$  to identify the two vertices. That is, we form the shrunk marked quiver-setting  $(Q_s^\bullet, \alpha_s)$  where  $Q_s^\bullet$  is the marked quiver on  $k - 1$  vertices  $\{v_1, \dots, \hat{v}_i, \dots, v_k\}$  and  $\alpha_s$  is the dimension vector with  $e_i$  removed.  $Q_s^\bullet$  has the following form in a neighborhood of the contracted vertex



In  $Q_s^\bullet$  we have for all  $k, l \neq i, j$  that  $\chi_{kl}^s = \chi_{kl}$ ,  $a_{kk}^s = a_{kk}$ ,  $m_{kk}^s = m_{kk}$  and the number of arrows and (marked) loops connected to  $v_j$  are determined as follows

- $\chi_{jk}^s = \chi_{ik} + \chi_{jk}$



- $\chi_{kj}^s = \chi_{ki} + \chi_{kj}$
- $a_{jj}^s = a_{ii} + a_{jj} + \chi_{ij} + \chi_{ji} - 1$
- $m_{jj}^s = m_{ii} + m_{jj}$

LEMMA 5.16.  $\alpha$  is the dimension vector of a simple representation of  $Q^\bullet$  if and only if  $\alpha_s$  is the dimension vector of a simple representation of  $Q_s^\bullet$ . Moreover,

$$\dim \operatorname{rep}_\alpha Q^\bullet / GL(\alpha) = \dim \operatorname{rep}_{\alpha_s} Q_s^\bullet / GL(\alpha_s)$$

PROOF. Fix an arrow  $\textcircled{i} \xleftarrow{a} \textcircled{j}$ . As  $e_i = e_j = e$  there is a Zariski open subset  $U \hookrightarrow \operatorname{rep}_\alpha Q^\bullet$  of points  $V$  such that  $V_a$  is invertible. By basechange in either  $v_i$  or  $v_j$  we can find a point  $W$  in its orbit such that  $W_a = \mathbb{1}_e$ . If we think of  $W_a$  as identifying  $\mathbb{C}^{e_i}$  with  $\mathbb{C}^{e_j}$  we can view the remaining maps of  $W$  as a representation in  $\operatorname{rep}_{\alpha_s} Q_s^\bullet$  and denote it by  $W^s$ . The map  $U \rightarrow \operatorname{rep}_{\alpha_s} Q_s^\bullet$  is well-defined and maps  $GL(\alpha)$ -orbits to  $GL(\alpha_s)$ -orbits. Conversely, given a representation  $W' \in \operatorname{rep}_{\alpha_s} Q_s^\bullet$  we can uniquely determine a representation  $W \in U$  mapping to  $W'$ . Both claims follow immediately from this observation.  $\square$

A marked quiver-setting can uniquely be shrunk to its *simplified form*, which has the characteristic property that no arrow-connected vertices can have the same dimension. The shrinking process has a converse operation which we will call *splitting of a vertex*. However, this splitting operation is usually not uniquely determined.

Before compiling a lists of marked-quiver settings in simplified form for a specific base-dimension  $d$ , we bound the components of  $\alpha$ .

PROPOSITION 5.17. Let  $\alpha = (e_1, \dots, e_k)$  be the dimension vector of a simple representation of  $Q$  and let  $1 - \chi_Q(\alpha, \alpha) = d = \dim \operatorname{rep}_\alpha Q / GL(\alpha)$ . Then, if  $e = \max e_i$ , we have that  $d \geq e + 1$ .


PROOF. By lemma 5.16 we may assume that  $(Q, \alpha)$  is brought in its simplified form, that is, no two arrow-connected vertices have the same dimension. Let  $\chi_{ii}$  denote the number of loops in a vertex  $v_i$ , then

$$-\chi_Q(\alpha, \alpha) = \begin{cases} \sum_i e_i (\sum_j \chi_{ij} e_j - e_i) \\ \sum_i e_i (\sum_j \chi_{ji} e_j - e_i) \end{cases}$$

and observe that the bracketed terms are positive by the requirement that  $\alpha$  is the dimension vector of a simple representation. We call them the incoming  $in_i$ , respectively outgoing  $out_i$ , contribution of the vertex  $v_i$  to  $d$ . Let  $v_m$  be a vertex with maximal vertex-dimension  $e$ .

$$in_m = e \left( \sum_{j \neq m} \chi_{jm} e_j + (\chi_{ii} - 1) e \right) \quad \text{and} \quad out_m = e \left( \sum_{j \neq m} \chi_{ij} e_j + (\chi_{ii} - 1) e \right)$$

If there are loops in  $v_m$ , then  $in_m \geq 2$  or  $out_m \geq 2$  unless the local structure of  $Q$  is



in which case  $in_m = e = out_m$ . Let  $v_i$  be the unique incoming vertex of  $v_m$ , then we have  $out_i \geq e - 1$ . But then,

$$d = 1 - \chi_Q(\alpha, \alpha) = 1 + \sum_j out_j \geq 2e$$

If  $v_m$  has no loops, consider the incoming vertices  $\{v_{i_1}, \dots, v_{i_s}\}$ , then


$$in_m = e\left(\sum_{j=1}^s \chi_{i_j m} e_{i_j} - e\right)$$

which is  $\geq e$  unless  $\sum \chi_{i_j m} e_{i_j} = e$ , but in that case we have

$$\sum_{j=1}^s out_{i_j} \geq e^2 - \sum_{j=1}^s e_{i_j}^2 \geq e$$

the last inequality because all  $e_{i_j} < e$ . In either case we have that  $d = 1 - \chi_Q(\alpha, \alpha) = 1 + \sum_i out_i = 1 + \sum_i in_i \geq e + 1$ .  $\square$

EXAMPLE 5.18. In a list of simplified marked quivers we are only interested in  $rep_\alpha Q^\bullet$  as  $GL(\alpha)$ -module and we call two setting *equivalent* if they determine the same  $GL(\alpha)$ -module. For example, the marked quiver-settings



determine the same  $\mathbb{C}^* \times GL_2$ -module, hence are equivalent.

THEOREM 5.19. *Let  $A$  be a Cayley-Hamilton order of degree  $n$  over a central normal variety  $X$  of degree  $d$ . Then, the local quiver of  $A$  in a point  $\xi \in X = iss_n^{tr} A$  belonging to the smooth locus  $Sm_{tr} A$  can be shrunk to one of a finite list of equivalence classes of simplified marked quiver-settings. For  $d \leq 4$ , the complete list is given in figure 1 where the boxed value is the dimension  $d$  of  $X$ .*

An immediate consequence is a noncommutative analogon of the fact that commutative smooth varieties have only one type of analytic (or étale) local behavior.

THEOREM 5.20. *There are only finitely many types of étale local behaviour of smooth Cayley-Hamilton orders of degree  $n$  over a central variety of dimension  $d$ .*

PROOF. The foregoing reduction shows that for fixed  $d$  there are only a finite number of marked quiver-settings shrunk to their simplified form. As  $\sum e_i \leq n$ , we can apply the splitting operations on vertices only a finite number of times.  $\square$

### 5.4. Low dimensional orders.

W. Schelter has proved in [68] that in dimension one, Cayley-smooth orders are hereditary. We give an alternative proof of this result using the étale local classification. The next result follows also by splitting the dimension 1 case in figure 1. We give a direct proof illustrating the type-stratification result of section 5.1.

THEOREM 5.21. *Let  $A$  be a Cayley-Hamilton order of degree  $n$  over an affine curve  $X = iss_n^{tr} A$ . If  $\xi \in Sm_{tr} A$ , then the étale local structure of  $A$  in  $\xi$  is determined by a marked quiver-setting which is an oriented cycle on  $k$  vertices with  $k \leq n$  and an unordered partition  $p = (d_1, \dots, d_k)$  having precisely  $k$  parts such that  $\sum_i d_i = n$  determining the dimensions of the simple components of  $V_\xi$ , see figure 2.*

PROOF. Let  $(Q^\bullet, \alpha)$  be the corresponding local marked quiver-setting. Because  $Q^\bullet$  is strongly connected, there exist oriented cycles in  $Q^\bullet$ . Fix one such cycle of length  $s \leq k$  and renumber the vertices of  $Q^\bullet$  such that the first  $s$  vertices make




FIGURE 1. The simplified local quivers for  $d \leq 4$




FIGURE 2. Cayley-smooth curve types.

up the cycle. If  $\alpha = (e_1, \dots, e_k)$ , then there exist semi-simple representations in  $rep_\alpha Q^\bullet$  with composition

$$\alpha_1 = (\underbrace{1, \dots, 1}_s, \underbrace{0, \dots, 0}_{k-s}) \oplus \epsilon_1^{\oplus e_1 - 1} \oplus \dots \oplus \epsilon_s^{\oplus e_s - 1} \oplus \epsilon_{s+1}^{\oplus e_{s+1}} \oplus \dots \oplus \epsilon_k^{\oplus e_k}$$

where  $\epsilon_i$  stands for the simple one-dimensional representation concentrated in vertex  $v_i$ . There is a one-dimensional family of simple representations of dimension

vector  $\alpha_1$ , hence the stratum of semi-simple representations in  $iss_\alpha Q^\bullet$  of representation type  $\tau = (1, \alpha_1; e_1 - 1, \epsilon_1; \dots; e_s - 1, \epsilon_s; e_{s+1}, \epsilon_{s+1}; e_k, \epsilon_k)$  is at least one-dimensional. However, as  $\dim iss_\alpha Q^\bullet = 1$  this can only happen if this semi-simple representation is actually simple. That is, when  $\alpha = \alpha_1$  and  $k = s$ .  $\square$


If  $V_\xi$  is the semi-simple  $n$ -dimensional representation of  $A$  corresponding to  $\xi$ , then

$$V_\xi = S_1 \oplus \dots \oplus S_k \quad \text{with} \quad \dim S_i = d_i$$

and the stabilizer subgroup is  $GL(\alpha) = \mathbb{C}^* \times \dots \times \mathbb{C}^*$  embedded in  $GL_n$  via the diagonal embedding

$$(\lambda_1, \dots, \lambda_k) \longrightarrow \text{diag}(\underbrace{\lambda_1, \dots, \lambda_1}_{d_1}, \dots, \underbrace{\lambda_k, \dots, \lambda_k}_{d_k})$$

Further, using basechange in  $rep_\alpha Q^\bullet$  we can bring every simple  $\alpha$ -dimensional representation of  $Q^\alpha$  in standard form



where  $x \in \mathbb{C}^*$  is the arrow from  $v_k$  to  $v_1$ . That is,  $\mathbb{C}[rep_\alpha Q^\bullet]^{GL(\alpha)} \simeq \mathbb{C}[x]$  proving that the quotient (or central) variety  $X$  must be smooth in  $\xi$  by the slice result. Moreover, as  $\widehat{A}_\xi \simeq \widehat{\mathbb{T}}_\alpha$  we have, using the numbering conventions of the vertices) the following block decomposition

$$\widehat{A}_\xi \simeq \begin{bmatrix} M_{d_1}(\mathbb{C}[[x]]) & M_{d_1 \times d_2}(\mathbb{C}[[x]]) & \dots & M_{d_1 \times d_k}(\mathbb{C}[[x]]) \\ M_{d_2 \times d_1}(x\mathbb{C}[[x]]) & M_{d_2}(\mathbb{C}[[x]]) & \dots & M_{d_2 \times d_k}(\mathbb{C}[[x]]) \\ \vdots & \vdots & \ddots & \vdots \\ M_{d_k \times d_1}(x\mathbb{C}[[x]]) & M_{d_k \times d_2}(x\mathbb{C}[[x]]) & \dots & M_{d_k}(\mathbb{C}[[x]]) \end{bmatrix}$$

From the local description of hereditary orders given in [67, Thm. 39.14] we deduce that  $A_\xi$  is an hereditary order. That is, we have the following characterization of the smooth locus

**PROPOSITION 5.22.** *Let  $A$  be a Cayley-Hamilton order of degree  $n$  over a central affine curve  $X$ . Then,  $Sm_{tr} A$  is the locus of points  $\xi \in X$  such that  $A_\xi$  is an hereditary order (in particular,  $\xi$  must be a smooth point of  $X$ ).*

**THEOREM 5.23.** *Let  $A$  be a Cayley-Hamilton central  $\mathcal{O}_X$ -order of degree  $n$  where  $X$  is a projective curve. Equivalent are*




FIGURE 3. Cayley-smooth surface types.

- (1)  $\mathcal{A}$  is a sheaf of Cayley-smooth orders
- (2)  $X$  is smooth and  $\mathcal{A}$  is a sheaf of hereditary  $\mathcal{O}_X$ -orders

We now turn to orders over surfaces. The next result can equally be proved using splitting and the classification of figure 1.

**THEOREM 5.24.** *Let  $A$  be a Cayley-Hamilton order of degree  $n$  over an affine surface  $X = \text{iss}_n^{tr} A$ . If  $\xi \in \text{Sm}_{tr} A$ , then the étale local structure of  $A$  in  $\xi$  is determined by a marked local quiver-setting  $A_{klm}$  on  $k + l + m \leq n$  vertices and an unordered partition  $p = (d_1, \dots, d_{k+l+m})$  of  $n$  with  $k + l + m$  non-zero parts determined by the dimensions of the simple components of  $V_\xi$  as in figure 3.*

**PROOF.** Let  $(Q^\bullet, \alpha)$  be the marked quiver-setting on  $r$  vertices with  $\alpha = (e_1, \dots, e_r)$  corresponding to  $\xi$ . As  $Q^\bullet$  is strongly connected and the quotient variety is two-dimensional,  $Q^\bullet$  must contain more than one oriented cycle, hence it contains a sub-quiver of type  $A_{klm}$ , possibly degenerated with  $k$  or  $l$  equal to zero. Order the first  $k + l + m$  vertices of  $Q^\bullet$  as indicated. One verifies that  $A_{klm}$  has simple representations of dimension vector  $(1, \dots, 1)$ . Assume that  $A_{klm}$  is a proper subquiver and denote  $s = k + l + m + 1$  then  $Q^\bullet$  has semi-simple representations in  $\text{rep}_\alpha Q^\bullet$  with dimension-vector decomposition

$$\alpha_1 = (\underbrace{1, \dots, 1}_{k+l+m}, 0, \dots, 0) \oplus \epsilon_1^{\oplus e_1 - 1} \oplus \dots \oplus \epsilon_{k+l+m}^{\oplus e_{k+l+m} - 1} \oplus \epsilon_s^{\oplus e_s} \oplus \dots \oplus \epsilon_r^{\oplus e_r}$$

Applying the formula for the dimension of the quotient variety shows that  $iss_{(1,\dots,1)} A_{klm}$  has dimension 2 so there is a two-dimensional family of such semi-simple representation in the two-dimensional quotient variety  $iss_\alpha Q^\bullet$ . This is only possible if this semi-simple representation is actually simple, whence  $r = k + l + m$ ,  $Q^\bullet = A_{klm}$  and  $\alpha = (1, \dots, 1)$ .  $\square$


If  $V_\xi$  is the semi-simple  $n$ -dimensional representation of  $A$  corresponding to  $\xi$ , then

$$V_\xi = S_1 \oplus \dots \oplus S_r \quad \text{with} \quad \dim S_i = d_i$$

and the stabilizer subgroup  $GL(\alpha) = \mathbb{C}^* \times \dots \times \mathbb{C}^*$  embedded diagonally in  $GL_n$

$$(\lambda_1, \dots, \lambda_r) \mapsto \text{diag}(\underbrace{\lambda_1, \dots, \lambda_1}_{d_1}, \dots, \underbrace{\lambda_r, \dots, \lambda_r}_{d_r})$$

By basechange in  $rep_\alpha A_{klm}$  we can bring every simple  $\alpha$ -dimensional representation in the following standard form



with  $x, y \in \mathbb{C}^*$  and as  $\mathbb{C}[iss_\alpha A_{klm}] = \mathbb{C}[rep_\alpha A_{klm}]^{GL(\alpha)}$  is the ring generated by traces along oriented cycles in  $A_{klm}$ , it is isomorphic to  $\mathbb{C}[x, y]$ . From the slice result one deduces that  $\xi$  must be a smooth point of  $X$  and because  $\widehat{A}_\xi \simeq \widehat{\mathbb{T}}_\alpha$  we deduce it must have the following block-decomposition

$$\widehat{A}_\xi \simeq \begin{array}{|c|c|c|} \hline & (1) & \\ \hline (x) & & (y) & (1) \\ \hline & & (1) & \\ \hline (x) & & (y) & (1) \\ \hline & & & (1) \\ \hline (x) & (y) & & (x, y) \\ \hline \end{array} \hookrightarrow M_n(\mathbb{C}[[x, y]])$$

$\underbrace{\hspace{1.5cm}}_k \quad \underbrace{\hspace{1.5cm}}_l \quad \underbrace{\hspace{1.5cm}}_m$

where at spot  $(i, j)$  with  $1 \leq i, j \leq k + l + m$  there is a block of dimension  $d_i \times d_j$  with entries the indicated ideal of  $\mathbb{C}[[x, y]]$ .




FIGURE 4. Proper semi-simples of  $A_{klm}$ .

DEFINITION 5.25. Let  $A$  be a Cayley-Hamilton central  $\mathbb{C}[X]$ -order of degree  $n$  in a central simple  $\mathbb{C}(X)$ - algebra  $\Delta$  of dimension  $n^2$ .

- (1)  $A$  is said to be *étale locally split* in  $\xi$  if and only if  $\widehat{A}_\xi$  is a central  $\hat{\mathcal{O}}_{X,x}$ -order in  $M_n(\hat{\mathcal{O}}_{X,x} \otimes_{\mathcal{O}_{X,x}} \mathbb{C}(X))$ .
- (2) The *ramification locus*  $\text{ram}_A$  of  $A$  is the locus of points  $\xi \in X$  such that

$$\frac{A}{\mathfrak{m}_\xi \text{Am}_\xi} \not\cong M_n(\mathbb{C})$$

The complement  $X - \text{ram}_A$  is the Azumaya locus  $X_{az}$  of  $A$ .

THEOREM 5.26. *Let  $\mathcal{A}$  be a Cayley-smooth central  $\mathcal{O}_X$ -order of degree  $n$  over a projective surface  $X$ . Then,*

- (1)  $X$  is smooth.
- (2)  $\mathcal{A}$  is étale locally split in all points of  $X$ .
- (3) The ramification divisor  $\text{ram}_{\mathcal{A}} \hookrightarrow X$  is either empty or consists of a finite number of isolated (possibly embedded) points and a reduced divisor having as its worst singularities normal crossings.

PROOF. (1) and (2) follow from the above local description of  $\mathcal{A}$ . As for (3) we have to compute the local quiver-settings in proper semi-simple representations of  $\text{rep}_\alpha A_{klm}$ . As simples have a strongly connected support, the decomposition types of these proper semi-simples are depicted in figure 4 with  $x, y \in \mathbb{C}^*$ . By the description of local quivers given in section 3 we see that they are respectively of the forms in figure 5. The associated unordered partitions are defined in the obvious way, that is, to the looped vertex one assigns the sum of the  $d_i$  belonging to the loop-contracted circuit and the other components of the partition are preserved. Using the étale local isomorphism between  $X$  in a neighborhood of  $\xi$  and of  $\text{iss}_\alpha A_{klm}$  in a neighborhood of the trivial representation, we see that the local picture of quiver-settings of  $\mathcal{A}$  in a neighborhood of  $\xi$  is described in figure 6 The Azumaya points are the points in which the quiver-setting is  $A_{001}$  (the two-loop quiver). From this local description the result follows if we take care of possibly degenerated cases.  $\square$




FIGURE 5. Local quivers for  $A_{klm}$ .





FIGURE 6. Local picture for  $A_{klm}$ .

EXAMPLE 5.27. An isolated point in  $\xi$  can occur if the quiver-setting in  $\xi$  is of type  $A_{00m}$  with  $m \geq 2$ , that is,



In the case of curves and surfaces, the central variety  $X$  of a Cayley-smooth model  $\mathcal{A}$  had to be smooth and that  $\mathcal{A}$  is étale locally split in every point  $\xi \in X$ . Both of these properties are no longer valid in higher dimensions.

LEMMA 5.28. For dimension  $d \geq 3$ , the center  $Z$  of a Cayley-smooth order of degree  $n$  can have singularities.







FIGURE 7. Central singularities can arise.

PROOF. Consider the marked quiver-setting of figure 7 which is allowed for dimension  $d = 3$  and degree  $n = 2$ . The quiver-invariants are generated by the traces along oriented cycles, that is by  $ac, ad, bc$  and  $bd$ . The coordinate ring is

$$\mathbb{C}[iss_\alpha Q] \simeq \frac{\mathbb{C}[x, y, z, v]}{(xv - yz)}$$

having a singularity in the origin. This example can be extended to dimensions  $d \geq 3$  by adding loops in one of the vertices.



□

LEMMA 5.29. For dimension  $d \geq 3$ , a Cayley-smooth algebra does not have to be locally étale split in every point of its central variety.


PROOF. Consider the following allowable quiver-setting for  $d = 3$  and  $n = 2$



The corresponding Cayley-smooth algebra  $A$  is generated by two generic  $2 \times 2$  trace zero matrices, say  $A$  and  $B$ . From the description of the trace algebra  $\mathbb{T}_2^2$  we see that its center is generated by  $A^2 = x, B^2 = z$  and  $AB + BA = y$ . Alternatively, we can identify  $A$  with the Clifford-algebra over  $\mathbb{C}[x, y, z]$  of the non-degenerate quadratic form

$$\begin{bmatrix} x & y \\ y & z \end{bmatrix}$$

This is a noncommutative domain and remains to be so over the formal power series  $\mathbb{C}[[x, y, z]]$ . That is,  $A$  cannot be split by an étale extension in the origin. More generally, whenever the local marked quiver contains vertices with dimension  $\geq 2$ , the corresponding Cayley-smooth algebra cannot be split by an étale extension as the local quiver-setting does not change and for a split algebra all vertex-dimensions have to be equal to 1. In particular, the Cayley-smooth algebra of degree 2 corresponding to the quiver-setting



cannot be split by an étale extension in the origin. Its corresponding dimension is

$$d = 3k + 4l - 3$$

whenever  $k + l \geq 2$  and all dimensions  $d \geq 3$  are obtained. □

**5.5. Noncommutative smooth surfaces.**

Let  $X$  be a projective surface, in this section we will characterize the central simple  $\mathbb{C}(X)$ -algebras  $\Delta$  allowing a *Cayley-smooth model*. We first need to perform a local calculation. Consider the ring of algebraic functions in two variables  $\mathbb{C}\{x, y\}$  and let  $X_{loc} = Spec \mathbb{C}\{x, y\}$ . There is only one codimension two subvariety :  $m = (x, y)$ . Let us compute the coniveau spectral sequence for  $X_{loc}$ . If  $K$  is the field of fractions of  $\mathbb{C}\{x, y\}$  and if we denote with  $k_p$  the field of fractions of  $\mathbb{C}\{x, y\}/p$  where  $p$  is a height one prime, we have as its first term

0	0	0	0	...
$H^2(K, \mu_n)$	$\oplus_p H^1(k_p, \mathbb{Z}_n)$	$\mu_n^{-1}$	0	...
$H^1(K, \mu_n)$	$\oplus_p \mathbb{Z}_n$	0	0	...
$\mu_n$	0	0	0	...

Because  $\mathbb{C}\{x, y\}$  is a unique factorization domain, we see that the map

$$H_{et}^1(K, \mu_n) = K^*/(K^*)^n \xrightarrow{\gamma} \oplus_p \mathbb{Z}_n$$

is surjective. Moreover, all fields  $k_p$  are isomorphic to the field of fractions of  $\mathbb{C}\{z\}$  whose only cyclic extensions are given by adjoining a root of  $z$  and hence they are all ramified in  $m$ . Therefore, the component maps

$$\mathbb{Z}_n = H_{et}^1(k_p, \mathbb{Z}_n) \xrightarrow{\beta_L} \mu^{-1}$$

are isomorphisms. But then, the second (and limiting) term of the spectral sequence has the form

0	0	0	0	...
$Ker \alpha$	$Ker \beta / Im \alpha$	0	0	...
$Ker \gamma$	0	0	0	...
$\mu_n$	0	0	0	...

Finally, we use the fact that  $\mathbb{C}\{x, y\}$  is strict Henselian whence has no proper étale extensions. But then,

$$H_{et}^i(X_{loc}, \mu_n) = 0 \text{ for } i \geq 1$$

and substituting this information in the spectral sequence we obtain that the top sequence of the coniveau spectral sequence

$$0 \longrightarrow Br_n K \xrightarrow{\alpha} \oplus_p \mathbb{Z}_n \longrightarrow \mathbb{Z}_n \longrightarrow 0$$

is exact. From this sequence we immediately obtain the following

LEMMA 5.30. *With notations as before, we have*

- (1) *Let  $U = X_{loc} - V(x)$ , then  $Br_n U = 0$*
- (2) *Let  $U = X_{loc} - V(xy)$ , then  $Br_n U = \mathbb{Z}_n$  with generator the quantum-plane algebra*

$$\mathbb{C}_\zeta[u, v] = \frac{\mathbb{C}\langle u, v \rangle}{(vu - \zeta uv)}$$

where  $\zeta$  is a primitive  $n$ -th root of one

Let  $\Delta$  be a central simple algebra of dimension  $n^2$  over a field  $L$  of transcendence degree 2. We want to determine when  $\Delta$  admits a *Cayley-smooth model*  $\mathcal{A}$ , that is, a sheaf of Cayley-smooth  $\mathcal{O}_X$ -algebras where  $X$  is a projective surface with functionfield  $\mathbb{C}(X) = L$ . It follows from theorem 5.26 that, if such a model exists,  $X$  must be a smooth projective surface. We may assume that  $X$  is a (commutative) smooth model for  $L$ . By the Artin-Mumford exact sequence 3.62 the class of  $\Delta$  in  $Br_n \mathbb{C}(X)$  is determined by the following geo-combinatorial data

- A finite collection  $\mathcal{C} = \{C_1, \dots, C_k\}$  of *irreducible curves* in  $X$ .
- A finite collection  $\mathcal{P} = \{P_1, \dots, P_l\}$  of *points* of  $X$  where each  $P_i$  is either an intersection point of two or more  $C_i$  or a singular point of some  $C_i$ .
- For each  $P \in \mathcal{P}$  the *branch-data*  $b_P = (b_1, \dots, b_{i_P})$  with  $b_i \in \mathbb{Z}_n = \mathbb{Z}/n\mathbb{Z}$  and  $\{1, \dots, i_P\}$  the different branches of  $\mathcal{C}$  in  $P$ . These numbers must satisfy the admissibility condition

$$\sum_i b_i = 0 \in \mathbb{Z}_n$$

for every  $P \in \mathcal{P}$

- for each  $C \in \mathcal{C}$  we fix a cyclic  $\mathbb{Z}_n$ -cover of smooth curves

$$D \longrightarrow \tilde{C}$$

of the desingularization  $\tilde{C}$  of  $C$  which is compatible with the branch-data.

If  $\mathcal{A}$  is a maximal  $\mathcal{O}_X$ -order in  $\Delta$ , then the ramification locus  $ram_{\mathcal{A}}$  coincides with the collection of curves  $\mathcal{C}$ . We fix such a maximal  $\mathcal{O}_X$ -order  $\mathcal{A}$  and investigate its Cayley-smooth locus.

PROPOSITION 5.31. *Let  $\mathcal{A}$  be a maximal  $\mathcal{O}_X$ -order in  $\Delta$  with  $X$  a projective smooth surface and with geo-combinatorial data  $(\mathcal{C}, \mathcal{P}, b, \mathcal{D})$  determining the class of  $\Delta$  in  $Br_n \mathbb{C}(X)$ .*

*If  $\xi \in X$  lies in  $X - \mathcal{C}$  or if  $\xi$  is a non-singular point of  $\mathcal{C}$ , then  $\mathcal{A}$  is Cayley-smooth in  $\xi$ .*

PROOF. If  $\xi \notin \mathcal{C}$ , then  $\mathcal{A}_\xi$  is an Azumaya algebra over  $\mathcal{O}_{X,x}$ . As  $X$  is smooth in  $\xi$ ,  $\mathcal{A}$  is Cayley-smooth in  $\xi$ . Alternatively, we know that Azumaya algebras are split by étale extensions, whence  $\hat{\mathcal{A}}_\xi \simeq M_n(\mathbb{C}[[x, y]])$  which shows that the behaviour of  $\mathcal{A}$  near  $\xi$  is controlled by the local data




and hence  $\xi \in Sm_{tr} \mathcal{A}$ . Next, assume that  $\xi$  is a nonsingular point of the ramification divisor  $\mathcal{C}$ . Consider the pointed spectrum  $X_\xi = \text{Spec } \mathcal{O}_{X,\xi} - \{\mathfrak{m}_\xi\}$ . The only prime ideals are of height one, corresponding to the curves on  $X$  passing through  $\xi$  and hence this pointed spectrum is a Dedekind scheme. Further,  $\mathcal{A}$  determines a maximal order over  $X_\xi$ . But then, tensoring  $\mathcal{A}$  with the strict henselization  $\mathcal{O}_{X,\xi}^{sh} \simeq \mathbb{C}\{x, y\}$  determines a sheaf of hereditary orders on the pointed spectrum  $\hat{X}_\xi = \text{Spec } \mathbb{C}\{x, y\} - \{(x, y)\}$  and we may choose the local variable  $x$  such that  $x$  is a local parameter of the ramification divisor  $\mathcal{C}$  near  $\xi$ .

Using the characterization result for hereditary orders over discrete valuation rings, given in [67, Thm. 39.14] we know the structure of this extended sheaf of hereditary orders over any height one prime of  $\hat{X}_\xi$ . Because  $\mathcal{A}_\xi$  is a reflexive (even a projective)  $\mathcal{O}_{X,\xi}$ -module, this height one information determines  $\mathcal{A}_\xi^{sh}$  or  $\hat{\mathcal{A}}_\xi$ . This proves that  $\mathcal{A}_\xi^{sh}$  must be isomorphic to the following blockdecomposition

$$\begin{bmatrix} M_{d_1}(\mathbb{C}\{x, y\}) & M_{d_1 \times d_2}(\mathbb{C}\{x, y\}) & \dots & M_{d_1 \times d_k}(\mathbb{C}\{x, y\}) \\ M_{d_2 \times d_1}(x\mathbb{C}\{x, y\}) & M_{d_2}(\mathbb{C}\{x, y\}) & \dots & M_{d_2 \times d_k}(\mathbb{C}\{x, y\}) \\ \vdots & \vdots & \ddots & \vdots \\ M_{d_k \times d_1}(x\mathbb{C}\{x, y\}) & M_{d_k \times d_2}(x\mathbb{C}\{x, y\}) & \dots & M_{d_k}(\mathbb{C}\{x, y\}) \end{bmatrix}$$

for a certain partition  $p = (d_1, \dots, d_k)$  of  $n$  having  $k$  parts. In fact, as we started out with a maximal order  $\mathcal{A}$  one can even show that all these integers  $d_i$  must be equal. This local form corresponds to the following quiver-setting



whence  $\xi \in Sm_{tr} A$  as this is one of the allowed surface settings.  $\square$

A maximal  $\mathcal{O}_X$ -order in  $\Delta$  can have at worst noncommutative singularities in the singular points of the ramification divisor  $\mathcal{C}$ . Theorem 5.26 a Cayley-smooth order over a surface has as ramification-singularities at worst normal crossings. We are always able to reduce to normal crossings by the following classical result on commutative surfaces, see for example [28, V.3.8].

**THEOREM 5.32** (Embedded resolution of curves in surfaces). *Let  $\mathcal{C}$  be any curve on the surface  $X$ . Then, there exists a finite sequence of blow-ups*

$$X' = X_s \longrightarrow X_{s-1} \longrightarrow \dots \longrightarrow X_0 = X$$

and, if  $f : X' \longrightarrow X$  is their composition, then the total inverse image  $f^{-1}(\mathcal{C})$  is a divisor with normal crossings.

Fix a series of blow-ups  $X' \xrightarrow{f} X$  such that the inverse image  $f^{-1}(\mathcal{C})$  is a divisor on  $X'$  having as worst singularities normal crossings. We will replace the Cayley-Hamilton  $\mathcal{O}_X$ -order  $\mathcal{A}$  by a Cayley-Hamilton  $\mathcal{O}_{X'}$ -order  $\mathcal{A}'$  where  $\mathcal{A}'$  is a sheaf of  $\mathcal{O}_{X'}$ -maximal orders in  $\Delta$ . In order to determine the ramification divisor of  $\mathcal{A}'$  we need to be able to keep track how the ramification divisor  $\mathcal{C}$  of  $\Delta$  changes if we blow up a singular point  $p \in \mathcal{P}$ .

**LEMMA 5.33.** *Let  $\tilde{X} \longrightarrow X$  be the blow-up of  $X$  at a singular point  $p$  of  $\mathcal{C}$ , the ramification divisor of  $\Delta$  on  $X$ . Let  $\tilde{\mathcal{C}}$  be the strict transform of  $\mathcal{C}$  and  $E$  the exceptional line on  $\tilde{X}$ . Let  $\mathcal{C}'$  be the ramification divisor of  $\Delta$  on the smooth model  $\tilde{X}$ . Then,*

- (1) *Assume the local branch data at  $p$  distribute in an admissible way on  $\tilde{\mathcal{C}}$ , that is,*


$$\sum_{i \text{ at } q} b_{i,p} = 0 \text{ for all } q \in E \cap \tilde{\mathcal{C}}$$

*where the sum is taken only over the branches at  $q$ . Then,  $\mathcal{C}' = \tilde{\mathcal{C}}$ .*

- (2) *Assume the local branch data at  $p$  do not distribute in an admissible way, then  $\mathcal{C}' = \tilde{\mathcal{C}} \cup E$ .*

**PROOF.** Clearly,  $\tilde{\mathcal{C}} \subset \mathcal{C}' \subset \tilde{\mathcal{C}} \cup E$ . By the Artin-Mumford sequence applied to  $X'$  we know that the branch data of  $\mathcal{C}'$  must add up to zero at all points  $q$  of  $\tilde{\mathcal{C}} \cap E$ . We investigate the two cases

1. : Assume  $E \subset \mathcal{C}'$ . Then, the  $E$ -branch number at  $q$  must be zero for all  $q \in \tilde{\mathcal{C}} \cap E$ . But there are no non-trivial étale covers of  $\mathbb{P}^1 = E$  so  $ram(\Delta)$  gives the trivial element in  $H_{et}^1(\mathbb{C}(E), \mu_n)$ , a contradiction. Hence  $\mathcal{C}' = \tilde{\mathcal{C}}$ .



2. : If at some  $q \in \tilde{\mathcal{C}} \cap E$  the branch numbers do not add up to zero, the only remedy is to include  $E$  in the ramification divisor and let the  $E$ -branch number be such that the total sum is zero in  $\mathbb{Z}_n$ .  $\square$

**THEOREM 5.34.** *Let  $\Delta$  be a central simple algebra of dimension  $n^2$  over a field  $L$  of transcendence degree two. Then, there exists a smooth projective surface  $S$  with functionfield  $\mathbb{C}(S) = L$  such that any maximal  $\mathcal{O}_S$ -order  $\mathcal{A}_S$  in  $\Delta$  has at worst a finite number of isolated noncommutative singularities. Each of these singularities is locally étale of quantum-plane type.*

**PROOF.** We take any projective smooth surface  $X$  with functionfield  $\mathbb{C}(X) = L$ . By the Artin-Mumford exact sequence, the class of  $\Delta$  determines a geo-combinatorial set of data

$$(\mathcal{C}, \mathcal{P}, b, \mathcal{D})$$

as before. In particular,  $\mathcal{C}$  is the ramification divisor  $\text{ram}(\Delta)$  and  $\mathcal{P}$  is the set of singular points of  $\mathcal{C}$ . We can separate  $\mathcal{P}$  in two subsets

- $\mathcal{P}_{unr} = \{P \in \mathcal{P} \text{ where all the branch-data } b_P = (b_1, \dots, b_{i_P}) \text{ are trivial, that is, all } b_i = 0 \text{ in } \mathbb{Z}_n\}$
- $\mathcal{P}_{ram} = \{P \in \mathcal{P} \text{ where some of the branch-data } b_P = (b_1, \dots, b_{i_P}) \text{ are non-trivial, that is, some } b_i \neq 0 \text{ in } \mathbb{Z}_n\}$

After a finite number of blow-ups we get a birational morphism  $S_1 \xrightarrow{\pi} X$  such that  $\pi^{-1}(\mathcal{C})$  has as its worst singularities normal crossings and all branches in points of  $\mathcal{P}$  are separated in  $S$ . Let  $\mathcal{C}_1$  be the ramification divisor of  $\Delta$  in  $S_1$ . By the foregoing argument we have

- If  $P \in \mathcal{P}_{unr}$ , then we have that  $\mathcal{C}' \cap \pi^{-1}(P)$  consists of smooth points of  $\mathcal{C}_1$ ,
- If  $P \in \mathcal{P}_{ram}$ , then  $\pi^{-1}(P)$  contains at least one singular points  $Q$  of  $\mathcal{C}_1$  with branch data  $b_Q = (a, -a)$  for some  $a \neq 0$  in  $\mathbb{Z}_n$ .

In fact, after blowing-up singular points  $Q'$  in  $\pi^{-1}(P)$  with trivial branch-data we obtain a smooth surface  $S \twoheadrightarrow S_1 \twoheadrightarrow X$  such that the only singular points of the ramification divisor  $\mathcal{C}'$  of  $\Delta$  have non-trivial branch-data  $(a, -a)$  for some  $a \in \mathbb{Z}_n$ . Then, take a maximal  $\mathcal{O}_S$ -order  $\mathcal{A}$  in  $\Delta$ . By the local calculation of  $Br_n \mathbb{C}\{x, y\}$  performed in the last section we know that locally étale  $\mathcal{A}$  is of quantum-plane type in these remaining singularities. As the quantum-plane is not étale locally split,  $\mathcal{A}$  is not Cayley-smooth in these finite number of singularities.  $\square$

In fact, the above proof gives a complete classification of the central simple algebras admitting a Cayley-smooth model.

**THEOREM 5.35.** *Let  $\Delta$  be a central simple  $\mathbb{C}(X)$ -algebra of dimension  $n^2$  determined by the geo-combinatorial data  $(\mathcal{C}, \mathcal{P}, b, \mathcal{D})$  given by the Artin-Mumford sequence. Then,  $\Delta$  admits a Cayley-smooth model if and only if all branch-data are trivial.*

**PROOF.** If all branch-data are trivial, the foregoing proof constructs a Cayley-smooth model of  $\Delta$ . Conversely, if  $\mathcal{A}$  is a Cayley-smooth  $\mathcal{O}_S$ -order in  $\Delta$  with  $S$  a smooth projective model of  $\mathbb{C}(X)$ , then  $\mathcal{A}$  is locally étale split in every point  $s \in S$ . But then, so is any maximal  $\mathcal{O}_S$ -order  $\mathcal{A}_{max}$  containing  $\mathcal{A}$ . By the foregoing arguments this can only happen if all branch-data are trivial.  $\square$

### 5.6. Complex moment map.

We fix a quiver  $Q$  on  $k$  vertices  $\{v_1, \dots, v_k\}$  and define the *opposite quiver*  $Q^\circ$  the quiver on  $\{v_1, \dots, v_k\}$  obtained by reversing all arrows in  $Q$ . That is, there is an arrow  $\textcircled{j} \xrightarrow{a^*} \textcircled{i}$  in  $Q^\circ$  for each arrow  $\textcircled{j} \xleftarrow{a} \textcircled{i}$  in the quiver  $Q$ . Fix a dimension vector  $\alpha = (a_1, \dots, a_k)$ , using the *trace pairings*

$$M_{a_i \times a_j} \times M_{a_j \times a_i} \longrightarrow \mathbb{C} \quad (V_{a^*}, V_a) \mapsto \text{tr}(V_{a^*} V_a)$$

we can identify the representation space  $\text{rep}_\alpha Q^\circ$  with the *dual space*  $(\text{rep}_\alpha Q)^* = \text{Hom}_{\mathbb{C}}(\text{rep}_\alpha Q, \mathbb{C})$ . Observe that the base change action of  $GL(\alpha)$  on  $\text{rep}_\alpha Q^\circ$  coincides with the action dual to that of  $GL(\alpha)$  on  $\text{rep}_\alpha Q$ .

The *dual quiver*  $Q^d$  is the superposition of the quivers  $Q$  and  $Q^\circ$ . Clearly, for an dimension vector  $\alpha$  we have

$$\text{rep}_\alpha Q^d = \text{rep}_\alpha Q \oplus \text{rep}_\alpha Q^\circ = \text{rep}_\alpha Q \oplus (\text{rep}_\alpha Q)^*$$

whence  $\text{rep}_\alpha Q^d$  can be viewed as the *cotangent bundle*  $T^*\text{rep}_\alpha Q$  on  $\text{rep}_\alpha Q$  with structural morphism projection on the first factor. Cotangent bundles are equipped with a canonical *symplectic structure*, see [17, Example 1.1.3] or chapter 8 for more details. The natural action of  $GL(\alpha)$  on  $\text{rep}_\alpha Q$  extends to an action of  $GL(\alpha)$  on  $T^*\text{rep}_\alpha Q$  preserving the symplectic structure and it coincides with the basechange action of  $GL(\alpha)$  on  $\text{rep}_\alpha Q^d$ . Such an action on the cotangent bundle gives rise to a *complex moment map*

$$T^*\text{rep}_\alpha Q \xrightarrow{\mu_{\mathbb{C}}} (\text{Lie } GL(\alpha))^*$$

Recall that  $\text{Lie } GL(\alpha) = M_\alpha(\mathbb{C}) = M_{a_1}(\mathbb{C}) \oplus \dots \oplus M_{a_k}(\mathbb{C})$ . Using the trace pairings on both sides, the complex moment map is the mapping

$$\text{rep}_\alpha Q^d \xrightarrow{\mu_{\mathbb{C}}} M_\alpha(\mathbb{C})$$

defined by

$$\mu_{\mathbb{C}}(V)_i = \sum_{\substack{a \in Q_a \\ t(a)=i}} V_a V_{a^*} - \sum_{\substack{a \in Q_a \\ s(a)=i}} V_{a^*} V_a$$

Observe that the image of the complex moment map is contained in  $M_\alpha^0(\mathbb{C})$  where

$$M_\alpha^0(\mathbb{C}) = \{(M_1, \dots, M_k) \in M_\alpha(\mathbb{C}) \mid \sum_i \text{tr}(M_i) = 0\} = \text{Lie } PGL(\alpha)$$

corresponding to the fact that the action of  $GL(\alpha)$  on  $T^*\text{rep}_\alpha Q$  is really a  $PGL(\alpha) = GL(\alpha)/C^*$  action.

**DEFINITION 5.36.** Elements of  $\mathbb{C}^k = \mathbb{C}^{Q_v}$  are called *weights*. If  $\lambda$  is a weight, one defines the *deformed preprojective algebra* of the quiver  $Q$  to be

$$\Pi_\lambda(Q) \stackrel{\text{dfn}}{=} \Pi_\lambda = \frac{\mathbb{C}Q^d}{c - \lambda}$$

where  $c$  is the *commutator element*

$$c = \sum_{a \in Q_a} [a, a^*]$$

in  $\mathbb{C}Q^d$  and where  $\lambda = (\lambda_1, \dots, \lambda_k)$  is identified with the element  $\sum_i \lambda_i v_i \in \mathbb{C}Q^d$ . the algebra  $\Pi(Q) = \Pi$  is known as the *preprojective algebra* of the quiver  $Q$ .

LEMMA 5.37. *The ideal  $(c - \lambda) \triangleleft \mathbb{C}Q^d$  is the same as the ideal with a generator*

$$\sum_{\substack{a \in Q_a \\ t(a)=i}} aa^* - \sum_{\substack{a \in Q_a \\ s(a)=i}} a^*a - \lambda_i v_i$$

for each vertex  $v_i \in Q_v$ .

PROOF. These elements are of the form  $v_j(c - \lambda)v_i$ , so they belong to the ideal  $(c - \lambda)$ . As  $c - \lambda$  is also the sum of them, the ideal they generate contains  $c - \lambda$ .  $\square$

That is,  $\alpha$ -dimensional representations of the deformed preprojective algebra  $\Pi_\lambda$  coincide with representations  $V \in \text{rep}_\alpha Q^d$  which satisfy

$$\sum_{\substack{a \in Q_a \\ t(a)=i}} V_a V_{a^*} - \sum_{\substack{a \in Q_a \\ s(a)=i}} V_{a^*} V_a = \lambda_i \mathbb{1}_{a_i}$$

for each vertex  $v_i$ . That is, we have an isomorphism between the scheme theoretic fiber of the complex moment map and the representation space

$$\underline{\text{rep}}_\alpha \Pi_\lambda = \mu_{\mathbb{C}}^{-1}(\lambda)$$

As the image of  $\mu_{\mathbb{C}}$  is contained in  $M_\alpha^0(\mathbb{C})$  we have in particular

LEMMA 5.38. *If  $\lambda \cdot \alpha = \sum_i \lambda_i a_i \neq 0$ , then there are no  $\alpha$ -dimensional representations of  $\Pi_\lambda$ .*

Because we have an embedding  $C_k \hookrightarrow \Pi_\lambda$ , the  $n$ -dimensional representations of the deformed preprojective algebra decompose into disjoint subvarieties

$$\underline{\text{rep}}_n \Pi_\lambda = \bigsqcup_{\alpha: \sum_i a_i = n} GL_n \times^{GL(\alpha)} \underline{\text{rep}}_\alpha \Pi_\lambda$$

Hence, in studying Cayley-smoothness of  $\Pi_\lambda$  we may reduce to the distinct components and hence to the study of  $\alpha$ -Cayley-smoothness, that is, smoothness in the category of  $C_k(\alpha)$ -algebras which are Cayley-Hamilton algebras of degree  $n = \sum_i a_i$ . Again, one can characterize this smoothness condition in a geometric way by the property that the restricted representation scheme  $\underline{\text{rep}}_\alpha$  is smooth. In the next section we will investigate this property for the preprojective algebra  $\Pi_0$ , in chapter 8 we will be able to extend these results to arbitrary  $\Pi_\lambda$ . In this section we will compute the dimension of these representation schemes. First, we will investigate the fibers of the structural map of the cotangent bundle, that is, the projection

$$T^* \text{rep}_\alpha Q \simeq \text{rep}_\alpha Q^d \longrightarrow \text{rep}_\alpha Q$$

PROPOSITION 5.39. *If  $V \in \text{rep}_\alpha Q$ , then there is an exact sequence*

$$0 \longrightarrow \text{Ext}_{\mathbb{C}Q}^1(V, V)^* \longrightarrow \text{rep}_\alpha Q^o \xrightarrow{c} M_\alpha(\mathbb{C}) \xrightarrow{t} \text{Hom}_{\mathbb{C}Q}(V, V)^* \longrightarrow 0$$

where  $c$  maps  $W = (W_{a^*})_{a^*} \in \text{rep}_\alpha Q^o$  to  $\sum_{a \in Q_a} [V_a, W_{a^*}]$  and  $t$  maps  $M = (M_i)_i \in M_{\text{alpha}}(\mathbb{C})$  to the linear map  $\text{Hom}_{\mathbb{C}Q}(V, V) \longrightarrow \mathbb{C}$  sending a morphism  $N = (N_i)_i$  to  $\sum_i \text{tr}(M_i N_i)$ .

PROOF. There is an exact sequence

$$0 \longrightarrow \text{Hom}_{\mathbb{C}Q}(V, V) \longrightarrow M_\alpha(\mathbb{C}) \xrightarrow{f} \text{rep}_\alpha Q \longrightarrow \text{Ext}_{\mathbb{C}Q}^1(V, V) \longrightarrow 0$$



where  $f$  sends  $M = (M_i)_i \in M_\alpha(\mathbb{C})$  to  $V' = (V'_a)_a$  with  $V'_a = M_{t(a)}V_a - V_aM_{s(a)}$ . By definition, the kernel of  $f$  is  $\text{Hom}_{\mathbb{C}Q}(V, V)$  and by the Euler form interpretation of theorem 3.29 we have

$$\dim_{\mathbb{C}} \text{Hom}_{\mathbb{C}Q}(V, V) - \dim_{\mathbb{C}} \text{Ext}_{\mathbb{C}Q}^1(V, V) = \chi_Q(\alpha, \alpha) = \dim_{\mathbb{C}} M_\alpha(\mathbb{C}) - \dim_{\mathbb{C}} \text{rep}_\alpha Q$$

so the cokernel of  $f$  has the same dimension as  $\text{Ext}_{\mathbb{C}Q}^1(V, V)$  and using the standard projective resolution of  $V$  one can show that it is naturally isomorphic to it. The required exact sequence follows by dualizing, using the trace pairing to identify  $\text{rep}_\alpha Q^o$  with  $(\text{rep}_\alpha Q)^*$  and  $M_\alpha(\mathbb{C})$  with its dual.  $\square$

This result allows us to give a characterization of the dimension vectors  $\alpha$  such that  $\underline{\text{rep}}_\alpha Q \neq \emptyset$ .

**THEOREM 5.40.** *For a weight  $\lambda \in \mathbb{C}^k$  and a representation  $V \in \text{rep}_\alpha Q$  the following are equivalent*

- (1)  $V$  extends to an  $\alpha$ -dimensional representation of the deformed preprojective algebra  $\Pi_\lambda$ .
- (2) For all dimension vectors  $\beta$  of direct summands  $W$  of  $V$  we have  $\lambda \cdot \beta = 0$ .

Moreover, if  $V \in \text{rep}_\alpha Q$  does lift, then  $\pi^{-1}(V) \simeq (\text{Ext}_{\mathbb{C}Q}^1(V, V))^*$ .

**PROOF.** If  $V$  lifts to a representation of  $\Pi_\lambda$ , then there is a representation  $W \in \text{rep}_\alpha Q^o$  mapping under  $c$  of proposition 5.39 to  $\lambda$ . But then, by exactness of the sequence in proposition 5.39  $\lambda$  must be in the kernel of  $t$ . In particular, for any morphism  $N = (N_i)_i \in \text{Hom}_{\mathbb{C}Q}(V, V)$  we have that  $\sum_i \lambda_i \text{tr}(N_i) = 0$ . In particular, let  $W$  be a direct summand of  $V$  (as  $Q$ -representation) and let  $N = (N_i)_i$  be the projection morphism  $V \twoheadrightarrow W \hookrightarrow V$ , then  $\sum_i \lambda_i \text{tr}(N_i) = \sum_i \lambda_i b_i$  where  $\beta = (b_1, \dots, b_k)$  is the dimension vector of  $W$ .

Conversely, it suffices to prove the liftability of any indecomposable representation  $W$  having a dimension vector  $\beta$  satisfying  $\lambda \cdot \beta = 0$ . Because the endomorphism ring of  $W$  is a local algebra, any endomorphism  $N = (N_i)_i$  of  $W$  is the sum of a nilpotent matrix and a scalar matrix whence  $\sum_i \lambda_i \text{tr}(N_i) = 0$ . But then considering the sequence of proposition 5.39 for  $\beta$  and considering  $\lambda$  as an element of  $M_{|\beta|}(\mathbb{C})$ , it lies in the kernel of  $t$  whence in the image of  $c$  and therefore  $W$  can be extended to a representation of  $\Pi_\lambda$ .

The last statement follows again from the exact sequence of proposition 5.39.  $\square$

In particular, if  $\alpha$  is a root for  $Q$  satisfying  $\lambda \cdot \alpha = 0$ , then there are  $\alpha$ -dimensional representations of  $\Pi_\lambda$ . Recall the definition of the *number of parameters* given in definition 4.36

$$\mu(X) = \max_d (\dim X_{(d)} - d)$$

where  $X_{(d)}$  is the union of all orbits of dimension  $d$ . We denote  $\mu(\text{rep}_\alpha^{\text{ind}} Q)$  for the  $GL(\alpha)$ -action on the indecomposables of  $\text{rep}_\alpha Q$  by  $p_Q(\alpha)$ . Recall that part of Kac's theorem 4.39 asserts that

$$p_Q(\alpha) = 1 - \chi_Q(\alpha, \alpha)$$

We will apply these facts to the determination of the dimension of the fibers of the complex moment map.

LEMMA 5.41. *Let  $U$  be a  $GL(\alpha)$ -stable constructible subset of  $rep_\alpha Q$  contained in the image of the projection map  $rep_\alpha Q^d \xrightarrow{\pi} rep_\alpha Q$ . Then,*

$$\dim \pi^{-1}(U) = \mu(U) + \alpha.\alpha - \chi_Q(\alpha, \alpha)$$

*If in addition  $U = \mathcal{O}(V)$  is a single orbit, then  $\pi^{-1}(U)$  is irreducible of dimension  $\alpha.\alpha - \chi_Q(\alpha, \alpha)$ .*

PROOF. Let  $V \in U_{(d)}$ , then by theorem 5.40, the fiber  $\pi^{-1}(V)$  is isomorphic to  $(Ext_{\mathbb{C}Q}^1(V, V))^*$  and has dimension  $\dim_{\mathbb{C}} End(V) - \chi_Q(\alpha, \alpha)$  by theorem 3.29 and

$$\dim_{\mathbb{C}} End(V) = \dim GL(\alpha) - \dim \mathcal{O}(V) = \alpha.\alpha - d.$$

Hence,  $\dim \pi^{-1}(U_{(d)}) = (\dim U_{(d)} - d) + \alpha.\alpha - \chi_Q(\alpha, \alpha)$ . If we now vary  $d$ , the result follows.

For the second assertion, suppose that  $\pi^{-1}(U) \longleftarrow Z_1 \sqcup Z_2$  with  $Z_i$  a  $GL(\alpha)$ -stable open subset, but then  $\pi^{-1}(V) \cap Z_i$  are non-empty disjoint open subsets of the irreducible variety  $\pi^{-1}(V)$ , a contradiction.  $\square$

THEOREM 5.42. *Let  $\lambda$  be a weight and  $\alpha$  a dimension vector such that  $\lambda.\alpha = 0$ . Then,*

$$\dim \underline{rep}_\alpha \Pi_\lambda = \dim \mu_{\mathbb{C}}^{-1}(\lambda) = \alpha.\alpha - \chi_Q(\alpha, \alpha) + m$$

where  $m$  is the maximum number among all

$$p_Q(\beta_1) + \dots + p_Q(\beta_r)$$

with  $r \geq 1$ , all  $\beta_i$  are (positive) roots such that  $\lambda.\beta_i = 0$  and  $\alpha = \beta_1 + \dots + \beta_r$ .

PROOF. Decompose  $rep_\alpha Q = \bigsqcup_r rep_\alpha(\tau)$  where  $rep_\alpha(\tau)$  are the representations decomposing as a direct sum of indecomposables of dimension vector  $\tau = (\beta_1, \dots, \beta_r)$ . By Kac's theorem 4.39 we have that

$$\mu(rep_\alpha(\tau)) = p_Q(\beta_1) + \dots + p_Q(\beta_r)$$

If some of the  $\beta_i$  are such that  $\lambda.\beta_i \neq 0$ , and  $\mu_{\mathbb{C}}^{-1}(\lambda) \xrightarrow{\pi} rep_\alpha Q$  is the projection then  $\pi^{-1}(rep_\alpha(\tau)) = \emptyset$  by lemma 5.38. Combining this with lemma 5.41 the result follows.  $\square$

DEFINITION 5.43. The set of  $\lambda$ -Schur roots  $S_\lambda$  is defined to be the set of  $\alpha \in \mathbb{N}^k$  such that  $p_Q(\alpha) \geq p_Q(\beta_1) + \dots + p_Q(\beta_r)$  for all decompositions  $\alpha = \beta_1 + \dots + \beta_r$  with  $\beta_i$  positive roots satisfying  $\lambda.\beta_i = 0$ .

$S_0$  is the set of  $\alpha \in \mathbb{N}^k$  such that  $p_Q(\alpha) \geq p_Q(\beta_1) + \dots + p_Q(\beta_r)$  for all decompositions  $\alpha = \beta_1 + \dots + \beta_r$  with  $\beta_i \in \mathbb{N}^k$

Observe that  $S_0$  consists of Schur roots for  $Q$ , for if

$$\tau_{can} = (e_1, \beta_1; \dots; e_s, \beta_s) = (\gamma_1, \dots, \gamma_t)$$

(the  $\gamma_j$  possibly occurring with multiplicities) is the canonical decomposition of  $\alpha$  with  $t \geq 2$  we have

$$\begin{aligned} p_Q(\alpha) &= 1 - \chi_Q(\alpha, \alpha) \\ &= 1 - \sum_{i,j} \chi_Q(\gamma_i, \gamma_j) \\ &= \sum_i (1 - \chi_Q(\gamma_i, \gamma_i)) - \sum_{i \neq j} \chi_Q(\gamma_i, \gamma_j) - (t-1) \\ &> \sum_i p_Q(\gamma_i) \end{aligned}$$

whence  $\alpha \notin S_0$ . This argument also shows that in the definition of  $S_0$  we could have taken all decompositions in positive roots, replacing the components  $\beta_i$  by their canonical decompositions.

**THEOREM 5.44.** *For  $\alpha \in \mathbb{N}^k$ , the following are equivalent :*

- (1) *The complex moment map  $\text{rep}_\alpha Q^d \xrightarrow{\mu_{\mathbb{C}}} \text{rep}_\alpha Q$  is flat.*
- (2)  *$\text{rep}_\alpha \Pi_0 = \mu_{\mathbb{C}}^{-1}(\underline{0})$  has dimension  $\alpha \cdot \alpha - 1 + 2p_Q(\alpha)$ .*
- (3)  *$\alpha \in S_0$ .*

**PROOF.** The dimensions of the relevant representation spaces are

$$\begin{cases} \dim \text{rep}_\alpha Q &= \alpha \cdot \alpha - \chi_Q(\alpha, \alpha) = \alpha \cdot \alpha - 1 + p_Q(\alpha) \\ \dim \text{rep}_\alpha Q^d &= 2\alpha \cdot \alpha - 2\chi_Q(\alpha, \alpha) = 2\alpha \cdot \alpha - 2 + 2p_Q(\alpha) \\ \dim M_\alpha^0(\mathbb{C}) &= \alpha \cdot \alpha - 1 \end{cases}$$

so the relative dimension of the complex moment map is  $d = \alpha \cdot \alpha - 1 + 2p_Q(\alpha)$ .

(1)  $\Rightarrow$  (2) : Because  $\mu_{\mathbb{C}}$  is flat, its image  $U$  is an open subset of  $M_\alpha^0(\mathbb{C})$  which obviously contains  $\underline{0}$ , but then the dimension of  $\mu_{\mathbb{C}}^{-1}(\underline{0})$  is equal to the relative dimension  $d$ .

(2)  $\Rightarrow$  (3) : Assume  $p_Q(\alpha) < \sum_i p_Q(\beta_i)$  for some decomposition  $\alpha = \beta_1 + \dots + \beta_s$  with  $\beta_i \in \mathbb{N}^k$ . Replacing each  $\beta_i$  by its canonical decomposition, we may assume that the  $\beta_i$  are actually positive roots. But then, theorem 5.42 implies that  $\mu_{\mathbb{C}}^{-1}(\underline{0})$  has dimension greater than  $d$ .

(3)  $\Rightarrow$  (1) : We have that  $\alpha$  is a Schur root. We claim that  $\text{rep}_\alpha Q^d \xrightarrow{\mu_{\mathbb{C}}} M_\alpha^0(\mathbb{C})$  is surjective. Let  $V \in \text{rep}_\alpha Q$  be a general representation, then  $\text{Hom}_{\mathbb{C}Q}(V, V) = \mathbb{C}$ . But then, the map  $c$  in proposition 5.39 has a one-dimensional cokernel. But as the image of  $c$  is contained in  $M_\alpha^0(\mathbb{C})$ , this shows that

$$\text{rep}_\alpha Q^0 \xrightarrow{c} M_\alpha^0(\mathbb{C})$$

is surjective from which the claim follows. Let  $M = (M_i)_i \in M_\alpha^0(\mathbb{C})$  and consider the projection

$$\mu_{\mathbb{C}}^{-1}(M) \xrightarrow{\tilde{\pi}} \text{rep}_\alpha Q$$

If  $U$  is a constructible  $GL(\alpha)$ -stable subset of  $\text{rep}_\alpha Q$ , then by an argument as in lemma 5.41 we have that

$$\dim \tilde{\pi}^{-1}(U) \leq \mu(U) + \alpha \cdot \alpha - \chi_Q(\alpha, \alpha)$$

But then, decomposing  $\text{rep}_\alpha Q$  into types  $\tau$  of direct sums of indecomposables, it follows from the assumption that  $\mu_{\mathbb{C}}^{-1}(M)$  has dimension at most  $d$ . But then by the dimension formula it must be equidimensional of dimension  $d$  whence flat.  $\square$

### 5.7. Preprojective algebras.

In this section we will determine the  $n$ -smooth locus of the preprojective algebra  $\Pi_0$ . By the étale local description of section 4.5 it is clear that we need to control the  $Ext^1$ -spaces of representations of  $\Pi_0$ .

PROPOSITION 5.45. *Let  $V$  and  $W$  be representations of  $\Pi_0$  of dimension vectors  $\alpha$  and  $\beta$ , then we have*

$$\dim_{\mathbb{C}} Ext_{\Pi_0}^1(V, W) = \dim_{\mathbb{C}} Hom_{\Pi_0}(V, W) + \dim_{\mathbb{C}} Hom_{\Pi_0}(W, V) - T_Q(\alpha, \beta)$$

PROOF. It is easy to verify by direct computation that  $V$  has a projective resolution as  $\Pi_0$ -module which starts as

$$\dots \longrightarrow \bigoplus_{i \in Q_v} \Pi_0 v_i \otimes v_i V \xrightarrow{f} \bigoplus_{\substack{j \leftarrow_a i \\ a \in Q_a^d}} \Pi_0 v_j \otimes v_i V \xrightarrow{g} \bigoplus_{i \in Q_v} \Pi_0 v_i \otimes v_i V \xrightarrow{h} V \longrightarrow 0$$

where  $f$  is defined by

$$f\left(\sum_i p_i \otimes m_i\right) = \sum_{\substack{j \leftarrow_a i \\ a \in Q_a}} (p_i a^* \otimes m_i - p_j \otimes a^* m_j)_a - (p_j a \otimes m_j - p_i \otimes a m_i)_{a^*}$$

where  $p_i \in \Pi_0 v_i$  and  $m_i \in v_i V$ . The map  $g$  is defined on the summand corresponding to an arrow  $j \leftarrow_a i$  in  $Q^d$  by

$$g(pa \otimes m) = (pa \otimes m)_i - (p \otimes am)_j$$

for  $p \in \Pi_0 v_j$  and  $m \in v_i V$ . the map  $h$  is the multiplication map. If we compute homomorphisms to  $W$  and use the identification

$$Hom_{\Pi_0}(\Pi_0 v_j \otimes v_i V, W) = Hom_{\mathbb{C}}(v_i V, v_j W)$$

we obtain a complex

$$0 \longrightarrow \bigoplus_{i \in Q_v} Hom_{\mathbb{C}}(v_i V, v_i W) \longrightarrow \bigoplus_{\substack{j \leftarrow_a i \\ a \in Q_a^d}} Hom_{\mathbb{C}}(v_i V, v_j W) \longrightarrow \bigoplus_{i \in Q_v} Hom_{\mathbb{C}}(v_i V, v_i W)$$

in which the left hand cohomology is  $Hom_{\Pi_0}(V, W)$  and the middle cohomology is  $Ext_{\Pi_0}^1(V, W)$ . Moreover, the alternating sum of the dimensions of the terms is  $T_Q(\alpha, \beta)$ . It remains to prove that the cokernel of the right hand side map has the same dimension as  $Hom_{\Pi_0}(W, V)$ . But using the trace pairing to identify

$$Hom_{\mathbb{C}}(M, N)^* = Hom_{\mathbb{C}}(N, M)$$

we obtain that the dual of this complex is

$$\bigoplus_{i \in Q_v} Hom_{\mathbb{C}}(v_i W, v_i V) \longrightarrow \bigoplus_{\substack{j \leftarrow_a i \\ a \in Q_a^d}} Hom_{\mathbb{C}}(v_i W, v_j V) \longrightarrow \bigoplus_{i \in Q_v} Hom_{\mathbb{C}}(v_i W, v_i V) \longrightarrow 0$$

and, up to changing the sign of components in the second direct sum corresponding to arrows which are not in  $Q$ , this is the same complex as the complex arising with  $V$  and  $W$  interchanged. From this the result follows.  $\square$

In order to determine the  $n$ -smooth locus we observe that the representation space decomposes into a disjoint union and we have quotient morphisms

$$\begin{array}{ccc} \underline{rep}_n \Pi_0 & \xrightarrow{=} & \bigsqcup_{\substack{\alpha=(a_1, \dots, a_k) \\ a_1 + \dots + a_k = n}} GL_n \times^{GL(\alpha)} \underline{rep}_\alpha \Pi_0 \\ \pi_n \downarrow & & \sqcup \pi_\alpha \downarrow \\ \underline{iss}_n \Pi_0 & \xrightarrow{=} & \bigsqcup_{\substack{\alpha=(a_1, \dots, a_k) \\ a_1 + \dots + a_k = n}} \underline{iss}_\alpha \Pi_0 \end{array}$$

Hence if  $\xi \in \underline{iss}_\alpha \Pi_0$  for  $\xi \in Sm_n \Pi_0$  it is necessary and sufficient that  $\underline{rep}_\alpha \Pi_0$  is smooth along  $\mathcal{O}(M_\xi)$  where  $M_\xi$  is the semi-simple  $\alpha$ -dimensional representation of  $\Pi_0$  corresponding to  $\xi$ . Assume that  $\xi$  is of type  $\tau = (e_1, \alpha_1; \dots; e_z, \alpha_z)$ , that is,

$$M_\xi = S_1^{\oplus e_1} \oplus \dots \oplus S_z^{\oplus e_z}$$

with  $S_i$  a simple  $\Pi_0$ -representation of dimension vector  $\alpha_i$ . Again, the normal space to the orbit  $\mathcal{O}(M_\xi)$  is determined by  $Ext_{\Pi_0}^1(M_\xi, M_\xi)$  and can be depicted by a local quiver setting  $(Q_\xi, \alpha_\xi)$  where  $Q_\xi$  is a quiver on  $z$  vertices and where  $\alpha_\xi = \alpha_\tau = (e_1, \dots, e_z)$ . Repeating the arguments of section 4.5 we have

LEMMA 5.46. *With notations as above,  $\xi \in Sm_n \Pi_0$  if and only if*

$$\dim GL(\alpha) \times^{GL(\alpha_\xi)} Ext_{\Pi_0}^1(M_\xi, M_\xi) = \dim_{M_\xi} \underline{rep}_\alpha \Pi_0$$

As we have enough information to compute both sides, we can prove :

THEOREM 5.47. *If  $\xi \in \underline{iss}_\alpha \Pi_0$  with  $\alpha = (a_1, \dots, a_k) \in S_0$  and  $\sum_i a_i = n$ , then  $\xi \in Sm_n \Pi_0$  if and only if  $M_\xi$  is a simple  $n$ -dimensional representation of  $\Pi_0$ .*

PROOF. Assume that  $\xi$  is a point of semi-simple representation type  $\tau = (e_1, \alpha_1; \dots; e_z, \alpha_z)$ , that is,

$$M_\xi = S_1^{\oplus e_1} \oplus \dots \oplus S_z^{\oplus e_z} \quad \text{with} \quad \dim(S_i) = \alpha_i$$

and  $S_i$  a simple  $\Pi_0$ -representation. Then, by proposition 5.45 we have

$$\begin{cases} \dim_{\mathbb{C}} Ext_{\Pi_0}^1(S_i, S_j) &= -T_Q(\alpha_i, \alpha_j) & i \neq j \\ \dim_{\mathbb{C}} Ext_{\Pi_0}^1(S_i, S_i) &= 2 - T_Q(\alpha_i, \alpha_i) \end{cases}$$

But then, the dimension of  $Ext_{\Pi_0}^1(M_\xi, M_\xi)$  is equal to

$$\sum_{i=1}^z (2 - T_Q(\alpha_i, \alpha_i)) e_i^2 + \sum_{i \neq j} e_i e_j (-T_Q(\alpha_i, \alpha_j)) = 2 \sum_{i=1}^z e_i - T_Q(\alpha, \alpha)$$

from which it follows immediately that

$$\dim GL(\alpha) \times^{GL(\alpha_\xi)} Ext_{\Pi_0}^1(M_\xi, M_\xi) = \alpha \cdot \alpha + \sum_{i=1}^z e_i^2 - T_Q(\alpha, \alpha)$$

On the other hand, as  $\alpha \in S_0$  we know from theorem 5.44 that

$$\dim \underline{rep}_\alpha \Pi_0 = \alpha \cdot \alpha - 1 + 2p_Q(\alpha) = \alpha \cdot \alpha - 1 + 2 + 2\chi_Q(\alpha, \alpha) = \alpha \cdot \alpha + 1 - T_Q(\alpha, \alpha)$$

But then, equality occurs if and only if  $\sum_i e_i^2 = 1$ , that is,  $\tau = (1, \alpha)$  or  $M_\xi$  is a simple  $n$ -dimensional representation of  $\Pi_0$ .  $\square$

In particular it follows that the preprojective algebra  $\Pi_0$  is *never* Quillen-smooth. Further, as  $\vec{v}_i = (0, \dots, 1, 0, \dots, 0)$  are dimension vectors of simple representations of  $\Pi_0$  it follows that  $\Pi_0$  is  $\alpha$ -smooth if and only if  $\alpha = \vec{v}_i$  for some  $i$ . In chapter 8 we will determine the dimension vectors of simple representations of the (deformed) preprojective algebras.

EXAMPLE 5.48. Let  $Q$  be an extended Dynkin diagram and  $\delta_Q$  the corresponding dimension vector. Then, we will show that  $\delta_Q$  is the dimension vector of a simple representation and  $\delta_Q \in S_0$ . Then, the dimension of the quotient variety

$$\begin{aligned} \dim \text{iss}_{\delta_Q} \Pi_0 &= \dim \text{rep}_{\delta_Q} \Pi_0 - \delta_Q \cdot \delta_Q + 1 \\ &= 2p_Q(\delta_Q) = 2 \end{aligned}$$

so it is a surface. The only other semi-simple  $\delta_Q$ -dimensional representation of  $\Pi_0$  is the trivial representation. By the theorem, this must be an isolated singular point of  $\text{iss}_{\delta_Q} Q$ . In fact, one can show that  $\text{iss}_{\delta_Q} \Pi_0$  is the Kleinian singularity corresponding to the extended Dynkin diagram  $Q$ .

### 5.8. Quantum groups.

In this section we will give another class of interesting noncommutative algebras which are as singular as possible : *quantum groups* at root of unity. Throughout we will use structural results on quantum groups which can be found in [19].

Let  $\mathfrak{g}$  be a *semi-simple Lie algebra* over  $\mathbb{C}$  of rank  $r$  with *Cartan matrix*  $C = (a_{ij}) \in M_r(\mathbb{Z})$  and vector  $d = (d_1, \dots, d_r) \in \mathbb{N}_+^r$  of relative prime integers such that  $d.C$  is symmetric.

The (simply connected form of the) *quantised enveloping algebra*  $U_\epsilon(\mathfrak{g})$  is the  $\mathbb{C}$ -algebra with generators

$$\{E_i, F_i, K_i^{\pm 1} \mid 1 \leq i, j \leq r\}$$

satisfying the following relations for all  $1 \leq i, j \leq r$  :

$$\begin{aligned} K_i K_j &= K_j K_i & K_i K_i^{-1} &= 1 = & K_i^{-1} K_i \\ & & K_i E_j &= & \epsilon^{d_i a_{ij}} E_j K_i \\ & & K_i F_j &= & \epsilon^{-d_i a_{ij}} F_j K_i \\ & & E_i F_j - F_j E_i &= & \delta_{ij} \frac{K_i - K_i^{-1}}{\epsilon^{d_i} - \epsilon^{-d_i}} \\ \sum_{s=0}^{1-a_{ij}} (-1)^s \begin{bmatrix} 1 - a_{ij} \\ s \end{bmatrix}_{d_i} E_i^{1-a_{ij}-s} E_j E_i^s &= & 0 \\ \sum_{s=0}^{1-a_{ij}} (-1)^s \begin{bmatrix} 1 - a_{ij} \\ s \end{bmatrix}_{d_i} F_i^{1-a_{ij}-s} F_j F_i^s &= & 0 \end{aligned}$$

where the symbols in squarebrackets denote Gaussian binomial coefficients for the parameter  $\epsilon$ , see [19].

The *Hopf structure* on this algebra is given by defining for all  $1 \leq i \leq r$

$$\begin{aligned} \Delta(E_i) &= E_i \otimes 1 + K_i \otimes E_i, \Delta(F) = F_i \otimes K_i^{-1} + 1 \otimes F_i, \Delta(K_i) = K_i \otimes K_i \\ S(E_i) &= -K_i^{-1} E_i, S(F) = -F_i K_i, S(K_i) = K_i^{-1} \\ \epsilon(E_i) &= 0, \epsilon(F_i) = 0, \epsilon(K_i) = 1. \end{aligned}$$


Now, let  $\epsilon$  be a primitive  $l$ -th root of unity, where  $l$  is odd and prime to 3 if  $\mathfrak{g}$  contains components of type  $G_2$ . We recall the following structural results on  $U_\epsilon(\mathfrak{g})$  from [19].

$U_\epsilon(\mathfrak{g})$  is an order with integrally closed center  $Z_\epsilon$  in a central simple algebra of dimension  $l^{2N}$  where  $N$  is the number of positive roots of  $\mathfrak{g}$ . The commutator in  $U_q(\mathfrak{g})$  induces a nontrivial *Poisson structure* on  $Z_\epsilon$ .


There exists a Poisson subalgebra  $Z_0$  of  $Z_\epsilon$  satisfying the following properties :

- (1)  $Z_0$  is a sub-Hopf algebra of  $U_\epsilon(\mathfrak{g})$ .
- (2)  $U_\epsilon(\mathfrak{g})$  is a free  $Z_0$ -module of rank  $l^{2N+r}$ .
- (3)  $Z_\epsilon$  is a free  $Z_0$ -module of rank  $l^r$ .

That is, we have the following inclusions



Being an affine commutative Hopf algebra,  $Z_0 \simeq \mathbb{C}[H]$  for an algebraic group  $H$  which can be described in the following way. Let  $T$  be a maximal torus of the simply connected Lie group  $G$  corresponding to  $\mathfrak{g}$ , and  $B_\pm$  the corresponding Borel subgroups. Then, we have a fiber diagram



where  $pr_\pm$  are the projections and  $pr_-^{-1} = (-)^{-1} \circ pr_-$ . The connection between  $H$  and  $G$  is given by the map

$$\sigma : H \longrightarrow G^0 \quad (h_-, h_+) \mapsto h_-^{-1} h_+$$

which is an unramified cover of the big cell  $G^0$  of  $G$  of degree  $2^r$ .

There is an (infinite dimensional) group  $\tilde{G}$  of analytic automorphisms of  $H$  such that its orbits in  $H$  are of the form

$$\sigma^{-1}(\mathcal{C} \cap G^0)$$

where  $\mathcal{C}$  is the conjugacy class of a non-central element in  $G$ . These orbits are also the symplectic leaves of  $H$  induced by the Poisson structure on  $Z_0 = \mathbb{C}[H]$ .

In this section we investigate to what extent the inclusion of Hopf algebras  $\mathbb{C}[H] \hookrightarrow U_\epsilon(\mathfrak{g})$  differs from a finite Hopf algebra extension between the coordinate rings of irreducible algebraic groups.

Recall the commutative situation. Let  $G$  be an irreducible *algebraic group* with coordinate ring  $\mathbb{C}[G]$ . Consider a subHopf algebra  $\mathbb{C}[H]$  such that  $\mathbb{C}[G]$  is a finite  $\mathbb{C}[H]$ -module. Corresponding to the inclusion of Hopf algebras  $\mathbb{C}[H] \hookrightarrow \mathbb{C}[G]$  is a projection of varieties (or group-schemes)  $G \twoheadrightarrow H$  which is a finite étale morphism, that is, unramified and smooth.

Since there is no suitable substitute for the variety corresponding to  $U_\epsilon(\mathfrak{g})$ , we will give a different geometric description of the extension  $\mathbb{C}[H] \hookrightarrow \mathbb{C}[G]$  which can be generalized to the non-commutative setting.

Consider the trace map  $tr$  on the finite Galois extension of degree  $d$  between the function fields  $\mathbb{C}(H) \hookrightarrow \mathbb{C}(G)$ . As  $\mathbb{C}[G]$  and  $\mathbb{C}[H]$  are integrally closed,  $tr$  restricts to a linear map on  $\mathbb{C}[G]$  with image  $tr \mathbb{C}[G] = \mathbb{C}[H]$ . Remark that  $tr$  satisfies the formal Cayley-Hamilton identity for  $d \times d$  matrices. That is,  $\mathbb{C}[G]$  is a Cayley-Hamilton algebra of degree  $d$  with a trace map  $tr$  such that  $tr \mathbb{C}[G] = \mathbb{C}[H]$ .

Consider the affine algebraic variety  $\underline{rep}_d^{tr} \mathbb{C}[G]$  with points the trace preserving algebra maps  $\mathbb{C}[G] \xrightarrow{\rho} M_d(\mathbb{C})$ .

By the reconstruction results of section 2 we can recover  $\mathbb{C}[G]$  and  $\mathbb{C}[H]$  from the  $GL_d$ -variety  $\underline{rep}_d^{tr} \mathbb{C}[G]$ .

- (1)  $\mathbb{C}[H]$  is the ring of polynomial invariants  $\mathbb{C}[\underline{rep}_d^{tr} \mathbb{C}[G]]^{GL_d}$ .
- (2)  $\mathbb{C}[G]$  is the ring of  $GL_d$ -equivariant maps  $\underline{rep}_d^{tr} \mathbb{C}[G] \rightarrow M_d(\mathbb{C})$ .

By the first fact, we have an algebraic quotient map

$$\underline{rep}_d^{tr} \mathbb{C}[G] \xrightarrow{\pi} H$$

Consider  $h \in H$  with corresponding maximal ideal  $m_h \triangleleft \mathbb{C}[H]$ . The fiber  $\pi^{-1}(h)$  contains a unique closed orbit which is the orbit of the unique (trace preserving)  $d$ -dimensional semi-simple  $\mathbb{C}[G]/m_h \mathbb{C}[G]$ -module. The full fiber  $\pi^{-1}(h)$  consists of all  $d$ -dimensional (trace preserving)  $\mathbb{C}[G]/m_h \mathbb{C}[G]$ -modules.

Because  $G \twoheadrightarrow H$  is an étale map,  $\mathbb{C}[G]/m_h \mathbb{C}[G] \simeq \mathbb{C} \oplus \dots \oplus \mathbb{C}$  ( $d$  components) is a semi-simple algebra and thus  $\pi^{-1}(h)$  is a single orbit isomorphic to  $GL_d/T_d$  where  $T_d$  is a maximal torus of  $GL_d$ . Hence,  $\underline{rep}_d^{tr} \mathbb{C}[G]$  is a principal fibration over  $H$  with fibres the homogeneous space  $GL_d/T_d$ .

**PROPOSITION 5.49.** *Let  $\mathbb{C}[H] \hookrightarrow \mathbb{C}[G]$  be a finite Hopf algebra extension of degree  $d$ , then  $\underline{rep}_d^{tr} \mathbb{C}[G]$  is a smooth (in particular, reduced)  $GL_d$ -variety with algebraic (even geometric) quotient variety  $H$ .*

We will investigate whether this result remains true if we replace  $\mathbb{C}[G]$  by a non-commutative Hopf algebra such as  $U_\epsilon(\mathfrak{g})$ .

We can define a trace map on  $U_\epsilon(\mathfrak{g})$  by taking the composition of the reduced trace map and the trace map of the extension  $\mathbb{C}[H] \hookrightarrow Z_\epsilon$ . Again, using the fact that both  $Z_\epsilon$  and  $Z_0 = \mathbb{C}[H]$  are integrally closed, this trace map  $tr$  is well defined with image  $tr U_\epsilon(\mathfrak{g}) = \mathbb{C}[H]$ . This time,  $tr$  satisfies the formal Cayley-Hamilton identity for  $d \times d$  matrices where  $d = l^{N+r}$ .

Again, we define the affine algebraic variety  $\underline{rep}_d^{tr} U_\epsilon(\mathfrak{g})$  of trace preserving algebra maps  $U_\epsilon(\mathfrak{g}) \xrightarrow{\rho} M_d(\mathbb{C})$ . This variety has a natural  $GL_d$ -action and orbits correspond to isomorphism classes of  $d$ -dimensional trace preserving  $U_\epsilon(\mathfrak{g})$ -modules.

We recover  $U_\epsilon(\mathfrak{g})$  and  $\mathbb{C}[H]$  from the  $GL_d$ -variety  $X_{U_\epsilon(\mathfrak{g})}$ .

- (1)  $\mathbb{C}[H]$  is the ring of polynomial invariants  $\mathbb{C}[\underline{rep}_d^{tr} U_\epsilon(\mathfrak{g})]^{GL_d}$ .



(2)  $U_\epsilon(\mathfrak{g})$  is the ring of  $GL_d$ -equivariant maps  $\underline{rep}_d^{tr} U_\epsilon(\mathfrak{g}) \longrightarrow M_d(\mathbb{C})$ .

We have an algebraic quotient variety

$$\underline{rep}_d^{tr} U_\epsilon(\mathfrak{g}) \xrightarrow{\pi} H$$

and the description of the fibers  $\pi^{-1}(h)$  is as above : it consists of all  $d$ -dimensional trace preserving  $\bar{U}_h = U_\epsilon(\mathfrak{g})/m_h U_\epsilon(\mathfrak{g})$ -modules and there is a unique closed orbit corresponding to a semi-simple  $U_\epsilon(\mathfrak{g})$ -module which we denote by

$$M_h^{ss} = S_1^{\oplus e_1} \oplus \dots \oplus S_k^{\oplus e_k}$$

where the  $S_i$  is a simple  $U_\epsilon(\mathfrak{g})$ -module of dimension  $d_i$  occurring in  $M_h^{ss}$  with multiplicity  $e_i$ . Clearly,  $d = \sum d_i e_i$ .

Recall that  $h$  is said to be a  $d$ -smooth point of  $U_\epsilon(\mathfrak{g})$  if and only if  $\underline{rep}_d^{tr} U_\epsilon(\mathfrak{g})$  is smooth along  $\pi^{-1}(h)$ , or equivalently, is smooth in  $M_h^{ss}$ .

The  $d$ -smooth locus  $Sm_{tr} U_\epsilon(\mathfrak{g})$  of  $U_\epsilon(\mathfrak{g})$  over  $\mathbb{C}[H]$  is the subset of all  $h \in H$  such that  $U_\epsilon(\mathfrak{g})$  is smooth in  $h$ .

An alternative geometric description of  $U_\epsilon(\mathfrak{g})$  is as follows. Because  $U_\epsilon(\mathfrak{g})$  is a free  $\mathbb{C}[H]$ -module of rank  $t = l^{dim \mathfrak{g}}$ , every  $\bar{U}_h$  is a  $\mathbb{C}$ -algebra of dimension  $t$ . This determines a map

$$H \xrightarrow{\phi} Alg_t$$

where  $Alg_t$  is the *variety of (structure constants of)  $t$ -dimensional associative algebras* with unit. One can recover  $U_\epsilon(\mathfrak{g})$  from  $\phi$  by taking  $\phi^*$  of the generic  $t$ -dimensional algebra over  $Alg_t$ .

For sufficiently general  $h$  we have that  $\bar{U}_h \simeq M_N(\mathbb{C}) \oplus \dots \oplus M_N(\mathbb{C})$  ( $r$  copies) a semi-simple algebra. This entails that  $\phi(H)$  is contained in the irreducible component of  $Alg_t$  which is the closure of the  $GL_t$ -orbit  $\mathcal{O}$  of  $M_N(\mathbb{C}) \oplus \dots \oplus M_N(\mathbb{C})$ . We define  $\phi^{-1}(\mathcal{O})$  as the *separability locus* of  $U_\epsilon(\mathfrak{g})$  over  $\mathbb{C}[H]$  and denote it with  $Sep U_\epsilon(\mathfrak{g})/\mathbb{C}[H]$ . Clearly, it is the Zariski open subset of  $H$  consisting of those  $h$  such that  $\bar{U}_h$  is a semi-simple algebra (and hence isomorphic to  $M_N(\mathbb{C}) \oplus \dots \oplus M_N(\mathbb{C})$ ). As the terminology suggests, it is also the locus over which  $U_\epsilon(\mathfrak{g})$  is a separable  $\mathbb{C}[H]$ -algebra. Observe that this is a natural extension of the notion of the Azumaya locus.

As in the commutative case, we see that  $\pi^{-1}(h)$  consists of a single orbit when  $h \in Sep U_\epsilon(\mathfrak{g})/\mathbb{C}[H]$  and that this orbit is the homogeneous space  $GL_d/T_r$  where  $T_r$  is the center of  $GL_N \times \dots \times GL_N \hookrightarrow GL_d$ . Hence,  $X_{U_\epsilon(\mathfrak{g})}$  is a principal fibration and therefore smooth over  $Sep U_\epsilon(\mathfrak{g})/\mathbb{C}[H]$ . Thus,

$$Sep U_\epsilon(\mathfrak{g})/\mathbb{C}[H] \hookrightarrow Sm_{tr} U_\epsilon(\mathfrak{g})/\mathbb{C}[H].$$

For general Hopf algebras, one expects the smooth locus to be larger. For example, if  $U_\epsilon(\mathfrak{b})$  is the *quantum Borel* of  $\mathfrak{sl}_2$  it follows from example 5.15 that its smooth locus is the whole of  $B$ . However, we will prove :

**THEOREM 5.50.** *The quantised enveloping algebra  $U_\epsilon(\mathfrak{g})$  is as singular as possible. That is,*

$$Sm U_\epsilon(\mathfrak{g})/\mathbb{C}[G] = Sep U_\epsilon(\mathfrak{g})/\mathbb{C}[H]$$

A similar result holds for  $O_\epsilon(G)$ , that is, the *quantum function algebra* of  $G$ .

We will outline the strategy of proof in the case of  $U_\epsilon(\mathfrak{g})$ . Assume that  $U_\epsilon(\mathfrak{g})$  is smooth in  $h$  then the normal space to  $\mathcal{O}(M_h^{ss})$  is given by a certain quiver setting  $(Q_h, \alpha_h)$ . Because the quotient variety  $\underline{iss}_d U_\epsilon(\mathfrak{g}) = H$  is smooth in  $h$  it follows


from the slice results that the ring of polynomial invariants  $\mathbb{C}[\text{rep}_{\alpha_h} Q_h]^{GL(\alpha_h)}$  of this quiver setting must be a polynomial algebra.

The problem to determine all *coregular (marked) quiver situations* is a hard one. However, we will prove the following rough classification.

**THEOREM 5.51.** *Let  $Q^\bullet$  be a strongly connected marked quiver and  $\alpha = (a_1, \dots, a_k)$  a dimension vector.*


*If the ring of polynomial invariants  $\mathbb{C}[\text{rep}_\alpha Q^\bullet]^{GL(\alpha)}$  is a polynomial ring, then we are in one of the following situations :*

- (1) *type 1 :  $\min_i a_i \leq 1$  or*
- (2) *type 2 :  $\min_i a_i = 2$  and  $Q^\bullet$  has the form  $\tilde{A}_k(+1)$  :*



*that is, the extended Dynkin diagram with cyclic orientation and one extra arrow (which may degenerate to a cycle and loop or to two loops, possibly marked), or*

- (3) *type 3 :  $\min_i a_i \geq 2$  and the marked quiver  $Q^\bullet$  is the extended Dynkin diagram  $\tilde{A}_k$  with cyclic orientation*




*(which may degenerate to one loop possibly marked).*

**PROOF.** Let us assume that  $\min_i a_i \geq 2$ . Because  $Q^\bullet$  is strongly connected  $(1, \dots, 1)$  is the dimension vector of a simple representation and there are infinitely many isoclasses of such representations.

Hence, in  $\text{rep}_\alpha Q^\bullet$  there are semi-simple representations of type

$$\tau = (1, (1, \dots, 1); 1, (1, \dots, 1); a_1 - 2, \delta_1; \dots; a_k - 2, \delta_k)$$

The local marked quiver  $Q_\tau^\bullet$  has at most  $k + 2$  vertices and the full subquiver on the first two vertices is of the following form



where there are  $a$  directed arrows between the vertices with

$$a = -\chi_Q((1, \dots, 1), (1, \dots, 1)) = \sum_i \left( \sum_j a_{ij} + a_{ii} + m_{ii} - 1 \right)$$


and  $u$  (resp.  $m$ ) loops (resp. marked loops) in the vertices where

$$\begin{aligned} u &= 1 - \chi_{Q^\bullet}^1((1, \dots, 1), (1, \dots, 1)) = \sum_{i,j} a_{ij} + \sum_i a_{ii} - k + 1 \\ m &= -\chi_{Q^\bullet}^2((1, \dots, 1), (1, \dots, 1)) = \sum_i m_{ii} \end{aligned}$$

and the dimension vector  $\alpha_\tau \mid \{w_1, w_2\} = (1, 1)$ . We claim that

$$\sum_i \left( \sum_j a_{ij} + a_{ii} + m_{ii} - 1 \right) \leq 1$$

If not, we have the following subquiver  $\mathbb{B}$  in  $Q_\tau^\bullet$  :



and we would have in  $\mathbb{C}[rep_{\alpha_\tau} Q_\tau^\bullet]^{GL(\alpha_\tau)}$  the invariants  $ux, uy, vx$  and  $vy$  which satisfy the relation  $(ux)(vy) = (uy)(vx)$ . But then, the ring of invariants of  $(Q_\tau^\bullet, \alpha_\tau)$  cannot be a polynomial ring hence neither can that be of  $(Q^\bullet, \alpha)$  by the slice results.

Therefore, at most one of the  $\sum_j a_{ij} + a_{ii} + m_{ii} - 1$  can be equal to one. Let us consider the different possibilities.

(a): There is just one vertex, in which case  $a_{11} + m_{11} - 1 \leq 1$  and we have the one or two loop quiver (one or both of the loops may be marked). If there is one (marked) loop there is no restriction on the dimension vector  $\alpha = (a_1)$ . If there are two (marked) loops, it is well known that the ring of polynomial invariants is a polynomial ring if and only if  $a_1 \leq 2$ , for example by an adaptation of the argument given below.

(b) : There are at least two vertices and all terms  $\sum_j a_{ij} + a_{ii} + m_{ii} - 1 = 0$ . As  $Q^\bullet$  is strongly connected this implies that all  $a_{ii} = m_{ii} = 0$  and for each  $i$  there is a unique arrow to another vertex.

Hence, we are in the case  $Q^\bullet = \tilde{A}_k$  ( $k \geq 2$ ) and there are no restrictions on the dimension vector for the invariants to be a polynomial ring.

(c) : There are at least two vertices and for a unique vertex  $v_i$  the sum  $\sum_i a_{ij} + a_{ii} + m_{ii} - 1 = 1$ , then we are in the case that  $Q^\bullet = \tilde{A}_k(+1)$  with the extra arrow possibly a (marked) loop starting in  $v_i$ .

In this case we still have to prove that  $\min_j a_j \leq 2$ . The case of an extra (marked) loop in  $v_i$  reduces easily to the two loop case treated before. So we may assume that there is a unique  $i' \neq i$  with  $v_{i'}$  the end point of the extra arrow.

First we reduce to the case of  $\tilde{A}_4(+1)$ . Fix the following four vertices :  $z_1 = v_i$ ,  $z_3 = v_{i'}$ ,  $z_2$  the vertex  $v_u$  on the oriented path from  $v_i$  to  $v_{i'}$  where  $a_u$  is minimal and likewise  $z_4$  is the vertex  $v_{u'}$  on the oriented path from  $v_{i'}$  to  $v_i$  where  $a_{u'}$  is minimal.

Observe that degenerate cases are possible if either  $a_i$  or  $a_{i'}$  is minimal but this only simplifies the argument given below. Let  $\beta = (a_i, a_u, a_{i'}, a_{u'})$  then we claim

that


$$\mathbb{C}[\text{rep}_\alpha \tilde{A}_k(+1)]^{GL(\alpha)} \simeq \mathbb{C}[\text{rep}_\beta \tilde{A}_4(+1)]^{GL(\beta)}$$

Indeed, classical invariant theory (see for example [45, Thm. II.4.1]) tells us that

$$(M_{a \times b}(\mathbb{C}) \oplus M_{b \times c}(\mathbb{C})) / GL_b \simeq M_{a \times c}(\mathbb{C})$$

if  $b \geq \min(a, c)$ . Iterating this reduction we obtain the claim.

Therefore, we only have to exclude the special case when  $Q^\bullet$  is of the following form



and the dimension vector  $\beta = (b_1, b_2, b_3, b_4)$  satisfies  $\min b_i \geq 3$ .

Then, one verifies that  $(2, 1, 2, 2)$  and  $(1, 1, 1, 1)$  are dimension vectors of simple representations and by assumption there are semi-simple representations in  $\text{rep}_\beta Q^\bullet$  of type

$$\tau = (1, (2, 1, 2, 2); 1, (1, 1, 1, 1); b_1 - 3, \delta_1; b_2 - 2, \delta_2; b_3 - 3, \delta_3; b_4 - 3, \delta_4)$$

Calculating the local quiver  $Q_\tau^\bullet$  we observe that it again contains a subquiver of the form  $\mathbb{B}$  and we can repeat the argument given above to exclude this case.  $\square$

Clearly, the first case is the hardest to classify. Even when all the dimension components are equal to one, a full classification of the settings where the ring of invariants is a polynomial ring is unknown at the moment.

This result drastically restricts the shapes of the strongly connected components of the quiver  $Q_h$  and of the dimension vector  $\alpha_h$ .

Let  $X_\epsilon$  be the affine variety determined by the center  $Z_\epsilon$  of  $U_\epsilon(\mathfrak{g})$  and let  $\{x_1, \dots, x_l\}$  be the set of points of  $X_\epsilon$  lying over  $h$ . One can prove, see [57, §4] that each  $x_i$  determines a strongly connected component  $\parallel_i$  of  $Q_h$  and that the restriction of  $\alpha_h$  to this component encodes the multiplicity of  $p_i$  over  $h$ . In particular, if  $x_i$  has multiplicity  $n_i$  over  $h$  then all components of  $\alpha_h \upharpoonright \parallel_i$  are multiples of  $n_i$ .

From this one can then show that only the first case of theorem 5.51 can occur and only if the multiplicity of  $p_i$  is equal to one, that is, if  $p_i$  is unramified over  $h$ . Therefore, we have

**THEOREM 5.52.** *With notations as above*

$$\text{Sep } U_\epsilon(\mathfrak{g})/\mathbb{C}[H] \hookrightarrow \text{Sm } U_\epsilon(\mathfrak{g})/\mathbb{C}[H] \hookrightarrow \text{Sm } Z_\epsilon/\mathbb{C}[H]$$

Here,  $\text{Sm } Z_\epsilon/\mathbb{C}[H]$  is the usual smooth locus of the commutative extension  $\mathbb{C}[H] \hookrightarrow Z_\epsilon$ . Therefore, if  $U_\epsilon(\mathfrak{g})$  is smooth in  $h$ , then there are precisely  $l^r$  points of  $X_\epsilon$  lying over  $h$  and  $X_\epsilon$  is smooth in all of them.

K. Brown and K. Goodearl have proved in [10] that the smooth locus of  $X_\epsilon$  coincides with the Azumaya locus of  $U_\epsilon(\mathfrak{g})$  over  $Z_\epsilon$ . But then,  $\bar{U}_h \simeq M_N(\mathbb{C}) \oplus \dots \oplus M_N(\mathbb{C})$  ( $s$  copies) whence  $U_\epsilon(\mathfrak{g})$  is separable over  $\mathbb{C}[H]$  in  $h$ , finishing the (outline) of the proof of theorem 5.50. For more details we refer to [57].

**References.**

The results of section 5.1 are due to L. Le Bruyn and C. Procesi [58]. The results of sections 5.2, 5.3, 5.4 and 5.5 are due to L. Le Bruyn, [55], [50]. Lemma 5.30 and proposition 5.31 are due to M. Artin [3], [4]. The results of section 5.6 are due to W. Crawley-Boevey and M. Holland [16], [14]. Proposition 5.45 is due to W. Crawley-Boevey [15] and theorem 5.47 is due to R. Bockland and L. Le Bruyn [8]. The structural results on quantum groups are due to C. De Concini and C. Procesi [19], the description of the Cayley-smooth locus is due to L. Le Bruyn [57].



## Nullcones.

### 6.1. Cornering matrices.

In this section we will outline the main idea of the Hesselink stratification of the nullcone [30] in the generic case, that is, the action of  $GL_n$  by simultaneous conjugation on  $m$ -tuples of matrices  $M_n^m = M_n \oplus \dots \oplus M_n$ . With  $Null_n^m$  we denote the nullcone of this action

$$Null_n^m = \{x = (A_1, \dots, A_m) \in M_n^m \mid \underline{0} = (0, \dots, 0) \in \overline{\mathcal{O}(x)}\}$$

It follows from the Hilbert criterium 2.17 that  $x = (A_1, \dots, A_m)$  belongs to the nullcone if and only if there is a one-parameter subgroup  $\mathbb{C}^* \xrightarrow{\lambda} GL_n$  such that

$$\lim_{t \rightarrow 0} \lambda(t) \cdot (A_1, \dots, A_m) = (0, \dots, 0).$$

We recall from proposition 2.14 that any one-parameter subgroup of  $GL_n$  is conjugated to one determined by an integral  $n$ -tuple  $(r_1, \dots, r_n) \in \mathbb{Z}^n$  by

$$\lambda(t) = \begin{bmatrix} t^{r_1} & & 0 \\ & \ddots & \\ 0 & & t^{r_n} \end{bmatrix}$$

Moreover, permuting the basis if necessary, we can conjugate this  $\lambda$  to one where the  $n$ -tuple is *dominant*, that is,  $r_1 \geq r_2 \geq \dots \geq r_n$ . By applying *permutation Jordan-moves*, that is, by simultaneously interchanging certain rows and columns in all  $A_i$ , we may therefore assume that the limit-formula holds for a dominant one-parameter subgroup  $\lambda$  of the maximal torus

$$T_n \simeq \underbrace{\mathbb{C}^* \times \dots \times \mathbb{C}^*}_n = \left\{ \begin{bmatrix} c_1 & & 0 \\ & \ddots & \\ 0 & & c_n \end{bmatrix} \mid c_i \in \mathbb{C}^* \right\} \hookrightarrow GL_n$$


of  $GL_n$ . Computing its action on a  $n \times n$  matrix  $A$  we obtain

$$\begin{bmatrix} t^{r_1} & & 0 \\ & \ddots & \\ 0 & & t^{r_n} \end{bmatrix} \begin{bmatrix} a_{11} & \dots & a_{1n} \\ \vdots & & \vdots \\ a_{n1} & \dots & a_{nn} \end{bmatrix} \begin{bmatrix} t^{-r_1} & & 0 \\ & \ddots & \\ 0 & & t^{-r_n} \end{bmatrix} = \begin{bmatrix} t^{r_1-r_1} a_{11} & \dots & t^{r_1-r_n} a_{1n} \\ \vdots & & \vdots \\ t^{r_n-r_1} a_{n1} & \dots & t^{r_n-r_n} a_{nn} \end{bmatrix}$$

But then, using dominance  $r_i \leq r_j$  for  $i \geq j$ , we see that the limit is only defined if  $a_{ij} = 0$  for  $i \geq j$ , that is, when  $A$  is a strictly upper triangular matrix. We have proved the first 'cornering' result.

**LEMMA 6.1.** *Any  $m$ -tuple  $x = (A_1, \dots, A_m) \in Null_n^m$  has a point in its orbit  $\mathcal{O}(x)$  under simultaneous conjugation  $x' = (A'_1, \dots, A'_m)$  with all  $A'_i$  strictly upper triangular matrices. In fact permutation Jordan-moves suffice to arrive at  $x'$ .*

For specific  $m$ -tuples  $x = (A_1, \dots, A_m)$  it might be possible to improve on this result. That is, we want to determine the smallest 'corner'  $C$  in the upper right hand corner of the matrix, such that all the component matrices  $A_i$  can be conjugated simultaneously to matrices  $A'_i$  having only non-zero entries in the corner  $C$



and no strictly smaller corner  $C'$  can be found with this property. Our first task will be to compile a list of the relevant corners and to define an order relation on this set. Consider the *weight space decomposition* of  $M_n^m$  for the action by simultaneous conjugation of the maximal torus  $T_n$ ,

$$M_n^m = \bigoplus_{1 \leq i, j \leq n} M_n^m(\pi_i - \pi_j) = \bigoplus_{1 \leq i, j \leq n} \mathbb{C}^{\oplus m}_{\pi_i - \pi_j}$$

where  $c = \text{diag}(c_1, \dots, c_n) \in T_m$  acts on any element of  $M_n^m(\pi_i - \pi_j)$  by multiplication with  $c_i c_j^{-1}$ , that is, the eigenspace  $M_n^m(\pi_i - \pi_j)$  is the space of the  $(i, j)$ -entries of the  $m$ -matrices. We call

$$\mathcal{W} = \{ \pi_i - \pi_j \mid 1 \leq i, j \leq n \}$$

the set of  $T_n$ -weights of  $M_n^m$ . Let  $x = (A_1, \dots, A_m) \in \text{Null}_n^m$  and consider the subset  $E_x \subset \mathcal{W}$  consisting of the elements  $\pi_i - \pi_j$  such that for at least one of the matrix components  $A_k$  the  $(i, j)$ -entry is non-zero. Repeating the argument above, we see that if  $\lambda$  is a one-parameter subgroup of  $T_n$  determined by the integral  $n$ -tuple  $(r_1, \dots, r_n) \in \mathbb{Z}^n$  such that  $\lim \lambda(t).x = \underline{0}$  we have

$$\forall \pi_i - \pi_j \in E_x \quad \text{we have} \quad r_i - r_j \geq 1$$

Conversely, let  $E \subset \mathcal{W}$  be a subset of weights, we want to determine the subset

$$\{ s = (s_1, \dots, s_n) \in \mathbb{R}^n \mid s_i - s_j \geq 1 \ \forall \pi_i - \pi_j \in E \}$$

and determine a point in this set, minimal with respect to the usual norm

$$\| s \| = \sqrt{s_1^2 + \dots + s_n^2}$$

Let  $s = (s_1, \dots, s_n)$  attain such a minimum. We can partition the entries of  $s$  in a disjoint union of *strings*

$$\{ p_i, p_i + 1, \dots, p_i + k_i \}$$

with  $k_i \in \mathbb{N}$  and subject to the condition that all the numbers  $p_{ij} \stackrel{\text{def}}{=} p_i + j$  with  $0 \leq j \leq k_i$  occur as components of  $s$ , possibly with a multiplicity that we denote by  $a_{ij}$ . We call a string  $\text{string}_i = \{ p_i, p_i + 1, \dots, p_i + k_i \}$  of  $s$  *balanced* if and only if

$$\sum_{s_k \in \text{string}_i} s_j = \sum_{j=0}^{k_i} a_{ij}(p_i + j) = 0$$

In particular, all balanced strings consists entirely of rational numbers. We have



LEMMA 6.2. *Let  $E \subset \mathcal{W}$ , then the subset of  $\mathbb{R}^n$  determined by*

$$\mathbb{R}_E^n = \{ (r_1, \dots, r_n) \mid r_i - r_j \geq 1 \ \forall \pi_i - \pi_j \in E \}$$

*has a unique point  $s_E = (s_1, \dots, s_n)$  of minimal norm  $\|s_E\|$ . This point is determined by the characteristic feature that all its strings are balanced. In particular,  $s_E \in \mathbb{Q}^n$ .*

PROOF. Let  $s$  be a minimal point for the norm in  $\mathbb{R}_E^n$  and consider a string of  $s$  and denote with  $S$  the indices  $k \in \{1, \dots, n\}$  such that  $s_k \in \text{string}$ . Let  $\pi_i - \pi_j \in E$ , then if only one of  $i$  or  $j$  belongs to  $S$  we have a strictly positive number  $a_{ij}$

$$s_i - s_j = 1 + r_{ij} \quad \text{with} \quad r_{ij} > 0$$

Take  $\epsilon_0 > 0$  smaller than all  $r_{ij}$  and consider the  $n$ -tuple

$$s_\epsilon = s + \epsilon(\delta_{1S}, \dots, \delta_{nS}) \quad \text{with} \quad \delta_{kS} = 1 \text{ if } k \in S \text{ and } 0 \text{ otherwise}$$

with  $|\epsilon| \leq \epsilon_0$ . Then,  $s_\epsilon \in \mathbb{R}_E^n$  for if  $\pi_i - \pi_j \in E$  and  $i$  and  $j$  both belong to  $S$  or both do not belong to  $S$  then  $(s_\epsilon)_i - (s_\epsilon)_j = s_i - s_j \geq 1$  and if one of  $i$  or  $j$  belong to  $S$ , then

$$(s_\epsilon)_i - (s_\epsilon)_j = 1 + r_{ij} \pm \epsilon \geq 1$$

by the choice of  $\epsilon_0$ . However, the norm of  $s_\epsilon$  is

$$\|s_\epsilon\| = \sqrt{\|s\|^2 + 2\epsilon \sum_{k \in S} s_k + \epsilon^2 \#S}$$

Hence, if the string would not be balanced,  $\sum_{k \in S} s_k \neq 0$  and we can choose  $\epsilon$  small enough such that  $\|s_\epsilon\| < \|s\|$ , contradicting minimality of  $s$ .  $\square$

For given  $n$  we have the following **algorithm** to compile the list  $\mathcal{S}_n$  of all dominant  $n$ -tuples  $(s_1, \dots, s_n)$  (that is,  $s_i \leq s_j$  whenever  $i \geq j$ ) having all its strings balanced.

- List all Young-diagrams  $\mathcal{Y}_n = \{Y_1, \dots\}$  having  $\leq n$  boxes.
- For every diagram  $Y_l$  fill the boxes with strictly positive integers subject to the rules
  - (1) the total sum is equal to  $n$
  - (2) no two rows are filled identically
  - (3) at most one row has length 1

This gives a list  $\mathcal{T}_n = \{T_1, \dots\}$  of tableaux.

- For every tableau  $T_l \in \mathcal{T}_n$ , for each of its rows  $(a_1, a_2, \dots, a_k)$  find a solution  $p$  to the linear equation

$$a_1x + a_2(x+1) + \dots + a_k(x+k) = 0$$

and define the  $\sum a_i$ -tuple of rational numbers

$$\underbrace{(p, \dots, p)}_{a_1}, \underbrace{(p+1, \dots, p+1)}_{a_2}, \dots, \underbrace{(p+k, \dots, p+k)}_{a_k}$$

Repeating this process for every row of  $T_l$  we obtain an  $n$ -tuple, which we then order.

The list  $\mathcal{S}_n$  will be the combinatorial object underlying the relevant corners and the stratification of the nullcone.

EXAMPLE 6.3. ( $\mathcal{S}_n$  for small  $n$ )

For  $n = 2$ , we have  $\boxed{1 \ 1}$  giving  $(\frac{1}{2}, -\frac{1}{2})$  and  $\boxed{2}$  giving  $(0, 0)$ . For  $n = 3$  we have five types

	tableau	$s_1$	$s_2$	$s_3$	$\ s\ ^2$
$\mathcal{S}_3 =$	$\boxed{1 \ 1 \ 1}$	1	0	-1	2
	$\boxed{1 \ 2}$	$\frac{1}{3}$	$\frac{1}{3}$	$-\frac{2}{3}$	$\frac{2}{3}$
	$\boxed{2 \ 1}$	$\frac{2}{3}$	$-\frac{1}{3}$	$-\frac{1}{3}$	$\frac{2}{3}$
	$\boxed{1 \ 1}$	1	0	-1	2
	$\boxed{3}$	0	0	0	0

$\mathcal{S}_4$  has eleven types

	tableau	$s_1$	$s_2$	$s_3$	$s_4$	$\ s\ ^2$
$\mathcal{S}_4 =$	$\boxed{1 \ 1 \ 1 \ 1}$	$\frac{3}{2}$	$\frac{1}{2}$	$-\frac{1}{2}$	$-\frac{3}{2}$	5
	$\boxed{2 \ 1 \ 1}$	$\frac{5}{4}$	$\frac{1}{4}$	$-\frac{3}{4}$	$-\frac{3}{4}$	$\frac{11}{4}$
	$\boxed{1 \ 1 \ 2}$	$\frac{3}{4}$	$\frac{3}{4}$	$-\frac{1}{4}$	$-\frac{5}{4}$	$\frac{11}{4}$
	$\boxed{1 \ 2 \ 1}$	1	0	0	-1	2
	$\boxed{2 \ 2}$	$\frac{1}{2}$	$\frac{1}{2}$	$-\frac{1}{2}$	$-\frac{1}{2}$	1
	$\boxed{3 \ 1}$	$\frac{3}{4}$	$-\frac{1}{4}$	$-\frac{1}{4}$	$-\frac{1}{4}$	$\frac{3}{4}$
	$\boxed{1 \ 3}$	$\frac{1}{4}$	$\frac{1}{4}$	$\frac{1}{4}$	$-\frac{3}{4}$	$\frac{3}{4}$
	$\boxed{1 \ 2}$	$\frac{1}{3}$	$\frac{1}{3}$	0	$-\frac{2}{3}$	$\frac{2}{3}$
	$\boxed{2 \ 1}$	$\frac{2}{3}$	0	$-\frac{1}{3}$	$-\frac{1}{3}$	$\frac{2}{3}$
	$\boxed{1 \ 1}$	1	0	0	-1	2
	$\boxed{4}$	0	0	0	0	0

Observe that we ordered the elements in  $\mathcal{S}_n$  according to  $\|s\|$ . The reader is invited to verify that  $\mathcal{S}_5$  has 28 different types.


To every  $s = (s_1, \dots, s_n) \in \mathcal{S}_n$  we associate the following data


- the *corner*  $C_s$  is the subspace of  $M_n^m$  consisting of those  $m$  tuples of  $n \times n$  matrices with zero entries except perhaps at position  $(i, j)$  where  $s_i - s_j \geq 1$ . A partial ordering is defined on these corners by the rule

$$C_{s'} < C_s \Leftrightarrow \|s'\| < \|s\|$$


- the *parabolic subgroup*  $P_s$  which is the subgroup of  $GL_n$  consisting of matrices with zero entries except perhaps at entry  $(i, j)$  when  $s_i - s_j \geq 0$ .
- the *Levi subgroup*  $L_s$  which is the subgroup of  $GL_n$  consisting of matrices with zero entries except perhaps at entry  $(i, j)$  when  $s_i - s_j = 0$ . Observe that  $L_s = \prod GL_{a_{ij}}$  where the  $a_{ij}$  are the multiplicities of  $p_i + j$ .

EXAMPLE 6.4. Using the sequence of types in the previous example, we have that the relevant corners and subgroup for  $3 \times 3$  matrices are





For  $4 \times 4$  matrices the relevant corners are



Returning to the corner-type of an  $m$ -tuple  $x = (A_1, \dots, A_m) \in Null_n^m$ , we have seen that  $E_x \subset \mathcal{W}$  determines a unique  $s_{E_x} \in \mathbb{Q}^n$  which up to permuting the entries an element  $s$  of  $\mathcal{S}_n$ . As permuting the entries of  $s$  translates into permuting rows and columns in  $M_n(\mathbb{C})$  we have

**THEOREM 6.5.** *Every  $x = (A_1, \dots, A_m) \in Null_n^m$  can be brought by permutation Jordan-moves to an  $m$ -tuple  $x' = (A'_1, \dots, A'_m) \in C_s$ . Here,  $s$  is the dominant reordering of  $s_{E_x}$  with  $E_x \subset \mathcal{W}$  the subset  $\pi_i - \pi_j$  determined by the non-zero entries at place  $(i, j)$  of one of the components  $A_k$ . The permutation of rows and columns is determined by the dominant reordering.*

The  $m$ -tuple  $s$  (or  $s_{E_x}$ ) determines a one-parameter subgroup  $\lambda_s$  of  $T_n$  where  $\lambda$  corresponds to the unique  $n$ -tuple of integers

$$(r_1, \dots, r_n) \in \mathbb{N}_+ s \cap \mathbb{Z}^n \quad \text{with} \quad \gcd(r_i) = 1$$

For any one-parameter subgroup  $\mu$  of  $T_n$  determined by an integral  $n$ -tuple  $\mu = (a_1, \dots, a_n) \in \mathbb{Z}^n$  and any  $x = (A_1, \dots, A_m) \in Null_n^m$  we define the integer

$$m(x, \mu) = \min \{a_i - a_j \mid x \text{ contains a non-zero entry in } M_n^m(\pi_i - \pi_j)\}$$

From the definition of  $\mathbb{R}_E^n$  it follows that the minimal value  $s_E$  and  $\lambda_{s_E}$  is

$$s_{E_x} = \frac{\lambda_{s_{E_x}}}{m(x, \lambda_{s_{E_x}})} \quad \text{and} \quad s = \frac{\lambda_s}{m(x, \lambda_s)}$$

We can now state to what extend  $\lambda_s$  is an optimal one-parameter subgroup of  $T_n$ .

**THEOREM 6.6.** *Let  $x = (A_1, \dots, A_m) \in Null_n^m$  and let  $\mu$  be a one-parameter subgroup contained in  $T_n$  such that  $\lim_{t \rightarrow 0} \lambda(t).x = \underline{0}$ , then*

$$\frac{\|\lambda_{s_{E_x}}\|}{m(x, \lambda_{s_{E_x}})} \leq \frac{\|\mu\|}{m(x, \mu)}$$

The proof follows immediately from the observation that  $\frac{\mu}{m(x, \mu)} \in \mathbb{R}_{E_x}^n$  and the minimality of  $s_{E_x}$ . Phrased differently, there is no simultaneous reordering of rows and columns that admit an  $m$ -tuple  $x'' = (A''_1, \dots, A''_m) \in C_{s'}$  for a corner  $C_{s'} < C_s$ . In section 4 we will improve this result.

**6.2. General subrepresentations.**

In the study of the Hesselink stratification of the nullcone and several moduli space problems, it is crucial to determine the dimension vectors of *general subrepresentations*. It follows from theorem 4.49 that this problem is equivalent to the calculation of  $ext(\alpha, \beta)$ . An inductive algorithm to do this was discovered by A. Schofield [70].

Recall that  $\alpha \hookrightarrow \beta$  iff a general representation  $W \in rep_\beta Q$  contains a subrepresentation  $S \hookrightarrow W$  of dimension vector  $\alpha$ . Similarly, we denote  $\beta \twoheadrightarrow \gamma$  if and only if a general representation  $W \in rep_\beta Q$  has a quotient-representation  $W \twoheadrightarrow T$  of dimension vector  $\gamma$ . As before,  $Q$  will be a quiver on  $k$ -vertices  $\{v_1, \dots, v_k\}$  and we denote dimension vectors  $\alpha = (a_1, \dots, a_k)$ ,  $\beta = (b_1, \dots, b_k)$  and  $\gamma = (c_1, \dots, c_k)$ . We will first determine the *rank of a general homomorphism*  $V \twoheadrightarrow W$  between representations  $V \in rep_\alpha Q$  and  $W \in rep_\beta Q$ . We denote

$$Hom(\alpha, \beta) = \bigoplus_{i=1}^k M_{b_i \times a_i} \quad \text{and} \quad Hom(V, \beta) = Hom(\alpha, \beta) = Hom(\alpha, W)$$

for any representations  $V$  and  $W$  as above. With these conventions we have

LEMMA 6.7. *There is an open subset  $Hom_m(\alpha, \beta) \hookrightarrow rep_\alpha Q \times rep_\beta Q$  and a dimension vector  $\gamma \stackrel{def}{=} rk \text{ hom}(\alpha, \beta)$  such that for all  $(V, W) \in Hom_{min}(\alpha, \beta)$*

- $dim_{\mathbb{C}} Hom_{\mathbb{C}Q}(V, W)$  is minimal, and
- $\{\phi \in Hom_{\mathbb{C}Q}(V, W) \mid rk \phi = \gamma\}$  is a non-empty Zariski open subset of  $Hom_{\mathbb{C}Q}(V, W)$ .

PROOF. Consider the subvariety  $Hom_Q(\alpha, \beta)$  of the trivial vectorbundle

$$\begin{array}{ccc} Hom_Q(\alpha, \beta) & \hookrightarrow & Hom(\alpha, \beta) \times rep_\alpha Q \times rep_\beta Q \\ & \searrow \phi & \downarrow pr \\ & & rep_\alpha Q \times rep_\beta Q \end{array}$$

of triples  $(\phi, V, W)$  such that  $V \xrightarrow{\phi} W$  is a morphism of representations of  $Q$ . The fiber  $\Phi^{-1}(V, W) = Hom_{\mathbb{C}Q}(V, W)$ . As the fiber dimension is upper semi-continuous, there is an open subset  $Hom_{min}(\alpha, \beta)$  of  $rep_\alpha Q \times rep_\beta Q$  consisting of points  $(V, W)$  where  $dim_{\mathbb{C}} Hom_{\mathbb{C}Q}(V, W)$  is minimal. For given dimension vector  $\delta = (d_1, \dots, d_k)$  we consider the subset

$$Hom_Q(\alpha, \beta, \delta) = \{(\phi, V, W) \in Hom_Q(\alpha, \beta) \mid rk \phi = \delta\} \hookrightarrow Hom_Q(\alpha, \beta)$$

This is a constructible subset of  $Hom_Q(\alpha, \beta)$  and hence there is a dimension vector  $\gamma$  such that  $Hom_Q(\alpha, \beta, \gamma) \cap \Phi^{-1}(Hom_{min}(\alpha, \beta))$  is constructible and dense in  $\Phi^{-1}(Hom_{min}(\alpha, \beta))$ . But then,

$$\Phi(Hom_Q(\alpha, \beta, \gamma) \cap \Phi^{-1}(Hom_{min}(\alpha, \beta)))$$

is constructible and dense in  $Hom_{min}(V, W)$ . Therefore it contains an open subset  $Hom_m(V, W)$  satisfying the requirements of the lemma.  $\square$

LEMMA 6.8. *Assume we have short exact sequences of representations of  $Q$*

$$\left\{ \begin{array}{l} 0 \longrightarrow S \longrightarrow V \longrightarrow X \longrightarrow 0 \\ 0 \longrightarrow Y \longrightarrow W \longrightarrow T \longrightarrow 0 \end{array} \right.$$

then there is a natural onto map

$$\text{Ext}_{\mathbb{C}Q}^1(V, W) \twoheadrightarrow \text{Ext}_{\mathbb{C}Q}^1(S, T)$$

PROOF. By lemma 4.40 we have surjective maps

$$\text{Ext}_{\mathbb{C}Q}^1(V, W) \twoheadrightarrow \text{Ext}_{\mathbb{C}Q}^1(V, T) \twoheadrightarrow \text{Ext}_{\mathbb{C}Q}^1(S, T)$$

from which the assertion follows.  $\square$

THEOREM 6.9. Let  $\gamma = rk \text{ hom}(\alpha, \beta)$  (with notations as in lemma 6.7), then

- (1)  $\alpha - \gamma \hookrightarrow \alpha \twoheadrightarrow \gamma \hookrightarrow \beta \twoheadrightarrow \beta - \gamma$
- (2)  $\text{ext}(\alpha, \beta) = -\chi_Q(\alpha - \gamma, \beta - \gamma) = \text{ext}(\alpha - \gamma, \beta - \gamma)$

PROOF. The first statement is obvious from the definitions, for if  $\gamma = rk \text{ hom}(\alpha, \beta)$ , then a general representation of dimension  $\alpha$  will have a quotient-representation of dimension  $\gamma$  (and hence a subrepresentation of dimension  $\alpha - \gamma$ ) and a general representation of dimension  $\beta$  will have a subrepresentation of dimension  $\gamma$  (and hence a quotient-representation of dimension  $\beta - \gamma$ ).

The strategy of the proof of the second statement is to compute the dimension of the subvariety of  $\text{Hom}(\alpha, \beta) \times rep_\alpha \times rep_\beta \times rep_\gamma$  defined by

$$H^{factor} = \{(\phi, V, W, X) \mid \begin{array}{ccc} V & \xrightarrow{\phi} & W \\ & \searrow & \nearrow \\ & X = Im \phi & \end{array} \text{ factors as representations} \}$$

in two different ways. Consider the intersection of the open set  $\text{Hom}_m(\alpha, \beta)$  determined by lemma 6.7 with the open set of couples  $(V, W)$  such that  $\dim \text{Ext}(V, W) = \text{ext}(\alpha, \beta)$  and let  $(V, W)$  lie in this intersection. In the previous section we have proved that

$$\dim \underline{Grass}_\gamma(W) = \chi_Q(\gamma, \beta - \gamma)$$

Let  $H$  be the subbundle of the trivial vectorbundle over  $\underline{Grass}_\gamma(W)$

$$\begin{array}{ccc} H & \hookrightarrow & \text{Hom}(\alpha, W) \times \underline{Grass}_\gamma(W) \\ & \searrow & \downarrow \\ & & \underline{Grass}_\gamma(W) \end{array}$$

consisting of triples  $(\phi, W, U)$  with  $\phi : \oplus_i \mathbb{C}^{a_i} \longrightarrow W$  a linear map such that  $Im(\phi)$  is contained in the subrepresentation  $U \hookrightarrow W$  of dimension  $\gamma$ . That is, the fiber over  $(W, U)$  is  $\text{Hom}(\alpha, U)$  and therefore has dimension  $\sum_{i=1}^k a_i c_i$ . With  $H^{full}$  we consider the open subvariety of  $H$  of triples  $(\phi, W, U)$  such that  $Im \phi = U$ . We have

$$\dim H^{full} = \sum_{i=1}^k a_i c_i + \chi_Q(\gamma, \beta - \gamma)$$

But then,  $H^{factor}$  is the subbundle of the trivial vectorbundle over  $H^{full}$

$$\begin{array}{ccc} H^{factor} & \hookrightarrow & rep_{\alpha} Q \times H^{full} \\ & \searrow \pi & \downarrow \\ & & H^{full} \end{array}$$

consisting of quadruples  $(V, \phi, W, X)$  such that  $V \xrightarrow{\phi} W$  is a morphism of representations, with image the subrepresentation  $X$  of dimension  $\gamma$ . The fiber of  $\pi$  over a triple  $(\phi, W, X)$  is determined by the property that for each arrow  $\textcircled{j} \xleftarrow{a} \textcircled{i}$  the following diagram must be commutative, where we decompose the vertex spaces  $V_i = X_i \oplus K_i$  for  $K = Ker \phi$

$$\begin{array}{ccc} X_i \oplus K_i & \xrightarrow{\begin{bmatrix} A & B \\ C & D \end{bmatrix}} & X_j \oplus K_j \\ \downarrow \begin{bmatrix} \mathbb{1}_{c_i} & 0 \end{bmatrix} & & \downarrow \begin{bmatrix} \mathbb{1}_{c_j} & 0 \end{bmatrix} \\ X_i & \xrightarrow{A} & X_j \end{array}$$

where  $A$  is fixed, giving the condition  $B = 0$  and hence the fiber has dimension equal to

$$\sum_{\textcircled{j} \xleftarrow{a} \textcircled{i}} (a_i - c_i)(a_j - c_j) + \sum_{\textcircled{j} \xleftarrow{a} \textcircled{i}} c_i(a_j - c_j) = \sum_{\textcircled{j} \xleftarrow{a} \textcircled{i}} a_i(a_j - c_j)$$

This gives our first formula for the dimension of  $H^{factor}$

$$H^{factor} = \sum_{i=1}^k a_i c_i + \chi_Q(\gamma, \beta - \gamma) + \sum_{\textcircled{j} \xleftarrow{a} \textcircled{i}} a_i(a_j - c_j)$$

On the other hand, we can consider the natural map  $H^{factor} \xrightarrow{\Phi} rep_{\alpha} Q$  defined by sending a quadruple  $(V, \phi, W, X)$  to  $V$ . the fiber in  $V$  is given by all quadruples  $(V, \phi, W, X)$  such that  $V \xrightarrow{\phi} W$  is a morphism of representations with  $Im \phi = X$  a representation of dimension vector  $\gamma$ , or equivalently

$$\Phi^{-1}(V) = \{V \xrightarrow{\phi} W \mid rk \phi = \gamma\}$$

Now, recall our restriction on the couple  $(V, W)$  giving at the beginning of the proof. There is an open subset  $max$  of  $rep_{\alpha} Q$  of such  $V$  and by construction  $max \hookrightarrow Im \Phi$ ,  $\Phi^{-1}(max)$  is open and dense in  $H^{factor}$  and the fiber  $\Phi^{-1}(V)$  is open and dense in  $Hom_{\mathbb{C}Q}(V, W)$ . This provides us with the second formula for the dimension of  $H^{factor}$

$$dim H^{factor} = dim rep_{\alpha} Q + hom(\alpha, W) = \sum_{\textcircled{j} \xleftarrow{a} \textcircled{i}} a_i a_j + hom(\alpha, \beta)$$

Equating both formulas we obtain the equality

$$\chi_Q(\gamma, \beta - \gamma) + \sum_{i=1}^k a_i c_i - \sum_{\substack{j \leftarrow \alpha \\ \textcircled{j} \leftarrow \textcircled{i}}} a_j c_j = \text{hom}(\alpha, \beta)$$

which is equivalent to

$$\chi_Q(\gamma, \beta - \gamma) + \chi_Q(\alpha, \gamma) - \chi_Q(\alpha, \beta) = \text{ext}(\alpha, \beta)$$

Now, for our  $(V, W)$  we have that  $\text{Ext}(V, W) = \text{ext}(\alpha, \beta)$  and we have exact sequences of representations

$$0 \longrightarrow S \longrightarrow V \longrightarrow X \longrightarrow 0 \quad 0 \longrightarrow X \longrightarrow W \longrightarrow T \longrightarrow 0$$

and using lemma 6.8 this gives a surjection  $\text{Ext}(V, W) \twoheadrightarrow \text{Ext}(S, T)$ . On the other hand we always have from the homological interpretation of the Euler form the first inequality

$$\begin{aligned} \dim_{\mathbb{C}} \text{Ext}(S, T) &\geq -\chi_Q(\alpha - \gamma, \beta - \gamma) = \chi_Q(\gamma, \beta - \gamma) - \chi_Q(\alpha, \beta) + \chi_Q(\alpha, \gamma) \\ &= \text{ext}(\alpha, \beta) \end{aligned}$$

As the last term is  $\dim_{\mathbb{C}} \text{Ext}(V, W)$ , this implies that the above surjection must be an isomorphism and that

$$\dim_{\mathbb{C}} \text{Ext}(S, T) = -\chi_Q(\alpha - \gamma, \beta - \gamma) \quad \text{whence} \quad \dim_{\mathbb{C}} \text{Hom}(S, T) = 0$$

But this implies that  $\text{hom}(\alpha - \gamma, \beta - \gamma) = 0$  and therefore  $\text{ext}(\alpha - \gamma, \beta - \gamma) = -\chi_Q(\alpha - \gamma, \beta - \gamma)$ . Finally,

$$\text{ext}(\alpha - \gamma, \beta - \gamma) = \dim \text{Ext}(S, T) = \dim \text{Ext}(V, W) = \text{ext}(\alpha, \beta)$$

finishing the proof. □

**THEOREM 6.10.** *For all dimension vectors  $\alpha$  and  $\beta$  we have*

$$\begin{aligned} \text{ext}(\alpha, \beta) &= \max_{\substack{\alpha' \hookrightarrow \alpha \\ \beta \twoheadrightarrow \beta'}} -\chi_Q(\alpha', \beta') \\ &= \max_{\beta \twoheadrightarrow \beta''} -\chi_Q(\alpha, \beta'') \\ &= \max_{\alpha'' \hookrightarrow \alpha} -\chi_Q(\alpha'', \beta) \end{aligned}$$

**PROOF.** Let  $V$  and  $W$  be representation of dimension vector  $\alpha$  and  $\beta$  such that  $\dim \text{Ext}(V, W) = \text{ext}(\alpha, \beta)$ . Let  $S \hookrightarrow V$  be a subrepresentation of dimension  $\alpha'$  and  $W \twoheadrightarrow T$  a quotient representation of dimension vector  $\beta'$ . Then, we have

$$\text{ext}(\alpha, \beta) = \dim_{\mathbb{C}} \text{Ext}(V, W) \geq \dim_{\mathbb{C}} \text{Ext}(S, T) \geq -\chi_Q(\alpha', \beta')$$

where the first inequality is lemma 6.8 and the second follows from the interpretation of the Euler form. Therefore,  $\text{ext}(\alpha, \beta)$  is greater or equal than all the terms in the statement of the theorem. The foregoing theorem asserts the first equality, as for  $\text{rk hom}(\alpha, \beta) = \gamma$  we do have that  $\text{ext}(\alpha, \beta) = -\chi_Q(\alpha - \gamma, \beta - \gamma)$ .

In the proof of the above theorem, we have found for sufficiently general  $V$  and  $W$  an exact sequence of representations

$$0 \longrightarrow S \longrightarrow V \longrightarrow W \longrightarrow T \longrightarrow 0$$

where  $S$  is of dimension  $\alpha - \gamma$  and  $T$  of dimension  $\beta - \gamma$ . Moreover, we have a commuting diagram of surjections

$$\begin{array}{ccc} \text{Ext}(V, W) & \twoheadrightarrow & \text{Ext}(V, T) \\ \downarrow & \dashrightarrow & \downarrow \\ \text{Ext}(S, W) & \twoheadrightarrow & \text{Ext}(S, T) \end{array}$$

and the dashed map is an isomorphism, hence so are all the epimorphisms. Therefore, we have

$$\begin{cases} \text{ext}(\alpha, \beta - \gamma) \leq \dim \text{Ext}(V, T) = \dim \text{Ext}(V, W) = \text{ext}(\alpha, \beta) \\ \text{ext}(\alpha - \gamma, \beta) \leq \dim \text{Ext}(S, W) = \dim \text{Ext}(V, W) = \text{ext}(\alpha, \beta) \end{cases}$$

Further, let  $T'$  be a sufficiently general representation of dimension  $\beta - \gamma$ , then it follows from  $\text{Ext}(V, T') \twoheadrightarrow \text{Ext}(S, T)$  that

$$\text{ext}(\alpha - \gamma, \beta - \gamma) \leq \dim \text{Ext}(S, T') \leq \dim \text{Ext}(V, T') = \text{ext}(\alpha, \beta - \gamma)$$

but the left term is equal to  $\text{ext}(\alpha, \beta)$  by the above theorem. But then, we have  $\text{ext}(\alpha, \beta) = \text{ext}(\alpha, \beta - \gamma)$ . Now, we may assume by induction that the theorem holds for  $\beta - \gamma$ . That is, there exists  $\beta - \gamma \twoheadrightarrow \beta''$  such that  $\text{ext}(\alpha, \beta - \gamma) = -\chi_Q(\alpha, \beta'')$ . Whence,  $\beta \twoheadrightarrow \beta''$  and  $\text{ext}(\alpha, \beta) = -\chi_Q(\alpha, \beta'')$  and the middle equality of the theorem holds. By a dual argument so does the last.  $\square$

This gives us the following inductive **algorithm** to find all the dimension vectors of general subrepresentations. Take a dimension vector  $\alpha$  and assume by induction we know for all  $\beta < \alpha$  the set of general subrepresentations  $\beta' \hookrightarrow \beta$ . Then,  $\beta \hookrightarrow \alpha$  if and only if

$$0 = \text{ext}(\beta, \alpha - \beta) = \max_{\beta' \hookrightarrow \beta} -\chi_Q(\beta', \alpha - \beta)$$

where the first equality comes from theorem 4.49 and the last from the above theorem.

### 6.3. Semistable representations.

Let  $Q$  be a quiver on  $k$  vertices  $\{v_1, \dots, v_k\}$  and fix a dimension vector  $\alpha$ . So far, we have considered the algebraic quotient map

$$\text{rep}_\alpha Q \twoheadrightarrow \text{iss}_\alpha Q$$


classifying closed  $GL(\alpha)$ -orbits in  $\text{rep}_\alpha Q$ , that is, isomorphism classes of semi-simple representations of dimension  $\alpha$ . We have seen that the invariant polynomial maps are generated by traces along oriented cycles in the quiver. Hence, if  $Q$  has no oriented cycles, the quotient variety  $\text{iss}_\alpha Q$  is reduced to one point corresponding to the semi-simple

$$S_1^{\oplus a_1} \oplus \dots \oplus S_k^{\oplus a_k}$$

where  $S_i$  is the trivial one-dimensional simple concentrated in vertex  $v_i$ . Still, in these cases one can often classify nice families of representations.



EXAMPLE 6.11. Consider the quiver setting



Then,  $rep_\alpha Q = \mathbb{C}^3$  and the action of  $GL(\alpha) = \mathbb{C}^* \times \mathbb{C}^*$  is given by  $(\lambda, \mu).(x, y, z) = (\frac{\lambda}{\mu}x, \frac{\lambda}{\mu}y, \frac{\lambda}{\mu}z)$ . The only closed  $GL(\alpha)$ -orbit in  $\mathbb{C}^3$  is  $(0, 0, 0)$  as the one-parameter subgroup  $\lambda(t) = (t, 1)$  has the property

$$\lim_{t \rightarrow 0} \lambda(t).(x, y, z) = (0, 0, 0)$$

so  $(0, 0, 0) \in \overline{\mathcal{O}(x, y, z)}$  for any representation  $(x, y, z)$ . Still, if we throw away the zero-representation, then we have a nice quotient map

$$\mathbb{C}^3 - \{(0, 0, 0)\} \xrightarrow{\pi} \mathbb{P}^2 \quad (x, y, z) \mapsto [x : y : z]$$

and as  $\mathcal{O}(x, y, z) = \mathbb{C}^*(x, y, z)$  we see that every  $GL(\alpha)$ -orbit is closed in this complement  $\mathbb{C}^3 - \{(0, 0, 0)\}$ . We will generalize such settings to arbitrary quivers.

A *character* of  $GL(\alpha)$  is an algebraic group morphism  $\chi : GL(\alpha) \longrightarrow \mathbb{C}^*$ . They are fully determined by an integral  $k$ -tuple  $\theta = (t_1, \dots, t_k) \in \mathbb{Z}^k$  where

$$GL(\alpha) \xrightarrow{\chi_\theta} \mathbb{C}^* \quad (g_1, \dots, g_k) \mapsto \det(g_1)^{t_1} \dots \det(g_k)^{t_k}$$

For a fixed  $\theta$  we can extend the  $GL(\alpha)$ -action to the space  $rep_\alpha Q \oplus \mathbb{C}$  by

$$GL(\alpha) \times rep_\alpha Q \oplus \mathbb{C} \longrightarrow rep_\alpha Q \oplus \mathbb{C} \quad g.(V, c) = (g.V, \chi_\theta^{-1}(g)c)$$

The coordinate ring  $\mathbb{C}[rep_\alpha Q \oplus \mathbb{C}] = \mathbb{C}[rep_\alpha Q][t]$  can be given a  $\mathbb{Z}$ -gradation by defining  $deg(t) = 1$  and  $deg(f) = 0$  for all  $f \in \mathbb{C}[rep_\alpha Q]$ . The induced action of  $GL(\alpha)$  on  $\mathbb{C}[rep_\alpha Q \oplus \mathbb{C}]$  preserves this gradation. Therefore, the ring of invariant polynomial maps

$$\mathbb{C}[rep_\alpha Q \oplus \mathbb{C}]^{GL(\alpha)} = \mathbb{C}[rep_\alpha Q][t]^{GL(\alpha)}$$

is also graded with homogeneous part of degree zero the ring of invariants  $\mathbb{C}[rep_\alpha Q]^{GL(\alpha)}$ . An invariant of degree  $n$ , say  $ft^n$  with  $f \in \mathbb{C}[rep_\alpha Q]$  has the characteristic property that

$$f(g.V) = \chi_\theta^n(g)f(V)$$

that is,  $f$  is a *semi-invariant* of weight  $\chi_\theta^n$ . That is, the graded decomposition of the invariant ring is

$$\mathbb{C}[rep_\alpha Q \oplus \mathbb{C}]^{GL(\alpha)} = R_0 \oplus R_1 \oplus \dots \quad \text{with} \quad R_i = \mathbb{C}[rep_\alpha Q]^{GL(\alpha), \chi_\theta^{n\theta}}$$

DEFINITION 6.12. With notations as above, the *moduli space of semi-stable quiver representations of dimension  $\alpha$*  is the projective variety

$$M_\alpha^{ss}(Q, \theta) = \text{proj } \mathbb{C}[rep_\alpha Q \oplus \mathbb{C}]^{GL(\alpha)} = \text{proj } \bigoplus_{n=0}^\infty \mathbb{C}[rep_\alpha Q]^{GL(\alpha), \chi_\theta^{n\theta}}$$

Recall that for a positively graded affine commutative  $\mathbb{C}$ -algebra  $R = \bigoplus_{i=0}^\infty R_i$ , the geometric points of the projective scheme  $\text{proj } R$  correspond to graded-maximal ideals  $\mathfrak{m}$  not containing the positive part  $R_+ = \bigoplus_{i=1}^\infty R_i$ . Intersecting  $\mathfrak{m}$  with the part of degree zero  $R_0$  determines a point of  $\text{spec } R_0$ , the affine variety with coordinate ring  $R_0$  and defines a structural morphism


$$\text{proj } R \longrightarrow \text{spec } R_0$$

The *Zariski closed subsets* of  $\text{proj } R$  are of the form

$$\mathbb{V}(I) = \{\mathfrak{m} \in \text{proj } R \mid I \subset \mathfrak{m}\}$$

for a homogeneous ideal  $I \triangleleft R$ . Further, recall that  $\text{proj } R$  can be covered by affine varieties of the form  $\mathbb{X}(f)$  with  $f$  a homogeneous element in  $R_+$ . The coordinate ring of this affine variety is the part of degree zero of the *graded localization*  $R_f^g$ . We refer to [28, II.2] for more details.

EXAMPLE 6.13. Consider again the quiver-situation



and character  $\theta = (-1, 1)$ , then the three coordinate functions  $x, y$  and  $z$  of  $\mathbb{C}[\text{rep}_\alpha Q]$  are semi-invariants of weight  $\chi_\theta$ . It is clear that the invariant ring is equal to

$$\mathbb{C}[\text{rep}_\alpha Q \oplus \mathbb{C}]^{GL(\alpha)} = \mathbb{C}[xt, yt, zt]$$

where the three generators all have degree one. That is,

$$M_\alpha^{ss}(Q, \theta) = \text{Proj } \mathbb{C}[xt, yt, zt] = \mathbb{P}^2$$


as desired,

We will now investigate which orbits in  $\text{rep}_\alpha Q$  are parameterized by the moduli space  $M_\alpha^{ss}(Q, \theta)$ .

DEFINITION 6.14. We say that a representation  $V \in \text{rep}_\alpha Q$  is  $\chi_\theta$ -semistable if and only if there is a semi-invariant  $f \in \mathbb{C}[\text{rep}_\alpha Q]^{GL(\alpha), \chi_\theta^n}$  for some  $n \geq 1$  such that  $f(V) \neq 0$ .

The subset of  $\text{rep}_\alpha Q$  consisting of all  $\chi_\theta$ -semistable representations will be denoted by  $\text{rep}_\alpha^{ss}(Q, \theta)$ .

Observe that  $\text{rep}_\alpha^{ss}(Q, \theta)$  is Zariski open (but it may be empty for certain  $(\alpha, \theta)$ ). We can lift a representation  $V \in \text{rep}_\alpha Q$  to points  $V_c = (V, c) \in \text{rep}_\alpha Q \oplus \mathbb{C}$  and use  $GL(\alpha)$ -invariant theory on this larger  $GL(\alpha)$ -module



Let  $c \neq 0$  and assume that the orbit closure  $\overline{\mathcal{O}(V_c)}$  does not intersect  $\mathbb{V}(t) = \text{rep}_\alpha Q \times \{0\}$ . As both are  $GL(\alpha)$ -stable closed subsets of  $\text{rep}_\alpha Q \oplus \mathbb{C}$  we know from the separation property of invariant theory, proposition 2.32, that this is equivalent to the existence of a  $GL(\alpha)$ -invariant function  $g \in \mathbb{C}[\text{rep}_\alpha Q \oplus \mathbb{C}]^{GL(\alpha)}$  such that  $g(\overline{\mathcal{O}(V_c)}) \neq 0$  but  $g(\mathbb{V}(t)) = 0$ .

We have seen that the invariant ring is graded, hence we may assume  $g$  to be homogeneous, that is, of the form  $g = ft^n$  for some  $n$ . But then,  $f$  is a semi-invariant

on  $\text{rep}_\alpha Q$  of weight  $\chi_\theta^n$  and we see that  $V$  must be  $\chi_\theta$ -semistable. Moreover, we must have that  $\theta(\alpha) = \sum_{i=1}^k t_i a_i = 0$ , for the one-dimensional central torus of  $GL(\alpha)$

$$\mu(t) = (t\mathbb{1}_{a_1}, \dots, t\mathbb{1}_{a_k}) \hookrightarrow GL(\alpha)$$

acts trivially on  $\text{rep}_\alpha Q$  but acts on  $\mathbb{C}$  via multiplication with  $\prod_{i=1}^k t^{-a_i t_i}$  hence if  $\theta(\alpha) \neq 0$  then  $\overline{\mathcal{O}(V_c)} \cap \mathbb{V}(t) \neq \emptyset$ .

More generally, we have from the strong form of the Hilbert criterium proved in theorem 2.17 that  $\overline{\mathcal{O}(V_c)} \cap \mathbb{V}(t) = \emptyset$  if and only if for every one-parameter subgroup  $\lambda(t)$  of  $GL(\alpha)$  we must have that  $\lim_{t \rightarrow 0} \lambda(t).V_c \notin \mathbb{V}(t)$ . We can also formulate this in terms of the  $GL(\alpha)$ -action on  $\text{rep}_\alpha Q$ . The composition of a one-parameter subgroup  $\lambda(t)$  of  $GL(\alpha)$  with the character

$$\mathbb{C}^* \xrightarrow{\lambda(t)} GL(\alpha) \xrightarrow{\chi_\theta} \mathbb{C}^*$$

is an algebraic group morphism and is therefore of the form  $t \longrightarrow t^m$  for some  $m \in \mathbb{Z}$  and we denote this integer by  $\theta(\lambda) = m$ . Assume that  $\lambda(t)$  is a one-parameter subgroup such that  $\lim_{t \rightarrow 0} \lambda(t).V = V'$  exists in  $\text{rep}_\alpha Q$ , then as

$$\lambda(t).(V, c) = (\lambda(t).V, t^{-m}c)$$

we must have that  $\theta(\lambda) \geq 0$  for the orbitclosure  $\overline{\mathcal{O}(V_c)}$  not to intersect  $\mathbb{V}(t)$ .

That is, we have the following characterization of  $\chi_\theta$ -semistable representations.

PROPOSITION 6.15. *The following are equivalent*

- (1)  $V \in \text{rep}_\alpha Q$  is  $\chi_\theta$ -semistable.
- (2) For  $c \neq 0$ , we have  $\overline{\mathcal{O}(V_c)} \cap \mathbb{V}(t) = \emptyset$ .
- (3) For every one-parameter subgroup  $\lambda(t)$  of  $GL(\alpha)$  we have  $\lim_{t \rightarrow 0} \lambda(t).V_c \notin \mathbb{V}(t) = \text{rep}_\alpha Q \times \{0\}$ .
- (4) For every one-parameter subgroup  $\lambda(t)$  of  $GL(\alpha)$  such that  $\lim_{t \rightarrow 0} \lambda(t).V$  exists in  $\text{rep}_\alpha Q$  we have  $\theta(\lambda) \geq 0$ .

Moreover, these cases can only occur if  $\theta(\alpha) = 0$ .

Assume that  $g = ft^n$  is a homogeneous invariant function for the  $GL(\alpha)$ -action on  $\text{rep}_\alpha Q \oplus \mathbb{C}$  and consider the affine open  $GL(\alpha)$ -stable subset  $\mathbb{X}(g)$ . The construction of the algebraic quotient and the fact that the invariant rings here are graded asserts that the closed  $GL(\alpha)$ -orbits in  $\mathbb{X}(g)$  are classified by the points of the graded localization at  $g$  which is of the form

$$(\mathbb{C}[\text{rep}_\alpha Q \oplus \mathbb{C}]^{GL(\alpha)})_g = R_f[h, h^{-1}]$$

for some homogeneous invariant  $h$  and where  $R_f$  is the coordinate ring of the affine open subset  $\mathbb{X}(f)$  in  $M_\alpha^{ss}(Q, \theta)$  determined by the semi-invariant  $f$  of weight  $\chi_\theta^n$ . Because the moduli space is covered by such open subsets we have

PROPOSITION 6.16. *The moduli space of  $\theta$ -semistable representations of  $\text{rep}_\alpha Q$*

$$M_\alpha^{ss}(Q, \theta)$$

*classifies closed  $GL(\alpha)$ -orbits in the open subset  $\text{rep}_\alpha^{ss}(Q, \theta)$  of all  $\chi_\theta$ -semistable representations of  $Q$  of dimension vector  $\alpha$ .*

EXAMPLE 6.17. In the foregoing example  $rep_\alpha^{ss}(Q, \theta) = \mathbb{C}^3 - \{(0, 0, 0)\}$  as for all these points one of the semi-invariant coordinate functions is non-zero. For  $\theta = (-1, 1)$  the lifted  $GL(\alpha) = \mathbb{C}^* \times \mathbb{C}^*$ -action to  $rep_\alpha Q \oplus \mathbb{C} = \mathbb{C}^4$  is given by

$$(\lambda, \mu).(x, y, z, t) = \left(\frac{\mu}{\lambda}x, \frac{\mu}{\lambda}y, \frac{\mu}{\lambda}z, \frac{\lambda}{\mu}t\right)$$

We have seen that the ring of invariants is  $\mathbb{C}[xt, yt, zt]$ . Consider the affine open set  $\mathbb{X}(xt)$  of  $\mathbb{C}^4$ , then the closed orbits in  $\mathbb{X}(xt)$  are classified by

$$\mathbb{C}[xt, yt, zt]_{xt}^g = \mathbb{C}\left[\frac{y}{x}, \frac{z}{x}\right][xt, \frac{1}{xt}]$$

and the part of degree zero  $\mathbb{C}[\frac{y}{x}, \frac{z}{x}]$  is the coordinate ring of the open set  $\mathbb{X}(x)$  in  $\mathbb{P}^2$ .

We have seen that closed  $GL_n$ -orbits in  $rep_n A$  correspond to semi-simple  $n$ -dimensional representations. We will now give a representation theoretic interpretation of closed  $GL(\alpha)$ -orbits in  $rep_\alpha^{ss}(Q, \theta)$ .

Again, the starting point is that one-parameter subgroups  $\lambda(t)$  of  $GL(\alpha)$  correspond to filtrations of representations. Let us go through the motions one more time. For  $\lambda : \mathbb{C}^* \rightarrow GL(\alpha)$  a one-parameter subgroup and  $V \in rep_\alpha Q$  we can decompose for every vertex  $v_i$  the vertex-space in weight spaces

$$V_i = \bigoplus_{n \in \mathbb{Z}} V_i^{(n)}$$

where  $\lambda(t)$  acts on the weight space  $V_i^{(n)}$  as multiplication by  $t^n$ . This decomposition allows us to define a filtration

$$V_i^{(\geq n)} = \bigoplus_{m \geq n} V_i^{(m)}$$

For every arrow  $\textcircled{i} \xleftarrow{a} \textcircled{j}$ ,  $\lambda(t)$  acts on the components of the arrow maps

$$V_i^{(n)} \xrightarrow{V_a^{m,n}} V_j^{(m)}$$

by multiplication with  $t^{m-n}$ . That is, a limit  $\lim_{t \rightarrow 0} V_a$  exists if and only if  $V_a^{m,n} = 0$  for all  $m < n$ , that is, if  $V_a$  induces linear maps

$$V_i^{(\geq n)} \xrightarrow{V_a} V_j^{(\geq n)}$$

Hence, a limiting representation exists if and only if the vertex-filtration spaces  $V_i^{(\geq n)}$  determine a subrepresentation  $V_n \hookrightarrow V$  for all  $n$ . That is, a one-parameter subgroup  $\lambda$  such that  $\lim_{t \rightarrow 0} \lambda(t).V$  exists determines a decreasing filtration of  $V$  by subrepresentations

$$\dots \longleftarrow V_n \longleftarrow V_{n+1} \longleftarrow \dots$$

Further, the limiting representation is then the associated graded representation

$$\lim_{t \rightarrow 0} \lambda(t).V = \bigoplus_{n \in \mathbb{Z}} \frac{V_n}{V_{n+1}}$$

where of course only finitely many of these quotients can be nonzero. For the given character  $\theta = (t_1, \dots, t_k)$  and a representation  $W \in rep_\beta Q$  we denote

$$\theta(W) = t_1 b_1 + \dots + t_k b_k \quad \text{where } \beta = (b_1, \dots, b_k)$$

Assume that  $\theta(V) = 0$ , then with the above notations, we have an interpretation of  $\theta(\lambda)$  as

$$\theta(\lambda) = \sum_{i=1}^k t_i \sum_{n \in \mathbb{Z}} \dim_{\mathbb{C}} V_i^{(n)} = \sum_{n \in \mathbb{Z}} n \theta\left(\frac{V_n}{V_{n+1}}\right) = \sum_{n \in \mathbb{Z}} \theta(V_n)$$

DEFINITION 6.18. A representation  $V \in \text{rep}_\alpha Q$  is said to be

- $\theta$ -semistable if  $\theta(V) = 0$  and for all subrepresentations  $W \hookrightarrow V$  we have  $\theta(W) \geq 0$ .
- $\theta$ -stable if  $V$  is  $\theta$ -semistable and if the only subrepresentations  $W \hookrightarrow V$  such that  $\theta(W) = 0$  are  $V$  and  $0$ .

PROPOSITION 6.19. For  $V \in \text{rep}_\alpha Q$  the following are equivalent

- (1)  $V$  is  $\chi_\theta$ -semistable.
- (2)  $V$  is  $\theta$ -semistable.

PROOF. (1)  $\Rightarrow$  (2) : Let  $W$  be a subrepresentation of  $V$  and let  $\lambda$  be the one-parameter subgroup associated to the filtration  $V \longleftarrow W \longleftarrow 0$ , then  $\lim_{t \rightarrow 0} \lambda(t).V$  exists whence by proposition 6.15.4 we have  $\theta(\lambda) \geq 0$ , but we have

$$\theta(\lambda) = \theta(V) + \theta(W) = \theta(W)$$

(2)  $\Rightarrow$  (1) : Let  $\lambda$  be a one-parameter subgroup of  $GL(\alpha)$  such that  $\lim_{t \rightarrow 0} \lambda(t).V$  exists and consider the induced filtration by subrepresentations  $V_n$  defined above. By assumption all  $\theta(V_n) \geq 0$ , whence

$$\theta(\lambda) = \sum_{n \in \mathbb{Z}} \theta(V_n) \geq 0$$

and again proposition 6.15.4 finishes the proof.  $\square$

LEMMA 6.20. Let  $V \in \text{rep}_\alpha Q$  and  $W \in \text{rep}_\beta Q$  be both  $\theta$ -semistable and

$$V \xrightarrow{f} W$$

a morphism of representations. Then,  $\text{Ker } f$ ,  $\text{Im } f$  and  $\text{Coker } f$  are  $\theta$ -semistable representations.

PROOF. Consider the two short exact sequences of representations of  $Q$

$$\begin{cases} 0 \longrightarrow \text{Ker } f \longrightarrow V \longrightarrow \text{Im } f \longrightarrow 0 \\ 0 \longrightarrow \text{Im } f \longrightarrow W \longrightarrow \text{Coker } f \longrightarrow 0 \end{cases}$$

As  $\theta(-)$  is additive, we have  $0 = \theta(V) = \theta(\text{Ker } f) + \theta(\text{Im } f)$  and as both are subrepresentations of  $\theta$ -semistable representations  $V$  resp.  $W$ , the right-hand terms are  $\geq 0$  whence are zero. But then, from the second sequence also  $\theta(\text{Coker } f) = 0$ . Being submodules of  $\theta$ -semistable representations,  $\text{Ker } f$  and  $\text{Im } f$  also satisfy  $\theta(S) \geq 0$  for all their subrepresentations  $U$ . Finally, a subrepresentation  $T \hookrightarrow \text{Coker } f$  can be lifted to a subrepresentation  $T' \hookrightarrow W$  and  $\theta(T) \geq 0$  follows from the short exact sequence  $0 \longrightarrow \text{Im } f \longrightarrow T' \longrightarrow T \longrightarrow 0$ .  $\square$

That is, the full subcategory  $\text{rep}^{ss}(Q, \theta)$  of  $\text{rep } Q$  consisting of all  $\theta$ -semistable representations is an Abelian subcategory and clearly the simple objects in  $\text{rep}^{ss}(Q, \theta)$  are precisely the  $\theta$ -stable representations. As this Abelian subcategory has the necessary finiteness conditions, one can prove a version of the Jordan-Hölder theorem. That is, every  $\theta$ -semistable representation  $V$  has a finite filtration

$$V = V_0 \longleftarrow V_1 \longleftarrow \dots \longleftarrow V_z = 0$$

of subrepresentation such that every factor  $\frac{V_i}{V_{i+1}}$  is  $\theta$ -stable. Moreover, the unordered set of these  $\theta$ -stable factors are uniquely determined by  $V$ .

**THEOREM 6.21.** *For a  $\theta$ -semistable representation  $V \in \text{rep}_\alpha Q$  the following are equivalent*

- (1) *The orbit  $\mathcal{O}(V)$  is closed in  $\text{rep}_\alpha^{ss}(Q, \alpha)$ .*
- (2)  *$V \simeq W_1^{\oplus e_1} \oplus \dots \oplus W_l^{\oplus e_l}$  with every  $W_i$  a  $\theta$ -stable representation.*

*That is, the geometric points of the moduli space  $M_\alpha^{ss}(Q, \theta)$  are in natural one-to-one correspondence with isomorphism classes of  $\alpha$ -dimensional representations which are direct sums of  $\theta$ -stable subrepresentations. The quotient map*

$$\text{rep}_\alpha^{ss}(Q, \theta) \longrightarrow M_\alpha^{ss}(Q, \theta)$$

*maps a  $\theta$ -semistable representation  $V$  to the direct sum of its Jordan-Hölder factors in the Abelian category  $\text{rep}^{ss}(Q, \theta)$ .*

**PROOF.** Assume that  $\mathcal{O}(V)$  is closed in  $\text{rep}_\alpha^{ss}(Q, \theta)$  and consider the  $\theta$ -semistable representation  $W = \text{gr}_{ss} V$ , the direct sum of the Jordan-Hölder factors in  $\text{rep}^{ss}(Q, \theta)$ . As  $W$  is the associated graded representation of a filtration on  $V$ , there is a one-parameter subgroup  $\lambda$  of  $GL(\alpha)$  such that  $\lim_{t \rightarrow 0} \lambda(t).V \simeq W$ , that is  $\mathcal{O}(W) \subset \overline{\mathcal{O}(V)} = \mathcal{O}(V)$ , whence  $W \simeq V$  and 2. holds.

Conversely, assume that  $V$  is as in 2. and let  $\mathcal{O}(W)$  be a closed orbit contained in  $\overline{\mathcal{O}(V)}$  (one of minimal dimension). By the Hilbert criterium there is a one-parameter subgroup  $\lambda$  in  $GL(\alpha)$  such that  $\lim_{t \rightarrow 0} \lambda(t).V \simeq W$ . Hence, there is a finite filtration of  $V$  with associated graded  $\theta$ -semistable representation  $W$ . As none of the  $\theta$ -stable components of  $V$  admits a proper quotient which is  $\theta$ -semistable (being a direct summand of  $W$ ), this shows that  $V \simeq W$  and so  $\mathcal{O}(V) = \mathcal{O}(W)$  is closed. The other statements are clear from this.  $\square$

Remains to determine the situations  $(\alpha, \theta)$  such that the corresponding moduli space  $M_\alpha^{ss}(Q, \theta)$  is non-empty, or equivalently, such that the Zariski open subset  $\text{rep}_\alpha^{ss}(Q, \theta) \hookrightarrow \text{rep}_\alpha Q$  is non-empty.

**THEOREM 6.22.** *Let  $\alpha$  be a dimension vector such that  $\theta(\alpha) = 0$ . Then,*

- (1)  *$\text{rep}_\alpha^{ss}(Q, \alpha)$  is a non-empty Zariski open subset of  $\text{rep}_\alpha Q$  if and only if for every  $\beta \hookrightarrow \alpha$  we have that  $\theta(\beta) \geq 0$ .*
- (2) *The  $\theta$ -stable representations  $\text{rep}_\alpha^s(Q, \alpha)$  are a non-empty Zariski open subset of  $\text{rep}_\alpha Q$  if and only if for every  $0 \neq \beta \hookrightarrow \alpha$  we have that  $\theta(\beta) > 0$*

The algorithm at the end of the last section gives an inductive procedure to calculate these conditions.

The graded algebra  $\mathbb{C}[\text{rep}_\alpha \oplus \mathbb{C}]^{GL(\alpha)}$  of all semi-invariants on  $\text{rep}_\alpha Q$  of weight  $\chi_\theta^n$  for some  $n \geq 0$  has as degree zero part the ring of polynomial invariants  $\mathbb{C}[\text{rep}_\alpha Q]^{GL(\alpha)}$ . This embedding determines a *proper morphism*

$$M_\alpha^{ss}(Q, \theta) \xrightarrow{\pi} \text{iss}_\alpha Q$$

which is onto whenever  $\text{rep}_\alpha^{ss}(Q, \alpha)$  is non-empty. In particular, if  $Q$  is a quiver without oriented cycles, then the moduli space of  $\theta$ -semistable representations of dimension vector  $\alpha$ ,  $M_\alpha^{ss}(Q, \theta)$ , is a projective variety.

### 6.4. Optimal corners.

We have proved that one can transform an  $m$ -tuple  $x = (A_1, \dots, A_m) \in \text{Null}_n^m$  by interchanging rows and columns to an  $m$ -tuple in corner-form  $C_s$ . However, it

is possible that another point in the orbit  $\mathcal{O}(x)$  say  $y = g.x = (B_1, \dots, B_m)$  can be transformed by permutation Jordan moves in a strictly smaller corner.

EXAMPLE 6.23. Consider one  $3 \times 3$  nilpotent matrix of the form

$$x = \begin{bmatrix} 0 & a & b \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \quad \text{with } ab \neq 0$$

Then,  $E_x = \{\pi_1 - \pi_2, \pi_1 - \pi_3\}$  and the corresponding  $s = s_{E_x} = (\frac{2}{3}, -\frac{1}{3}, -\frac{1}{3})$  so  $x$  is clearly of corner type

$$C_s = \begin{array}{|c|c|c|} \hline & \blacksquare & \blacksquare \\ \hline & & \\ \hline & & \\ \hline \end{array}$$

However,  $x$  is a nilpotent matrix of rank 1 and by the Jordan-normalform we can conjugate it in standard form, that is, there is some  $g \in GL_3$  such that

$$y = g.x = gxg^{-1} = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

For this  $y$  we have  $E_y = \{\pi_1 - \pi_2\}$  and the corresponding  $s_{E_y} = (\frac{1}{2}, -\frac{1}{2}, 0)$ , which can be brought into standard dominant form  $s' = (\frac{1}{2}, 0, -\frac{1}{2})$  by interchanging the two last entries. Hence, by interchanging the last two rows and columns,  $y$  is indeed of corner type

$$C_{s'} = \begin{array}{|c|c|c|} \hline & & \blacksquare \\ \hline & & \\ \hline & & \\ \hline \end{array}$$

and we have that  $C_{s'} < C_s$ .


We required the Jordan-normalform to produce this example. As there are no known canonical forms for  $m$  tuples of  $n \times n$  matrices, it is a difficult problem to determine the optimal corner type in general.

DEFINITION 6.24. We say that  $x = (A_1, \dots, A_m) \in Null_n^m$  is of *optimal corner type*  $C_s$  if after reordering rows and columns,  $x$  is of corner type  $C_s$  and there is no point  $y = g.x$  in the orbit which is of corner type  $C_{s'}$  with  $C_{s'} < C_s$ .

We can give an elegant solution to the problem of determining the optimal corner type of an  $m$ -tuple in  $Null_n^m$  by using the results on  $\theta$ -semistable representations proved in the foregoing section. We assume that  $x = (A_1, \dots, A_m)$  is brought into corner type  $C_s$  with  $s = (s_1, \dots, s_n) \in \mathcal{S}_n$ . We will associate a quiver-representation to  $x$ . As we are interested in checking whether we can transform  $x$  to a smaller corner-type, it is intuitively clear that the *border* region of  $C_s$  will be important.

- the *border*  $B_s$  is the subspace of  $C_s$  consisting of those  $m$ -tuples of  $n \times n$  matrices with zero entries except perhaps at entries  $(i, j)$  where  $s_i - s_j = 1$ .

EXAMPLE 6.25. For  $3 \times 3$  matrices we have the following corner-types  $C_s$  having border-regions  $B_s$  and associated Levi-subgroups  $L_s$






FIGURE 1. Corners and borders for  $4 \times 4$  matrices.

For  $4 \times 4$  matrices the relevant data are given in figure 1

From these examples, it is clear that the action of the Levi-subgroup  $L_s$  on the border  $B_s$  is a quiver-setting. In general, let  $s \in \mathcal{S}_n$  be determined by the tableau  $T_s$ , then the associated quiver-setting  $(Q_s, \alpha_s)$  is


- $Q_s$  is the quiver having as many connected components as there are rows in the tableau  $T_s$ . If the  $i$ -th row in  $T_s$  is

$$(a_{i0}, a_{i1}, \dots, a_{ik_i})$$

then the corresponding string of entries in  $s$  is of the form

$$\underbrace{\{p_i, \dots, p_i\}}_{a_{i0}}, \underbrace{\{p_i + 1, \dots, p_i + 1\}}_{a_{i1}}, \dots, \underbrace{\{p_i + k_i, \dots, p_i + k_i\}}_{a_{ik_i}}$$

and the  $i$ -th component of  $Q_s$  is defined to be the quiver  $Q_i$  on  $k_i + 1$  vertices having  $m$  arrows between the consecutive vertices, that is  $Q_i$  is



- the dimension vector  $\alpha_i$  for the  $i$ -th component quiver  $Q_i$  is equal to the  $i$ -th row of the tableau  $T_s$ , that is

$$\alpha_i = (a_{i0}, a_{i1}, \dots, a_{ik_i})$$

and the total dimension vector  $\alpha_s$  is the collection of these component dimension vectors.



- the character  $GL(\alpha_s) \xrightarrow{\chi_s} \mathbb{C}^*$  is determined by the integral  $n$ -tuple  $\theta_s = (t_1, \dots, t_n) \in \mathbb{Z}^n$  where if entry  $k$  corresponds to the  $j$ -th vertex of the  $i$ -th component of  $Q_s$  we have

$$t_k = n_{ij} \stackrel{\text{def}}{=} d \cdot (p_i + j)$$

where  $d$  is the least common multiple of the numerators of the  $p_i$ 's for all  $i$ . Equivalently, the  $n_{ij}$  are the integers appearing in the description of the one-parameter subgroup  $\lambda_s = (r_1, \dots, r_n)$  grouped together according to the ordering of vertices in the quiver  $Q_s$ . Recall that the character  $\chi_s$  is then defined to be

$$\chi_s(g_1, \dots, g_n) = \prod_{i=1}^n \det(g_i)^{t_i}$$

or in terms of  $GL(\alpha_s)$  it sends an element  $g_{ij} \in GL(\alpha_s)$  to  $\prod_{i,j} \det(g_{ij})^{n_{ij}}$ .

PROPOSITION 6.26. *The action of the Levi-subgroup  $L_s = \prod_{i,j} GL_{\alpha_{ij}}$  on the border  $B_s$  coincides with the base-change action of  $GL(\alpha_s)$  on the representation space  $\text{rep}_{\alpha_s} Q_s$ . The isomorphism*

$$B_s \longrightarrow \text{rep}_{\alpha_s} Q_s$$

is given by sending an  $m$ -tuple of border  $B_s$ -matrices  $(A_1, \dots, A_m)$  to the representation in  $\text{rep}_{\alpha_s} Q_s$  where the  $j$ -th arrow between the vertices  $v_a$  and  $v_{a+1}$  of the  $i$ -th component quiver  $Q_i$  is given by the relevant block in the matrix  $A_j$ .

EXAMPLE 6.27. We illustrate these definitions with a few of examples for  $4 \times 4$  matrices

tableau	$L_s$	$B_s$	$\theta_s$	$(Q_s, \alpha_s, \theta_s)$
$\begin{array}{ c c c } \hline 2 & 1 & 1 \\ \hline \end{array}$			$(5, 1, -3, -3)$	$\begin{array}{c} 5 & & 1 & & -3 \\ \textcircled{1} \leftarrow [m] \leftarrow \textcircled{1} \leftarrow [m] \leftarrow \textcircled{2} \end{array}$
$\begin{array}{ c c c } \hline 1 & 2 & 1 \\ \hline \end{array}$			$(1, 0, 0, -1)$	$\begin{array}{c} 1 & & 0 & & -1 \\ \textcircled{1} \leftarrow [m] \leftarrow \textcircled{2} \leftarrow [m] \leftarrow \textcircled{1} \end{array}$
$\begin{array}{ c c } \hline 1 & 2 \\ \hline 1 \\ \hline \end{array}$			$(1, 1, 0, -2)$	$\begin{array}{c} 1 & & -2 \\ \textcircled{2} \leftarrow [m] \leftarrow \textcircled{1} \\ 0 \\ \textcircled{1} \end{array}$

Using these conventions we can now state the main result of this section, giving a solution to the problem of optimal corners.

THEOREM 6.28. *Let  $x = (A_1, \dots, A_m) \in \text{Null}_n^m$  be of corner type  $C_s$ . Then,  $x$  is of optimal corner type  $C_s$  if and only if under the natural maps*

$$C_s \longrightarrow B_s \xrightarrow{\cong} \text{rep}_{\alpha_s} Q_s$$

(the first map forgets the non-border entries)  $x$  is mapped to a  $\theta_s$ -semistable representation in  $\text{rep}_{\alpha_s} Q_s$ .

**6.5. Hesselink stratification.**

Every orbit in  $Null_n^m$  has a representative  $x = (A_1, \dots, A_m)$  with all  $A_i$  strictly upper triangular matrices. That is, if  $N \subset M_n$  is the subspace of strictly upper triangular matrices, then the action map determines a surjection

$$GL_n \times N^m \xrightarrow{ac} Null_n^m$$

Recall that the standard *Borel subgroup*  $B$  is the subgroup of  $GL_n$  consisting of all upper triangular matrices and consider the action of  $B$  on  $GL_n \times M_n^m$  determined by

$$b.(g, x) = (gb^{-1}, b.x)$$

Then,  $B$ -orbits in  $GL_n \times N^m$  are mapped under the action map  $ac$  to the same point in the nullcone  $Null_n^m$ . Consider the morphisms

$$GL_n \times M_n^m \xrightarrow{\pi} GL_n/B \times M_n^m$$

which sends a point  $(g, x)$  to  $(gB, g.x)$ . The quotient  $GL_n/B$  is called a *flag variety* and is a projective manifold. Its points are easily seen to correspond to complete *flags*

$$\mathcal{F} : 0 \subset F_1 \subset F_2 \subset \dots \subset F_n = \mathbb{C}^n \quad \text{with} \quad \dim_{\mathbb{C}} F_i = i$$

of subspaces of  $\mathbb{C}^n$ . For example, if  $n = 2$  then  $GL_2/B \simeq \mathbb{P}^1$ . Consider the fiber  $\pi^{-1}$  of a point  $(\bar{g}, (B_1, \dots, B_m)) \in GL_n/B \times M_n^m$ . These are the points

$$(h, (A_1, \dots, A_m)) \quad \text{such that} \quad \begin{cases} g^{-1}h & = b \in B \\ bA_i b^{-1} & = g^{-1}B_i g \quad \text{for all } 1 \leq i \leq m. \end{cases}$$

Therefore, the fibers of  $\pi$  are precisely the  $B$ -orbits in  $GL_n \times M_n^m$ . That is, there exists a quotient variety for the  $B$ -action on  $GL_n \times M_n^m$  which is the trivial vectorbundle of rank  $mn^2$

$$\mathcal{T} = GL_n/B \times M_n^m \xrightarrow{p} GL_n/B$$

over the flag variety  $GL_n/B$ . We will denote with  $GL_n \times^B N^m$  the image of the subvariety  $GL_n \times N^m$  of  $GL_n \times M_n^m$  under this quotient map. That is, we have a commuting diagram

$$\begin{array}{ccc} GL_n \times N^m & \hookrightarrow & GL_n \times M_n^m \\ \downarrow & & \downarrow \\ GL_n \times^B N^m & \hookrightarrow & GL_n/B \times M_n^m \end{array}$$

Hence,  $\mathcal{V} = GL_n \times^B N^m$  is a sub-bundle of rank  $m \cdot \frac{n(n-1)}{2}$  of the trivial bundle  $\mathcal{T}$  over the flag variety. Note however that  $\mathcal{V}$  itself is not trivial as the action of  $GL_n$  does not map  $N^m$  to itself.

**THEOREM 6.29.** *Let  $U$  be the open subvariety of  $m$ -tuples of strictly upper triangular matrices  $N^m$  consisting of those tuples such that one of the component matrices has rank  $n - 1$ . The action map  $ac$  induces the commuting diagram of figure 2. The upper map is an isomorphism of  $GL_n$ -varieties for the action on fiber bundles to be left multiplication in the first component.*

$$\begin{array}{ccc}
 GL_n \times^B U & \xrightarrow{\simeq} & GL_n.U \\
 \downarrow & & \downarrow \\
 GL_n \times^B N^m & \xrightarrow{ac} & Null_n^m
 \end{array}$$

FIGURE 2. Resolution of the nullcone.

Therefore, there is a natural one-to-one correspondence between  $GL_n$ -orbits in  $GL_n.U$  and  $B$ -orbits in  $U$ . Further,  $ac$  is a desingularization of the nullcone and  $Null_n^m$  is irreducible of dimension

$$(m + 1) \frac{n(n - 1)}{2}.$$

PROOF. Let  $A \in N$  be a strictly upper triangular matrix of rank  $n - 1$  and  $g \in GL_n$  such that  $gAg^{-1} \in N$ , then  $g \in B$  as one verifies by first bringing  $A$  into Jordan-normal form  $J_n(0)$ . This implies that over a point  $x = (A_1, \dots, A_m) \in U$  the fiber of the action map

$$GL_n \times N^m \xrightarrow{ac} Null_n^m$$

has dimension  $\frac{n(n-1)}{2} = \dim B$ . Over all other points the fiber has at least dimension  $\frac{n(n-1)}{2}$ . But then, by the dimension formula we have

$$\dim Null_n^m = \dim GL_n + \dim N^m - \dim B = (m + 1) \frac{n(n - 1)}{2}$$

Over  $GL_n.U$  this map is an isomorphism of  $GL_n$ -varieties. Irreducibility of  $Null_n^m$  follows from surjectivity of  $ac$  as  $\mathbb{C}[Null_n^m] \hookrightarrow \mathbb{C}[GL_n] \otimes \mathbb{C}[N^m]$  and the latter is a domain. These facts imply that the induced action map

$$GL_n \times^B N^m \xrightarrow{ac} Null_n^m$$

is birational and as the former is a smooth variety (being a vectorbundle over the flag manifold), this is a desingularization.  $\square$

EXAMPLE 6.30. Let  $n = 2$  and  $m = 1$ . We have seen in chapter 3 that  $Null_2^1$  is a cone in 3-space with the singular top the orbit of the zero-matrix and the open complement the orbit of

$$\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

In this case the flag variety is  $\mathbb{P}^1$  and the fiber bundle  $GL_2 \times^B N$  has rank one. The action map is depicted in figure 3 and is a  $GL_2$ -isomorphism over the complement of the fiber of the top.

Theorem 6.29 gives us a complexity-reduction, both in the dimension of the acting group and in the dimension of the space acted upon, from

- $GL_n$ -orbits in the nullcone  $Null_n^m$ , to
- $B$ -orbits in  $N^m$ .

at least on the stratum  $GL_n.U$  described before. The aim of the *Hesselink stratification* of the nullcone is to extend this reduction also to the complement.

Let  $s \in \mathcal{S}_n$  and let  $C_s$  be the vectorspace of all  $m$ -tuples in  $M_n^m$  which are of corner-type  $C_s$ . We have seen that there is a Zariski open subset (but, possibly




FIGURE 3. Resolution of  $Null_2^1$ .

empty)  $U_s$  of  $C_s$  consisting of  $m$ -tuples of optimal corner type  $C_s$ . Observe that the action of conjugation of  $GL_n$  on  $M_n^m$  induces an action of the associated parabolic subgroup  $P_s$  on  $C_s$ .

DEFINITION 6.31. The *Hesslink stratum*  $S_s$  associated to  $s$  is the subvariety  $GL_n.U_s$  where  $U_s$  is the open subset of  $C_s$  consisting of the optimal  $C_s$ -type tuples.

The results of the foregoing section imply, similar to the proof of theorem 6.29, the complexity-reduction

THEOREM 6.32. *With notations as before we have a commuting diagram*

$$\begin{array}{ccc}
 GL_n \times^{P_s} U_s & \xrightarrow{\cong} & S_s \\
 \downarrow & & \downarrow \\
 GL_n \times^{P_s} C_s & \xrightarrow{ac} & \overline{S}_s
 \end{array}$$

where  $ac$  is the action map,  $\overline{S}_s$  is the Zariski closure of  $S_s$  in  $Null_n^m$  and the upper map is an isomorphism of  $GL_n$ -varieties.

Here,  $GL_n/P_s$  is the flag variety associated to the parabolic subgroup  $P_s$  and is a projective manifold. The variety  $GL_n \times^{P_s} C_s$  is a vectorbundle over the flag variety  $GL_n/P_s$  and is a subbundle of the trivial bundle  $GL_n \times^{P_s} M_n^m$ .

Therefore, the Hesslink stratum  $S_s$  is an irreducible smooth variety of dimension

$$\begin{aligned}
 \dim S_s &= \dim GL_n/P_s + rk GL_n \times^{P_s} C_s \\
 &= n^2 - \dim P_s + \dim_{\mathbb{C}} C_s
 \end{aligned}$$

and there is a natural one-to-one correspondence between the  $GL_n$ -orbits in  $S_s$  and the  $P_s$ -orbits in  $U_s$ .

Moreover, the vectorbundle  $GL_n \times^{P_s} C_s$  is a desingularization of  $\overline{S}_s$  hence 'feels' the gluing of  $S_s$  to the remaining strata. Finally, the ordering of corners has the geometric interpretation

$$\overline{S}_s \subset \bigcup_{\|s'\| \leq \|s\|} S_{s'}$$

We have seen that  $U_s = p^{-1} \text{rep}_{\alpha_s}^{ss}(Q_s, \theta_s)$  where  $C_s \xrightarrow{p} B_s$  is the canonical projection forgetting the non-border entries. As the action of the parabolic subgroup  $P_s$  restricts to the action of its Levi-part  $L_s$  on  $B_s = \text{rep}_{\alpha_s} Q$  we have a canonical projection

$$U_s/P_s \xrightarrow{p} M_{\alpha_s}^{ss}(Q_s, \theta_s)$$

to the moduli space of  $\theta_s$ -semistable representations in  $\text{rep}_{\alpha_s} Q_s$ . As none of the components of  $Q_s$  admits cycles, these moduli spaces are projective varieties. For small values of  $m$  and  $n$  these moduli spaces give good approximations to the study of the orbits in the nullcone.


**EXAMPLE 6.33.** (Nullcone of  $m$ -tuples of  $2 \times 2$  matrices)

In chapter 3 we have proved by brute force that the orbits in  $\text{Null}_2^m$  correspond to points on  $\mathbb{P}^1$  together with one extra orbit, the zero representation. For arbitrary  $m$ , the relevant strata-information for  $\text{Null}_2^m$  is contained in the following table

tableau	$s$	$B_s = C_s$	$P_s$	$(Q_s, \alpha_s, \theta_s)$
$\begin{array}{ c c } \hline 1 & 1 \\ \hline \end{array}$	$(\frac{1}{2}, -\frac{1}{2})$	$\begin{array}{ c c } \hline \blacksquare & \blacksquare \\ \hline \end{array}$	$\begin{array}{ c c } \hline \bullet & \bullet \\ \hline \end{array}$	$\begin{array}{c} 1 \leftarrow m \rightarrow -1 \\ \textcircled{1} \end{array}$
$\begin{array}{ c } \hline 2 \\ \hline \end{array}$	$(0, 0)$	$\begin{array}{ c c } \hline & \\ \hline \end{array}$	$\begin{array}{ c c } \hline \bullet & \bullet \\ \hline \end{array}$	$\begin{array}{c} 0 \\ \textcircled{2} \end{array}$

Because  $B_s = C_s$  we have that the orbit space  $U_s/P_s \simeq M_{\alpha_s}^{ss}(Q_s, \theta_s)$ . For the first stratum, every representation in  $\text{rep}_{\alpha_s} Q_s$  is  $\theta_s$ -semistable except the zero-representation (as it contains a subrepresentation of dimension  $\beta = (1, 0)$  and  $\theta_s(\beta) = -1 < 0$ ). The action of  $L_s = \mathbb{C}^* \times \mathbb{C}^*$  on  $\mathbb{C}^m - 0$  has as orbit space  $\mathbb{P}^{m-1}$ , classifying the orbits in the maximal stratum. The second stratum consists of one point, the zero representation.

**EXAMPLE 6.34.** A more interesting application, illustrating all of the general phenomena, is the description of orbits in the nullcone of two  $3 \times 3$  matrices. H. Kraft described them in [44, p. 202] by brute force. The orbit space decomposes as a disjoint union of tori and can be represented by the picture



Here, each node corresponds to a torus of dimension the right-hand side number in the bottom row. A point in this torus represents an orbit with dimension the left-hand side number. The top letter is included for classification purposes. That is, every orbit has a unique representant in the

following list of couples of  $3 \times 3$  matrices  $(A, B)$ . The top letter gives the torus, the first 2 rows give the first two rows of  $A$  and the last two rows give the first two rows of  $B$ ,  $x, y \in \mathbb{C}^*$

$a$ 0 1 0 0 0 1 0 x 0 0 0 y	$b$ 0 1 0 0 0 1 0 0 0 0 0 x	$c$ 0 1 0 0 0 1 0 x 0 0 0 0	$d$ 0 1 0 0 0 1 0 x y 0 0 x	$e$ 0 1 0 0 0 1 0 x 0 0 0 0	$f$ 0 0 0 0 0 1 0 1 0 0 0 x	$g$ 0 1 0 0 0 0 0 0 0 0 0 1	$h$ 0 1 0 0 0 1 0 0 x 0 0 0	$i$ 0 0 x 0 0 0 0 1 0 0 0 1
$j$ 0 0 0 0 0 1 0 1 0 0 0 0	$k$ 0 0 1 0 0 0 0 1 0 0 0 0	$l$ 0 0 0 0 0 1 0 0 1 0 0 0	$m$ 0 0 1 0 0 0 0 1 0 0 0 0	$n$ 0 0 0 0 0 0 0 1 0 0 0 1	$o$ 0 1 0 0 0 0 0 x 0 0 0 0	$p$ 0 1 0 0 0 0 0 0 0 0 0 0	$q$ 0 0 0 0 0 0 0 1 0 0 0 0	$r$ 0 0 0 0 0 0 0 0 0 0 0 0

We will now derive this result from the above description of the Hesselink stratification. To begin, the relevant data concerning  $\mathcal{S}_3$  is summarized in the following table

tableau	$s$	$B_s, C_s$	$P_s$	$(Q_s, \alpha_s, \theta_s)$
$\begin{bmatrix} 1 & 1 & 1 \end{bmatrix}$	$(1, 0, -1)$			
$\begin{bmatrix} 1 & 2 \end{bmatrix}$	$(\frac{1}{3}, \frac{1}{3}, -\frac{2}{3})$			
$\begin{bmatrix} 2 & 1 \end{bmatrix}$	$(\frac{2}{3}, -\frac{1}{3}, -\frac{1}{3})$			
$\begin{bmatrix} 1 & 1 \\ 1 \end{bmatrix}$	$(\frac{1}{2}, 0, -\frac{1}{2})$			
$\begin{bmatrix} 3 \end{bmatrix}$	$(0, 0, 0)$			

For the last four corner types,  $B_s = C_s$  whence the orbit space  $U_s/P_s$  is isomorphic to the moduli space  $M_{\alpha_s}^{ss}(Q_s, \theta_s)$ . Consider the quiver-setting



If the two arrows are not linearly independent, then the representation contains a proper subrepresentation of dimension-vector  $\beta = (1, 1)$  or  $(1, 0)$  and in both cases  $\theta_s(\beta) < 0$  whence the representation is not  $\theta_s$ -semistable. If the two arrows are linearly independent, we can use the  $GL_2$ -component to bring them in the form  $(\begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \end{bmatrix})$ , whence  $M_{\alpha_s}^{ss}(Q_s, \alpha_s)$  is reduced to one point, corresponding to the matrix-couple of type  $l$


$$\left( \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \right)$$

A similar argument, replacing linear independence by common zero-vector shows that also the quiver-setting corresponding to the tableau  $\begin{bmatrix} 2 & 1 \end{bmatrix}$  has one point as its moduli space, the matrix-tuple of type  $k$ . Incidentally, this shows that the corners corresponding to the tableaux  $\begin{bmatrix} 2 & 1 \\ 1 \end{bmatrix}$  or  $\begin{bmatrix} 1 & 2 \end{bmatrix}$  cannot be optimal when  $m = 1$  as then the row or column vector always has a kernel or

cokernel whence cannot be  $\theta_s$ -semistable. This of course corresponds to the fact that the only orbits in  $Null_3^1$  are those corresponding to the Jordan-matrixes

$$\begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix} \quad \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \quad \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$


which are respectively of corner type  $\overline{1111}$ ,  $\overline{11}$  and  $\overline{3}$ , whence the two other types do not occur. Next, consider the quiver setting



A representation in  $rep_{\alpha_s} Q_s$  is  $\theta_s$ -semistable if and only if the two maps are not both zero (otherwise, there is a subrepresentation of dimension  $\beta = (1, 0)$  with  $\theta_s(\beta) < 0$ ). The action of  $GL(\alpha_s) = \mathbb{C}^* \times \mathbb{C}^*$  on  $\mathbb{C}^2 - \underline{0}$  has a  $s$  orbit space  $\mathbb{P}^1$  and they are represented by matrix-couples

$$\left( \begin{bmatrix} 0 & 0 & a \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & b \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \right)$$

with  $[a : b] \in \mathbb{P}^1$  giving the types  $a, p$  and  $q$ . Clearly, the stratum  $\overline{3}$  consists just of the zero-matrix, which is type  $r$ . Remains to investigate the quiver-setting



Again, one easily verifies that a representation in  $rep_{\alpha_s} Q_s$  is  $\theta_s$ -semistable if and only if  $(a, b) \neq (0, 0) \neq (c, d)$  (for otherwise one would have subrepresentations of dimensions  $(1, 1, 0)$  or  $(1, 0, 0)$ ). The corresponding  $GL(\alpha_s)$ -orbits are classified by

$$M_{\alpha_s}^{ss}(Q_s, \theta_s) \simeq \mathbb{P}^1 \times \mathbb{P}^1$$

corresponding to the matrix-couples of types  $a, b, c, e, f, g, j, k$  and  $n$

$$\left( \begin{bmatrix} 0 & c & 0 \\ 0 & 0 & a \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & d & 0 \\ 0 & 0 & b \\ 0 & 0 & 0 \end{bmatrix} \right)$$

where  $[a : b]$  and  $[c : d]$  are points in  $\mathbb{P}^1$ . In this case, however,  $C_s \neq B_s$  and we need to investigate the fibers of the projection

$$U_s/P_s \xrightarrow{p} M_{\alpha_s}^{ss}(Q_s, \alpha_s)$$

Now,  $P_s$  is the Borel subgroup of upper triangular matrices and one verifies that the following two couples

$$\left( \begin{bmatrix} 0 & c & 0 \\ 0 & 0 & a \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & d & 0 \\ 0 & 0 & b \\ 0 & 0 & 0 \end{bmatrix} \right) \text{ and } \left( \begin{bmatrix} 0 & c & x \\ 0 & 0 & a \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & d & y \\ 0 & 0 & b \\ 0 & 0 & 0 \end{bmatrix} \right)$$

lie in the same  $B$ -orbit if and only if  $\det \begin{bmatrix} a & c \\ b & d \end{bmatrix} \neq 0$ , that is, if and only if  $[a : b] \neq [c : d]$  in  $\mathbb{P}^1$ . Hence, away from the diagonal  $p$  is an isomorphism. On the diagonal one can again verify by direct computation that the fibers of  $p$  are isomorphic to  $\mathbb{C}$ , giving rise to the cases  $d, h$  and  $i$  in the classification.

The connection between this approach and Kraft's result is depicted in figure 4. The picture on the left is Kraft's toric degeneration picture where we enclosed all orbits belonging to the same Hesselink strata, that is, having the same optimal corner type. The dashed region enclosed the orbits which do not come from the moduli spaces  $M_{\alpha_s}^{ss}(Q_s, \theta_s)$ , that is, those coming from the





FIGURE 4. Nullcone of couples of  $3 \times 3$  matrices.

projection  $U_s/P_s \rightarrow M_{\alpha_s}^{ss}(Q_s, \theta_s)$ . The picture on the right gives the ordering of the relevant corners.

EXAMPLE 6.35. We see that we get most orbits in the nullcone from the moduli spaces  $M_{\alpha_s}^{ss}(Q_s, \theta_s)$ . The reader is invited to work out the orbits in  $Null_4^2$ . We list here the moduli spaces of the relevant corners

corner	$M_{\alpha_s}^{ss}(Q_s, \theta_s)$	corner	$M_{\alpha_s}^{ss}(Q_s, \theta_s)$	corner	$M_{\alpha_s}^{ss}(Q_s, \theta_s)$
	$\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$		$\mathbb{P}^1$		$\mathbb{P}^1$
	$\mathbb{P}^3 \sqcup \mathbb{P}^1 \times \mathbb{P}^1 \sqcup \mathbb{P}^1 \times \mathbb{P}^1$		$\mathbb{P}^1 \sqcup S^2(\mathbb{P}^1)$		$\mathbb{P}^0$
	$\mathbb{P}^1$		$\mathbb{P}^1$		$\mathbb{P}^0$


Observe that two potential corners are missing in this list. This is because we have the following quiver setting for the corner



and there are no  $\theta_s$ -semistable representations as the two maps have a common kernel, whence a subrepresentation of dimension  $\beta = (1, 0)$  and  $\theta_s(\beta) < 0$ . A similar argument holds for the other



missing corner and quiver setting



For general  $n$ , a similar argument proves that the corners associated to the tableaux  $\begin{bmatrix} 1 & n \end{bmatrix}$  and  $\begin{bmatrix} n & 1 \end{bmatrix}$  are not optimal for tuples in  $Null_{n+1}^m$  unless  $m \geq n$ . It is also easy to see that with  $m \geq n$  all relevant corners appear in  $Null_{n+1}^m$ , that is all potential Hesselink strata are non-empty.

### 6.6. Cornering quiver representations.

In this section we generalize the results on matrices to representation of arbitrary quivers. In the final section we will see that we can reduce the study of the orbits in the fibers of

$$\text{rep}_n A \longrightarrow \text{iss}_n A$$

over a point  $\xi \in Sm_n A$  to a nullcone problem for the local quiver setting. Let  $Q$  be a (marked) quiver on  $k$  vertices  $\{v_1, \dots, v_k\}$  and fix a dimension vector  $\alpha = (a_1, \dots, a_k)$  and denote the total dimension  $\sum_{i=1}^k a_i$  by  $a$ . A representation  $V \in \text{rep}_\alpha Q$  is said to belong to the nullcone  $Null_\alpha Q$  if the trivial representation  $\underline{0} \in \overline{\mathcal{O}(V)}$ . Equivalently, all polynomial invariants are zero when evaluated in  $V$ , that is, the traces of all oriented cycles in  $Q$  are zero in  $V$ . By the Hilbert criterium 2.17 for  $GL(\alpha)$ ,  $V \in Null_\alpha Q$  if and only if there is a one-parameter subgroup

$$\mathbb{C}^* \xrightarrow{\lambda} GL(\alpha) = \begin{bmatrix} GL_{a_1} & & \\ & \ddots & \\ & & GL_{a_k} \end{bmatrix} \hookrightarrow GL_a$$

such that  $\varprojlim \lambda(t).V = \underline{0}$ . Up to conjugation in  $GL(\alpha)$ , or equivalently, replacing  $V$  by another point in the orbit  $\mathcal{O}(V)$ , we may assume that  $\lambda$  lies in the maximal torus  $T_a$  of  $GL(\alpha)$  (and of  $GL_a$ ) and can be represented by an integral  $a$ -tuple  $(r_1, \dots, r_a) \in \mathbb{Z}^a$  such that

$$\lambda(t) = \begin{bmatrix} t^{r_1} & & \\ & \ddots & \\ & & t^{r_a} \end{bmatrix}$$


We have to take the vertices into account, so we decompose the integer interval  $[1, 2, \dots, a]$  into *vertex intervals*  $I_{v_i}$  such that

$$[1, 2, \dots, a] = \sqcup_{i=1}^k I_{v_i} \quad \text{with} \quad I_{v_i} = \left[ \sum_{j=1}^{i-1} a_j + 1, \dots, \sum_{j=1}^i a_j \right]$$

If we recall that the weights of  $T_a$  are isomorphic to  $\mathbb{Z}^a$  having canonical generators  $\pi_p$  for  $1 \leq p \leq a$  we can decompose the representation space into weight spaces

$$\text{rep}_\alpha Q = \bigoplus_{\pi_{pq} = \pi_q - \pi_p} \text{rep}_\alpha Q(\pi_{pq})$$

where the eigenspace of  $\pi_{pq}$  is non-zero if and only if for  $p \in I_{v_i}$  and  $q \in I_{v_j}$ , there is an arrow



in the quiver  $Q$ . Call  $\pi_\alpha Q$  the set of weights  $\pi_{pq}$  which have non-zero eigenspace in  $\text{rep}_\alpha Q$ . Using this weight space decomposition we can write every representation as  $V = \sum_{p,q} V_{pq}$  where  $V_{pq}$  is a vector of the  $(p, q)$ -entries of the maps  $V(a)$  for all arrows  $a$  in  $Q$  from  $v_i$  to  $v_j$ . Using the fact that the action of  $T_a$  on  $\text{rep}_\alpha Q$  is induced by conjugation, we deduce as before that for  $\lambda$  determined by  $(r_1, \dots, r_a)$

$$\lim_{t \rightarrow 0} \lambda(t).V = \underline{0} \iff r_q - r_p \geq 1 \text{ whenever } V_{pq} \neq 0$$

Again, we define the corner type  $C$  of the representation  $V$  by defining the subset of real  $a$ -tuples

$$E_V = \{(x_1, \dots, x_a) \in \mathbb{R}^a \mid x_q - x_p \geq 1 \forall V_{pq} \neq 0\}$$

and determine a minimal element  $s_V$  in it, minimal with respect to the usual norm on  $\mathbb{R}^a$ . Similar to the case of matrices considered before, it follows that  $s_V$  is a uniquely determined point in  $\mathbb{Q}^a$ , having the characteristic property that its entries can be partitioned into strings

$$\underbrace{\{p_l, \dots, p_l\}}_{a_{l0}} \underbrace{\{p_l + 1, \dots, p_l + 1\}}_{a_{l1}} \dots \underbrace{\{p_l + k_l, \dots, p_l + k_l\}}_{a_{lk_l}} \quad \text{with all } a_{lm} \geq 1$$

which are balanced, that is  $\sum_{m=0}^{k_l} a_{lm}(p_l + m) = 0$ .

Note however that this time we are not allowed to bring  $s_V$  into dominant form, as we can only permute base-vectors of the vertex-spaces. That is, we can only use the action of the *vertex-symmetric groups*

$$S_{a_1} \times \dots \times S_{a_k} \hookrightarrow S_a$$

to bring  $s_V$  into *vertex dominant form*, that is if  $s_V = (s_1, \dots, s_a)$  then

$$s_q \leq s_p \quad \text{whenever } p, q \in I_{v_i} \text{ for some } i \text{ and } p < q$$

We compile a list  $\mathcal{S}_\alpha$  of such rational  $a$ -tuples by the following **algorithm**

- Start with the list  $\mathcal{S}_a$  of matrix corner types.
- For every  $s \in \mathcal{S}_a$  consider all permutations  $\sigma \in S_a / (S_{a_1} \times \dots \times S_{a_k})$  such that  $\sigma.s = (s_{\sigma(1)}, \dots, s_{\sigma(a)})$  is vertex dominant.
- Take  $\mathcal{H}_\alpha$  to be the list of the distinct  $a$ -tuples  $\sigma.s$  which are vertex dominant.
- Remove  $s \in \mathcal{H}_\alpha$  whenever there is an  $s' \in \mathcal{H}_\alpha$  such that

$$\pi_s Q = \{\pi_{pq} \in \pi_\alpha Q \mid s_q - s_p \geq 1\} \subset \pi_{s'} Q = \{\pi_{pq} \in \pi_\alpha Q \mid s'_q - s'_p \geq 1\}$$

and  $\|s\| > \|s'\|$ .

- The list  $\mathcal{S}_\alpha$  are the remaining entries  $s$  from  $\mathcal{H}_\alpha$ .

For  $s \in \mathcal{S}_\alpha$ , we define associated data similar to the case of matrices


- The *corner*  $C_s$  is the subspace of  $\text{rep}_\alpha Q$  such that all arrow matrices  $V_b$ , when viewed as  $a \times a$  matrices using the partitioning in vertex-entries, have only non-zero entries at spot  $(p, q)$  when  $s_q - s_p \geq 1$ .
- The *border*  $B_s$  is the subspace of  $\text{rep}_\alpha Q$  such that all arrow matrices  $V_b$ , when viewed as  $a \times a$  matrices using the partitioning in vertex-entries, have only non-zero entries at spot  $(p, q)$  when  $s_q - s_p = 1$ .

- The *parabolic subgroup*  $P_s(\alpha)$  is the intersection of  $P_s \subset GL_a$  with  $GL(\alpha)$  embedded along the diagonal.  $P_s(\alpha)$  is a parabolic subgroup of  $GL(\alpha)$ , that is, contains the product of the Borels  $B(\alpha) = B_{a_1} \times \dots \times B_{a_k}$ .
- The *Levi-subgroup*  $L_s(\alpha)$  is the intersection of  $L_s \subset GL_a$  with  $GL(\alpha)$  embedded along the diagonal.


We say that a representation  $V \in \text{rep}_\alpha Q$  is of *corner type*  $C_s$  whenever  $V \in C_s$ .

**THEOREM 6.36.** *By permuting the vertex-bases, every representation  $V \in \text{rep}_\alpha Q$  can be brought to a corner type  $C_s$  for a uniquely determined  $s$  which is a vertex-dominant reordering of  $s_V$ .*

**EXAMPLE 6.37.** Consider the following quiver setting



Then, the relevant corners have the following block decomposition



Again, we solve the problem of *optimal corner representations* by introducing a new quiver setting.


Fix a type  $s \in \mathcal{S}_\alpha Q$  and let  $J_1, \dots, J_u$  be the distinct strings partitioning the entries of  $s$ , say with

$$J_l = \underbrace{\{p_l, \dots, p_l\}}_{\sum_{i=1}^k b_{i,l0}} \underbrace{\{p_l + 1, \dots, p_l + 1\}}_{\sum_{i=1}^k b_{i,l1}} \dots \underbrace{\{p_l + k_l, \dots, p_l + k_l\}}_{\sum_{i=1}^k b_{i,lk_l}}$$

where  $b_{i,lm}$  is the number of entries  $p \in I_{v_i}$  such that  $s_p = p_l + m$ . To every string  $l$  we will associate a quiver  $Q_{s,l}$  and dimension vector  $\alpha_{s,l}$  as follows

- $Q_{s,l}$  has  $k \cdot (k_l + 1)$  vertices labeled  $(v_i, m)$  with  $1 \leq i \leq k$  and  $0 \leq m \leq k_l$ .
- In  $Q_{s,l}$  there are as many arrows from vertex  $(v_i, m)$  to vertex  $(v_j, m + 1)$  as there are arrows in  $Q$  from vertex  $v_i$  to vertex  $v_j$ . There are no arrows between  $(v_i, m)$  and  $(v_j, m')$  if  $m' - m \neq 1$ .
- The dimension-component of  $\alpha_{s,l}$  in vertex  $(v_i, m)$  is equal to  $b_{i,lm}$ .

EXAMPLE 6.38. For the above quiver, all component quivers  $Q_{s,l}$  are pieces of the quiver below



Clearly, we only need to consider that part of the quiver  $Q_{s,l}$  where the dimensions of the vertex spaces are non-zero.

The quiver-setting  $(Q_s, \alpha_s)$  associated to a type  $s \in \mathcal{S}_\alpha Q$  will be the disjoint union of the string quiver-settings  $(Q_{s,l}, \alpha_{s,l})$  for  $1 \leq l \leq u$ .

THEOREM 6.39. *With notations as before, for  $s \in \mathcal{S}_\alpha Q$  we have isomorphisms*

$$\begin{cases} B_s & \simeq \text{rep}_{\alpha_s} Q_s \\ L_s(\alpha) & \simeq GL(\alpha_s) \end{cases}$$

Moreover, the base-change action of  $GL(\alpha_s)$  on  $\text{rep}_{\alpha_s} Q_s$  coincides under the isomorphisms with the action of the Levi-subgroup  $L_s(\alpha)$  on the border  $B_s$ .

In order to determine the representations in  $\text{rep}_{\alpha_s} Q_s$  which have optimal corner type  $C_s$  we define the following character on the Levi-subgroup

$$L_s(\alpha) = \prod_{l=1}^u \times_{i=1}^k \times_{m=0}^{k_l} GL_{b_{i,lm}} \xrightarrow{\chi^{\theta_s}} \mathbb{C}^*$$

determined by sending a tuple  $(g_{i,lm})_{ilm} \rightarrow \prod_{ilm} \det g_{i,lm}^{m_{i,lm}}$  where the exponents are determined by

$$\theta_s = (m_{i,lm})_{ilm} \quad \text{where} \quad m_{i,lm} = d(p_l + m)$$

with  $d$  the least common multiple of the numerators of the rational numbers  $p_l$  for all  $1 \leq l \leq u$ .

THEOREM 6.40. *Consider a representation  $V \in \text{Null}_\alpha Q$  of corner type  $C_s$ . Then,  $V$  is of optimal corner type  $C_s$  if and only if under the natural maps*

$$C_s \xrightarrow{\pi} B_s \xrightarrow{\simeq} \text{rep}_{\alpha_s} Q_s$$

$V$  is mapped to a  $\theta_s$ -semistable representation in  $\text{rep}_{\alpha_s} Q_s$ . If  $U_s$  is the open subvariety of  $C_s$  consisting of all representations of optimal corner type  $C_s$ , then

$$U_s = \pi^{-1} \text{rep}_{\alpha_s}^{ss}(Q_s, \theta_s)$$

For the corresponding Hesselink stratum  $S_s = GL(\alpha).U_s$  we have the commuting diagram

$$\begin{array}{ccc} GL(\alpha) \times^{P_s(\alpha)} U_s & \xrightarrow{\simeq} & S_s \\ \downarrow & & \downarrow \\ GL(\alpha) \times^{P_s(\alpha)} C_s & \xrightarrow{ac} & \overline{S_s} \end{array}$$

where  $ac$  is the action map,  $\overline{S_s}$  is the Zariski closure of  $S_s$  in  $\text{Null}_\alpha Q$  and the upper map is an isomorphism as  $GL(\alpha)$ -varieties.

Here,  $GL(\alpha)/P_s(\alpha)$  is the flag variety associated to the parabolic subgroup  $P_s(\alpha)$  and is a projective manifold. The variety  $GL(\alpha) \times^{P_s(\alpha)} C_s$  is a vectorbundle over the flag variety  $GL(\alpha)/P_s(\alpha)$  and is a subbundle of the trivial bundle  $GL(\alpha) \times^{P_s(\alpha)} \text{rep}_\alpha Q$ .

Hence, the Hesselink stratum  $S_s$  is an irreducible smooth variety of dimension

$$\begin{aligned} \dim S_s &= \dim GL(\alpha)/P_s(\alpha) + \text{rk } GL(\alpha) \times^{P_s(\alpha)} C_s \\ &= \sum_{i=1}^k a_i^2 - \dim P_s(\alpha) + \dim_{\mathbb{C}} C_s \end{aligned}$$

and there is a natural one-to-one correspondence between the  $GL(\alpha)$ -orbits in  $S_s$  and the  $P_s(\alpha)$ -orbits in  $U_s$ .

Moreover, the vectorbundle  $GL(\alpha) \times^{P_s(\alpha)} C_s$  is a desingularization of  $\overline{S_s}$  hence 'feels' the gluing of  $S_s$  to the remaining strata. The ordering of corners has the geometric interpretation


$$\overline{S_s} \subset \bigcup_{\|s'\| \leq \|s\|} S_{s'}$$

Finally, because  $P_s(\alpha)$  acts on  $B_s$  by the restriction to its subgroup  $L_s(\alpha) = GL(\alpha_s)$  we have a projection from the orbit space










$$U_s/P_s \xrightarrow{P} M_{\alpha_s}^{ss}(Q_s, \theta_s)$$

to the moduli space of  $\theta_s$ -semistable quiver representations.







EXAMPLE 6.41. Above we have listed the relevant corner-types for the nullcone of the quiver-setting



In the table below we list the data of the three irreducible components of  $Null_\alpha Q/GL(\alpha)$  corresponding to the three maximal Hesselink strata :

$C_s, B_s$	$L_s$	$(Q_s, \alpha_s, \theta_s)$	$M_{\alpha_s}^{ss}(Q_s, \theta_s)$
 $(1, 0, -1)$			$\mathbb{P}^1$
 $(0, -1, 1)$			$\mathbb{P}^1$
 $(1, -1, 0)$			$\mathbb{P}^0$

There are 6 other Hesselink strata consisting of precisely one orbit. Finally, two possible corner-types do not appear as there are no  $\theta_s$ -semistable representations for the corresponding quiver setting

$C_s, B_s$	$L_s$	$(Q_s, \alpha_s, \theta_s)$	$M_{\alpha_s}^{ss}(Q_s, \theta_s)$
 $(\frac{1}{3}, \frac{1}{3}, -\frac{2}{3})$			$\emptyset$
 $(-\frac{1}{3}, -\frac{1}{3}, \frac{2}{3})$			$\emptyset$

### 6.7. Simultaneous conjugacy classes.

We have come a long way from our bare hands description of the simultaneous conjugacy classes of couples of  $2 \times 2$  matrices in section SimultaneousConjugacyClassesSection. In this section we will summarize what we have learned so far to approach this hopeless problem.

Having studied the nullcone of arbitrary quiver settings, we want to use these results to study the representations of Quillen-smooth algebras. Let  $A$  be an affine  $\mathbb{C}$ -algebra and  $M_\xi$  is a semi-simple  $n$ -dimensional module such that the representation variety  $\underline{rep}_n A @ \mathfrak{n}$  is smooth in  $M_\xi$ , that is  $\xi \in Sm_n A$ . Let  $M_\xi$  be of


representation type  $\tau = (e_1, d_1; \dots; e_k, d_k)$ , that is,

$$M_\xi = S_1^{\oplus e_1} \oplus \dots \oplus S_k^{\oplus e_k}$$

with distinct simple components  $S_i$  of dimension  $d_i$  and occurring in  $M_\xi$  with multiplicity  $e_i$ , then the  $GL(\alpha) = \text{Stab } M_\xi$ -structure on the normal space  $N_\xi$  to the orbit  $\mathcal{O}(M_\xi)$  is isomorphic to that of the representation space

$$\text{rep}_\alpha Q^\bullet$$

of a certain marked quiver on  $k$  vertices. The slice theorem asserts the existence of a slice  $S_\xi \xrightarrow{\phi} N_\xi$  and a commuting diagram



in a neighborhood of  $\xi \in \text{iss}_n A @ \mathbf{n}$  on the right and a neighborhood of the image  $\underline{0}$  of the trivial representation in  $N_\xi/GL(\alpha)$  on the left. In this diagram, the vertical maps are the quotient maps, all diagonal maps are étale and the upper ones are  $GL_n$ -equivariant. In particular, there is a  $GL_n$ -isomorphism between the fibers

$$\pi_2^{-1}(\underline{0}) \simeq \pi_1^{-1}(\xi)$$

Because  $\pi_2^{-1}(\underline{0}) \simeq GL_n \times^{GL(\alpha)} \pi^{-1}(\underline{0})$  with  $\pi$  is the quotient morphism for the marked quiver representations  $N_\xi = \text{rep}_\alpha Q^\bullet \xrightarrow{\pi} \text{iss}_\alpha Q^\bullet = N_x/GL(\alpha)$  we have a  $GL_n$ -isomorphism

$$\pi_1^{-1}(\xi) \simeq GL_n \times^{GL(\alpha)} \pi^{-1}(\underline{0})$$

That is, there is a natural one-to-one correspondence between

- $GL_n$ -orbits in the fiber  $\pi_1^{-1}(\xi)$ , that is, isomorphism classes of  $n$ -dimensional representations of  $A$  with Jordan-Hölder decomposition  $M_\xi$ , and
- $GL(\alpha)$ -orbits in  $\pi^{-1}(\underline{0})$ , that is, the nullcone of the marked quiver  $\text{Null}_\alpha Q^\bullet$ .

Summarizing we have the following

**THEOREM 6.42.** *Let  $A$  be an affine Quillen-smooth  $\mathbb{C}$ -algebra and  $M_\xi$  a semi-simple  $n$ -dimensional representation of  $A$ . Then, the isomorphism classes of  $n$ -dimensional representations of  $A$  with Jordan-Hölder sum isomorphic to  $M_\xi$  are given by the  $GL(\alpha)$ -orbits in the nullcone  $\text{Null}_\alpha Q^\bullet$  of the local marked quiver setting.*

The problem of classifying simultaneous conjugacy classes of  $m$ -tuples of  $n \times n$  matrices, is the same as  $n$ -dimensional representations of the Quillen-smooth algebra  $\mathbb{C}\langle x_1, \dots, x_m \rangle$ . To study semi-simple representations, one considers the quotient map

$$M_n^m = \text{rep}_n \mathbb{C}\langle x_1, \dots, x_m \rangle \xrightarrow{\pi} \text{iss}_n \mathbb{C}\langle x_1, \dots, x_m \rangle = \text{iss}_n^m$$

Fix a point  $\xi \in \text{iss}_n^m$  and assume that the corresponding semi-simple  $n$ -dimensional representation  $M_\xi$  is of representation type  $\tau = (e_1, d_1; \dots; e_k, d_k)$ .

We have shown that the coordinate ring  $\mathbb{C}[\text{iss}_n^m] = \mathbb{N}_n^m$  is the *necklace algebra*, that is, is generated by traces of monomials in the generic  $n \times n$  matrices  $X_1, \dots, X_m$  of length bounded by  $n^2 + 1$ . Further, if we collect all  $M_\xi$  with representation type  $\tau$  in the subset  $\text{iss}_n^m(\tau)$ , then

$$\text{iss}_n = \bigsqcup_{\tau} \text{iss}_n^m(\tau)$$

is a finite stratification of  $\text{iss}_n^m$  into locally closed smooth algebraic subvarieties.

We have an ordering on the representation types  $\tau' < \tau$  indicating that the stratum  $\text{iss}_n^m(\tau')$  is contained in the Zariski closure of  $\text{iss}_n^m(\tau)$ . This order relation is induced by the *direct ordering*

$$\tau' = (e'_1, d'_1; \dots; e'_{k'}, d'_{k'}) <^{dir} \tau = (e_1, d_1; \dots; e_k, d_k)$$


if there is a permutation  $\sigma$  of  $[1, 2, \dots, k']$  and there are numbers

$$1 = j_0 < j_1 < j_2 \dots < j_k = k'$$

such that for every  $1 \leq i \leq k$  we have the following relations

$$\begin{cases} e_i d_i &= \sum_{j=j_{i-1}+1}^{j_i} e'_{\sigma(j)} d'_{\sigma(j)} \\ e_i &\leq e'_{\sigma(j)} \text{ for all } j_{i-1} < j \leq j_i \end{cases}$$

EXAMPLE 6.43. The order relation on the representation types of dimension  $n = 4$  has the following Hasse diagram.



Because  $\text{iss}_n^m$  is irreducible, there is an open stratum corresponding to the simple representations, that is type  $(1, n)$ . The sub-generic strata are all of the form

$$\tau = (1, m_1; 1, m_2) \quad \text{with} \quad m_1 + m_2 = n.$$

The (in)equalities describing the locally closed subvarieties  $\text{iss}_n^m(\tau)$  can (in principle) be deduced from the theory of trace identities. Remains to study the local structure of the quotient variety  $\text{iss}_n^m$  near  $\xi$  and the description of the fibers  $\pi^{-1}(\xi)$ .



Both problems can be tackled by studying the local quiver setting  $(Q_\xi, \alpha_\xi)$  corresponding to  $\xi$  which describes the  $GL(\alpha_\xi) = Stab(M_\xi)$ -module structure of the normal space to the orbit of  $M_\xi$ . If  $\xi$  is of representation type  $\tau = (e_1, d_1; \dots; e_k, d_k)$  then the local quiver  $Q_\xi$  has  $k$ -vertices  $\{v_1, \dots, v_k\}$  corresponding to the  $k$  distinct simple components  $S_1, \dots, S_k$  of  $M_\xi$  and the number of arrows (resp. loops) from  $v_i$  to  $v_j$  (resp. in  $v_i$ ) are given by the dimensions

$$\dim_{\mathbb{C}} Ext^1(S_i, S_j) \quad \text{resp.} \quad \dim_{\mathbb{C}} Ext^1(S_i, S_i)$$

and these numbers can be computed from the dimensions of the simple components,

$$\begin{cases} \# \text{ } \textcircled{i} \xleftarrow{a} \textcircled{i} & = (m-1)d_i d_j \\ \# \text{ } \textcircled{i} & = (m-1)d_i^2 + 1 \end{cases}$$


Further, the local dimension vector  $\alpha_\xi$  is given by the multiplicities  $(e_1, \dots, e_k)$ . The étale local structure of  $iss_n^m$  in a neighborhood of  $\xi$  is the same as that of the quotient variety  $iss_{\alpha_\xi} Q_\xi$  in a neighborhood of  $\underline{0}$ . The local algebra of the latter is generated by traces along oriented cycles in the quiver  $Q_\xi$ . A direct application is

**PROPOSITION 6.44.** *For  $m \geq 2$ ,  $\xi$  is a smooth point of  $iss_n^m$  if and only if  $M_\xi$  is a simple representation, unless  $(m, n) = (2, 2)$  in which case  $iss_2^2 \simeq \mathbb{C}^5$  is a smooth variety.*

**PROOF.** If  $\xi$  is of representation type  $(1, n)$ , the local quiver setting  $(Q_\xi, \alpha_\xi)$  is



where  $d = (m-1)n^2 + 1$ , whence the local algebra is the formal power series ring in  $d$  variables and so  $iss_n^m$  is smooth in  $\xi$ . Because the singularities form a Zariski closed subvariety of  $iss_n^m$ , the result follows if we prove that all points  $\xi$  lying in sub-generic strata, say of type  $(1, m_1; 1, m_2)$  are singular. In this case the local quiver setting is equal to



where  $a = (m-1)m_1 m_2$  and  $l_i = (m-1)m_i^2 + 1$ . Let us denote the arrows from  $v_1$  to  $v_2$  by  $x_1, \dots, x_a$  and those from  $v_2$  to  $v_1$  by  $y_1, \dots, y_a$ . If  $(m, n) \neq (2, 2)$  then  $a \geq 2$ , but then we have traces along cycles

$$\{x_i y_j \mid 1 \leq i, j \leq a\}$$

that is, the polynomial ring of invariants is the polynomial algebra in  $l_1 + l_2$  variables (the traces of the loops) over the homogeneous coordinate ring of the Segre embedding

$$\mathbb{P}^{a-1} \times \mathbb{P}^{a-1} \hookrightarrow \mathbb{P}^{a^2-1}$$

which has a singularity at the top (for example we have equations of the form  $(x_1 y_2)(x_2 y_1) - (x_1 y_1)(x_2 y_2)$ ). Thus, the local algebra of  $iss_n^m$  cannot be a formal power series ring in  $\xi$  whence  $iss_n^m$  is singular in  $\xi$ . We have seen in section 1.2 that for the exceptional case  $iss_2^2 \simeq \mathbb{C}^5$ .  $\square$

To determine the fibers of the quotient map  $M_n^m \xrightarrow{\pi} iss_n^m$  we have to study the nullcone of this local quiver setting,  $Null_{\alpha_\xi} Q_\xi$ . Observe that the quiver  $Q_\xi$  has loops in every vertex and arrows connecting each ordered pair of vertices, whence we do not have to worry about potential corner-type removals. Denote  $\sum e_i = z \leq n$  and let  $\mathcal{C}_z$  be the set of all  $s = (s_1, \dots, s_z) \in \mathbb{Q}^z$  which are disjoint unions of strings of the form

$$\{p_i, p_i + 1, \dots, p_i + k_i\}$$

where  $l_i \in \mathbb{N}$ , all intermediate numbers  $p_i + j$  with  $j \leq k_i$  do occur as components in  $s$  with multiplicity  $a_{ij} \geq 1$  and  $p_i$  satisfies the balance-condition

$$\sum_{j=0}^{k_i} a_{ij}(p_i + j) = 0$$

for every string in  $s$ . For fixed  $s \in \mathcal{C}_z$  we can distribute the components  $s_i$  over the vertices of  $Q_\xi$  ( $e_j$  of them to vertex  $v_j$ ) in all possible ways modulo the action of the small Weyl group  $S_{e_1} \times \dots \times S_{e_k} \hookrightarrow S_z$ . That is, we can rearrange the  $s_i$ 's belonging to a fixed vertex such that they are in decreasing order. This gives us the list  $\mathcal{S}_{\alpha_\xi}$  or  $\mathcal{S}_\tau$  of all corner-types in  $Null_{\alpha_\xi} Q_\xi$ . For each  $s \in \mathcal{S}_{\alpha_\xi}$  we then construct the corner-quiver setting

$$(Q_{\xi \ s}, \alpha_{\xi \ s}, \theta_{\xi \ s})$$

and study the Hesselink strata  $S_s$  which actually do appear, which is equivalent to verifying whether there are  $\theta_{\xi \ s}$ -semistable representations in  $rep_{\alpha_{\xi \ s}} Q_{\xi \ s}$ . We have given a purely combinatorial way to settle this (in general quite hard) problem of optimal corner-types.

That is, we can determine which Hesselink strata  $S_s$  actually occur in  $\pi^{-1}(\xi) \simeq Null_{\alpha_{xi}} Q_\xi$ . The  $GL(\alpha_{\xi \ s})$ -orbits in the stratum  $S_s$  are in natural one-to-one correspondence with the orbits under the associated parabolic subgroup  $P_s$  acting on the semistable representations

$$U_s = \pi^{-1} rep_{\alpha_{\xi \ s}}^{ss}(Q_{\xi \ s}, \theta_{\xi \ s})$$

and there is a natural projection morphism from the corresponding orbit-space

$$U_s/P_s \xrightarrow{p_s} M_{\alpha_{\xi \ s}}^{ss}(Q_{\xi \ s}, \theta_{\xi \ s})$$

to the moduli space of  $\theta_{\xi \ s}$ -semistable representations. The remaining (hard) problem in the classification of  $m$ -tuples of  $n \times n$  matrices under simultaneous conjugation is the description of the fibers of this projection map  $p_s$ .

**EXAMPLE 6.45.** ( $m$ -tuples of  $2 \times 2$  matrices)

There are three different representation types  $\tau$  of 2-dimensional representations of  $\mathbb{C}\langle x_1, \dots, x_m \rangle$  with corresponding local quiver settings  $(Q_\tau, \alpha_\tau)$  given in figure 5 The defining (in)equalities of the strata  $iss_2^m(\tau)$  are given by  $k \times k$  minors (with  $k \leq 4$  of the symmetric  $m \times m$  matrix

$$\begin{bmatrix} tr(x_1^0 x_1^0) & \dots & tr(x_1^0 x_m^0) \\ \vdots & & \vdots \\ tr(x_m^0 x_1^0) & \dots & tr(x_m^0 x_m^0) \end{bmatrix}$$

where  $x_i^0 = x_i - \frac{1}{2}tr(x_i)$  is the generic trace zero matrix. These facts follow from the description of the trace algebras  $T_2^m$  as polynomial algebras over the generic Clifford algebras of rank  $\leq 4$  (determined by the above symmetric matrix) and the classical matrix decomposition of Clifford algebras over  $\mathbb{C}$ . For more details we refer to [48].




FIGURE 5. Local quiver settings for  $2 \times 2$  matrices.




FIGURE 6. Moduli spaces for type  $2_b$ .

To study the fibers  $M_2^m \rightarrow iss_2^m$  we need to investigate the different Hesselink strata in the nullcones of these local quiver settings. Type  $2_a$  has just one potential corner type corresponding to  $s = (0) \in \mathcal{S}_1$  and with corresponding corner-quiver setting

$$\begin{matrix} 0 \\ \textcircled{1} \end{matrix}$$

which obviously has  $\mathbb{P}^0$  (one point) as corresponding moduli (and orbit) space. This corresponds to the fact that for  $\xi \in iss_2^m(1, 2)$ ,  $M_\xi$  is simple and hence the fiber  $\pi^{-1}(\xi)$  consists of the closed orbit  $\mathcal{O}(M_\xi)$ .

For type  $2_b$  the list of figure 6 gives the potential corner-types  $C_s$  together with their associated corner-quiver settings and moduli spaces (note that as  $B_s = C_s$  in all cases, these moduli spaces describe the full fiber) That is, for  $\xi \in iss_2^m(1, 1; 1, 1)$ , the fiber  $\pi^{-1}(\xi)$  consists of the unique closed orbit  $\mathcal{O}(M_\xi)$  (corresponding to the  $\mathbb{P}^0$ ) and two families  $\mathbb{P}^{m-2}$  of non-closed orbits. Observe that in the special case  $m = 2$  we recover the two non-closed orbits found in section 1.2.




FIGURE 7. Local quiver settings for  $3 \times 3$  matrices.

Finally, for type  $2_c$ , the fibers are isomorphic to the nullcones of  $m$ -tuples of  $2 \times 2$  matrices. We have the following list of corner-types, corner-quiver settings and moduli spaces. Again, as  $B_s = C_s$  in all cases, these moduli spaces describe the full fiber.

$s$	$B_s, C_s$	$(Q_s, \alpha_s, \theta_s)$	$M_{\alpha_s}^{ss}(Q_s, \theta_s)$
$(\frac{1}{2}, -\frac{1}{2})$			$\mathbb{P}^{m-1}$
$(0, 0)$			$\mathbb{P}^0$


whence the fiber  $\pi^{-1}(\xi)$  consists of the closed orbit, together with a  $\mathbb{P}^{m-1}$ -family of non-closed orbits. Again, in the special case  $m = 2$ , we recover the  $\mathbb{P}^1$ -family found in section 1.2.

**EXAMPLE 6.46.** ( $m$ -tuples of  $3 \times 3$  matrices)

There are 5 different representation-types for 3-dimensional representations. Their associated local quiver settings are given in figure 7. For each of these types we can perform an analysis of the nullcones as before. We leave the details to the interested reader and mention only the end-result

- For type  $3_a$  the fiber is one closed orbit.
- For type  $3_b$  the fiber consists of the closed orbit together with two  $\mathbb{P}^{2m-3}$ -families of non-closed orbits.

- For type  $3_c$  the fiber consists of the closed orbit together with twelve  $\mathbb{P}^{m-2} \times \mathbb{P}^{m-2}$ -families and one  $\mathbb{P}^{m-2}$ -family of non-closed orbits.
- For type  $3_d$  the fiber consists of the closed orbit together with four  $\mathbb{P}^{m-1} \times \mathbb{P}^{m-2}$ -families, one  $\mathbb{P}^{m-2} \times \mathbb{P}^{m-2}$ -family, two  $\mathbb{P}^{m-2}$ -families, one  $\mathbb{P}^{m-1}$ -family and two  $M$ -families of non-closed orbits determined by moduli spaces of quivers, where  $M$  is the moduli space of the following quiver setting



together with some additional orbits coming from the projection maps  $p_s$ .

- For type  $3_e$  we have to study the nullcone of  $m$ -tuples of  $3 \times 3$  matrices, which can be done as in the case of couples but for  $m \geq 3$  the two extra strata do occur.

We see that in this case the only representation-types where the fiber is not fully determined by moduli spaces of quivers are  $3_d$  and  $3_e$ .

### 6.8. Representation fibers.

Let  $A$  be a Cayley-Hamilton algebra of degree  $n$  and consider the algebraic quotient map

$$\underline{rep}_n^{tr} A \xrightarrow{\pi} \underline{iss}_n^{tr} A$$

from the variety of  $n$ -dimensional trace preserving representations to the variety classifying isomorphism classes of trace preserving  $n$ -dimensional semi-simple representations. Assume  $\xi \in Sm_{tr} A \hookrightarrow \underline{iss}_n^{tr} A$ . That is, the representation variety  $\underline{rep}_n^{tr} A$  is smooth along the  $GL_n$ -orbit of  $M_\xi$  where  $M_\xi$  is the semi-simple representation determined by  $\xi \in \underline{iss}_n^{tr} A$ . We have seen that the local structure of  $A$  and  $\underline{rep}_n^{tr} A$  near  $\xi$  is fully determined by a local marked quiver setting  $(Q_\xi^\bullet, \alpha_\xi)$ . That is, we have a  $GL_n$ -isomorphism between the fiber of the quotient map, that is, the  $n$ -dimensional trace preserving representation degenerating to  $M_\xi$

$$\pi^{-1}(\xi) \simeq GL_n \times^{GL(\alpha_\xi)} Null_{\alpha_\xi} Q_\xi$$

and the nullcone of the marked quiver-setting. In this section we will apply the results on nullcones to the study of these representation fibers  $\pi^{-1}(\xi)$ .

Observe that all the facts on nullcones of quivers extend verbatim to marked quivers  $Q^\bullet$  using the underlying quiver  $Q$  with the proviso that we drop all loops in vertices with vertex-dimension 1 which get a marking in  $Q^\bullet$ . This is clear as nilpotent quiver representations obviously have zero trace along each oriented cycle, in particular in each loop.

The examples given before illustrate that a complete description of the nullcone is rather cumbersome. For this reason we restrict here to the determination of the number of irreducible components and their dimensions in the representation fibers. Modulo the  $GL_n$ -isomorphism above this study amounts to describing the irreducible components of  $Null_{\alpha_\xi} Q_\xi$  which are determined by the maximal corner-types  $C_s$ , that is such that the set of weights in  $C_s$  is maximal among subsets of  $\pi_{\alpha_{xi}} Q_\xi$  (and hence  $\|s\|$  is maximal among  $\mathcal{S}_{\alpha_\xi} Q_\xi$ ).

To illustrate our strategy, consider the case of curve orders. In section 5.4 we proved that if  $A$  is a Cayley-Hamilton order of degree  $n$  over an affine curve  $X = \underline{iss}_n^t A$  and if  $\xi \in Sm_{tr} A$ , then the local quiver setting  $(Q, \alpha)$  is determined by an oriented cycle  $Q$  on  $k$  vertices with  $k \leq n$  being the number of distinct simple components of  $M_\xi$ , the dimension vector  $\alpha = (1, \dots, 1)$  as in figure 8 and an unordered partition  $p = (d_1, \dots, d_k)$  having precisely  $k$  parts such that  $\sum_i d_i =$




FIGURE 8. Local quiver settings for curve orders.


$n$ , determining the dimensions of the simple components of  $M_\xi$ . Fixing a cyclic ordering of the  $k$ -vertices  $\{v_1, \dots, v_k\}$  we have that the set of weights of the maximal torus  $T_k = \mathbb{C}^* \times \dots \times \mathbb{C}^* = GL(\alpha)$  occurring in  $rep_\alpha Q$  is the set

$$\pi_\alpha Q = \{\pi_{k1}, \pi_{12}, \pi_{23}, \dots, \pi_{k-1k}\}$$

Denote  $K = \sum_{i=0}^{k-1} i = \frac{k(k-1)}{2}$  and consider the one string vector

$$s = (\dots, k - 2 - \frac{K}{k}, k - 1 - \frac{K}{k}, \underbrace{-\frac{K}{k}}_i, 1 - \frac{K}{k}, 2 - \frac{K}{k}, \dots)$$

then  $s$  is balanced and vertex-dominant,  $s \in \mathcal{S}_\alpha Q$  and  $\pi_s Q = \Pi$ . To check whether the corresponding Hesselink strata in  $Null_\alpha Q$  is nonempty we have to consider the associated quiver-setting  $(Q_s, \alpha_s, \theta_s)$  which is



It is well known and easy to verify that  $rep_{\alpha_s} Q_s$  has an open orbit with representative all arrows equal to 1. For this representation all proper subrepresentations have dimension vector  $\beta = (0, \dots, 0, 1, \dots, 1)$  and hence  $\theta_s(\beta) > 0$ . That is, the representation is  $\theta_s$ -stable and hence the corresponding Hesselink stratum  $S_s \neq \emptyset$ . Finally, because the dimension of  $rep_{\alpha_s} Q_s$  is  $k - 1$  we have that the dimension of this component in the representation fiber  $\pi^{-1}(x)$  is equal to

$$\dim GL_n - \dim GL(\alpha) + \dim rep_{\alpha_s} Q_s = n^2 - k + k - 1 = n^2 - 1$$

which completes the proof of the following

**THEOREM 6.47.** *Let  $A$  be a Cayley-Hamilton order of degree  $n$  over an affine curve  $X$  such that  $A$  is smooth in  $\xi \in X$ . Then, the representation fiber  $\pi^{-1}(\xi)$  has exactly  $k$  irreducible components of dimension  $n^2 - 1$ , each the closure of one orbit. In particular, if  $A$  is Cayley-smooth over  $X$ , then the quotient map*

$$\underline{rep}_n^t A \xrightarrow{\pi} \underline{iss}_n^t A = X$$

is flat, that is, all fibers have the same dimension  $n^2 - 1$ .




FIGURE 9. Local quiver settings for surface orders.

For Cayley-Hamilton orders over surfaces, the situation is slightly more complicated. From section 5.4 we recall that if  $A$  is a Cayley-Hamilton order of degree  $n$  over an affine surface  $S = \text{is}_n^t A$  and if  $A$  is smooth in  $\xi \in X$ , then the local structure of  $A$  is determined by a quiver setting  $(Q, \alpha)$  where  $\alpha = (1, \dots, 1)$  and  $Q$  is a two-circuit quiver on  $k+l+m \leq n$  vertices, corresponding to the distinct simple components of  $M_\xi$  as in figure 9 and an unordered partition  $p = (d_1, \dots, d_{k+l+m})$  of  $n$  with  $k+l+m$  non-zero parts determined by the dimensions of the simple components of  $M_\xi$ . With the indicated ordering of the vertices we have that

$$\pi_\alpha Q = \left\{ \pi_i \ i+1 \mid \begin{cases} 1 & \leq i \leq k-1 \\ k+1 & \leq i \leq k+l-1 \\ k+l+1 & \leq i \leq k+l+m-1 \end{cases} \right\} \\ \cup \{ \pi_k \ k+l+1, \pi_{k+l} \ k+l+1, \pi_{k+l+m} \ 1, \pi_{k+l+m} \ k+1 \}$$

As the weights of a corner cannot contain all weights of an oriented cycle in  $Q$  we have to consider the following two types of potential corner-weights  $\Pi$  of maximal cardinality

- (outer type) :  $\Pi = \pi_\alpha Q - \{ \pi_a, \pi_b \}$  where  $a$  is an edge in the interval  $[v_1, \dots, v_k]$  and  $b$  is an edge in the interval  $[v_{k+1}, \dots, v_{k+l}]$ .
- (inner type) :  $\Pi = \pi_\alpha Q - \{ \pi_c \}$  where  $c$  is an edge in the interval  $[v_{k+l+1}, v_{k+l+m}]$ .

There are  $2 + (k-1)(l-1)$  different subsets  $\Pi$  of outer type, each occurring as the set of weights of a corner  $C_s$ , that is  $\Pi = \pi_s Q$  for some  $s \in \mathcal{S}_\alpha Q$ . The two exceptional cases correspond to

$$\begin{cases} \Pi_1 & = \pi_\alpha Q - \{ \pi_{k+l+m} \ 1, \pi_{k+l} \ k+l+1 \} \\ \Pi_2 & = \pi_\alpha Q - \{ \pi_{k+l+m} \ k+1, \pi_k \ k+l+1 \} \end{cases}$$

which are of the form  $\pi_{s_i} Q$  with associated border quiver-setting  $(Q_{s_i}, \alpha_{s_i}, \theta_{s_i})$  where  $\alpha_{s_i} = (1, \dots, 1)$ ,  $Q_{s_i}$  are the full line subquivers of  $Q$  given in figure 10 with starting point  $v_1$  (resp.  $v_{k+1}$ ). The corresponding  $s_i \in \mathcal{S}_\alpha Q$  is a single string with




FIGURE 10. Border quiver settings.

minimal entry

$$-\frac{\sum_{i=0}^{k+l+m-1} i}{k+l+m} = -\frac{k+l+m-1}{2} \quad \text{at place} \quad \begin{cases} 1 \\ k+1 \end{cases}$$

and going with increments equal to one along the unique path. Again, one verifies that  $rep_{\alpha_s} Q_s$  has a unique open and  $\theta_s$ -stable orbit, whence these Hesselink strata do occur and the border  $B_s$  is the full corner  $C_s$ . The corresponding irreducible component in  $\pi^{-1}(\xi)$  has therefore dimension equal to  $n^2 - 1$  and is the closure of a unique orbit. The remaining  $(k-1)(l-1)$  subsets  $\Pi$  of outer type are of the form


$$\Pi_{ij} = \pi_{\alpha} Q - \{\pi_{i \ i+1}, \pi_{j \ j+1}\}$$

with  $1 \leq i \leq k-1$  and  $k+1 \leq j \leq k+l-1$ . We will see in a moment that they are again of type  $\pi_s Q$  for some  $s \in \mathcal{S}_{\alpha} Q$  with associated border quiver-setting






There are  $m-1$  different subsets  $\Pi_u$  of inner type, where for  $k+l+1 \leq u < k+l+m$  we define  $\Pi_u = \pi_\alpha Q - \{\pi_u \ u+1\}$ , that is dropping an edge in the middle



First assume that  $k = l$ . In this case we can walk through the quiver (with notations as before)



and hence the full subquiver of  $Q$  is part of a corner quiver-setting  $(Q_s, \alpha_s, \theta_s)$  where  $\alpha = (1, \dots, 1)$  and where  $s$  has as its minimal entry  $-\frac{x}{k+l+m}$  where


$$x = \sum_{i=1}^a i + 2 \sum_{i=1}^b (a+i) + \sum_{i=1}^c (a+b+i)$$

In this case we see that  $rep_{\alpha_s} Q_s$  has  $\theta_s$ -stable representations, in fact, there is a  $\mathbb{P}^1$ -family of such orbits. The corresponding Hesselink stratum is nonempty and the irreducible component of  $\pi^{-1}(\xi)$  determined by it has dimension

$$\dim GL_n - \dim GL(\alpha) + \dim rep_{\alpha_s} Q_s = n^2 - (k+l+m) + (k+l+m) = n^2$$

If  $l < k$ , then  $\Pi_u = \pi_s Q$  for some  $s \in \mathcal{S}_\alpha Q$  but this time the border quiver-setting  $(Q_s, \alpha_s, \theta_s)$  is determined by  $\alpha_s = (1, \dots, 1)$  and  $Q_s$  the full subquiver of  $Q$  by also

dropping the arrow corresponding to  $\pi_{k+l+1, k+l}$ , that is



If  $Q_s$  is this quiver (without the dashed arrow) then  $B_s = rep_{\alpha_s} Q_s$  and it contains an open orbit of a  $\theta_s$ -stable representation. Observe that  $s$  is determined as the one string vector with minimal entry  $-\frac{x}{k+l+m}$  where

$$x = \sum_{i=1}^a i + 2 \sum_{i=1}^b (a+i) + \sum_{i=1}^c (a+b+i) + \sum_{i=1}^d (a+b+c+i)$$

However, in this case  $B_s \neq C_s$  and we can identify  $C_s$  with  $rep_{\alpha_s} Q'_s$  where  $Q'_s$  is  $Q_s$  together with the dashed arrow. There is an  $\mathbb{A}^1$ -family of orbits in  $C_s$  mapping to the  $\theta_s$ -stable representation. In particular, the Hesselink stratum exists and the corresponding irreducible component in  $\pi^{-1}(\xi)$  has dimension equal to

$$\dim GL_n - \dim GL(\alpha) + \dim C_s = n^2 - (k+l+m) + (k+l+m) = n^2.$$

This concludes the proof of the description of the representation fibers of smooth orders over surfaces, summarized in the following result.

**THEOREM 6.48.** *Let  $A$  be a Cayley-Hamilton order of degree  $n$  over an affine surface  $X = iss_n^t A$  and assume that  $A$  is smooth in  $\xi \in X$  of local type  $(A_{klm}, \alpha)$ . Then, the representation fiber  $\pi^{-1}(\xi)$  has exactly  $2 + (k-1)(l-1) + (m-1)$  irreducible components of which  $2 + (k-1)(l-1)$  are of dimension  $n^2 - 1$  and are closure of one orbit and the remaining  $m-1$  have dimension  $n^2$  and are closures of a one-dimensional family of orbits. In particular, if  $A$  is Cayley-smooth, then the algebraic quotient map*

$$rep_n^t A \xrightarrow{\pi} iss_n^t A = X$$

*is flat if and only if all local quiver settings of  $A$  have quiver  $A_{klm}$  with  $m = 1$ .*

**References.**

The stratification of the nulcone used in this chapter is due to W. Hesselink [30]. The results in section 6.2 are due to A. Schofield [70]. The results in section 6.3 are due to A. King [38]. The proof of theorem 6.28 uses results of F. Kirwan [39]. Theorem 6.29 is due to H-P. Kraft [44], theorem 6.32 and theorem 6.40 are special cases of Hesselinks result [30]. All other results are due to L. Le Bruyn [51], [52] and [56].



## geo @<sub>n</sub>

Having studied the local structure of noncommutative manifolds, we will present some classes of examples in this part. Sum-varieties are families  $(X_n)_n$  of commutative varieties which are locally controlled by noncommutative algebras and are connected with morphisms

$$X_{n_1} \times \dots \times X_{n_i} \xrightarrow{+} X_{\sum n_i}$$

Such families often appear in moduli space problems of vectorbundles, with  $X_n$  the moduli space of semistable rank  $n$  bundles (with additional invariants) and the connecting morphisms induced by the direct sum of bundles. A related class of examples are the moduli spaces  $M_\alpha^{ss}(Q, \theta)$  of  $\theta$ -semistable representations of a quiver  $Q$ . In particular, if  $Q$  has no oriented cycles, each of these moduli spaces is a projective variety. As an example we associate to the problem of describing all finite dimensional representations of the torus knot groups (for example, the braid group  $B_3$ ) a sum family of projective varieties and study their local properties with the tools developed before.

The Quillen-smooth algebras controlling an object in **geo** @<sub>n</sub> are often derived (for example, by universal localization) from a fixed Quillen-smooth algebra  $A$ . We develop the formal theory of noncommutative differential forms, de Rham cohomology, symplectic structures etc. for a Quillen-smooth algebra  $A$ . In particular, for the path algebra  $\mathbb{C}Q^d$  of the double of a quiver, the noncommutative functions  $\mathfrak{dR}^0$  are equipped with a natural Poisson-bracket. This infinite dimensional Lie algebra is called the necklace Lie algebra because the necklace words in the quiver  $Q^d$  form a vectorspace bases and the Lie bracket is determined by opening and re-assembling the necklaces. These Lie algebras are associated to infinite dimensional automorphism groups acting naturally on certain moduli spaces.

In the final chapter we present a second possibility to construct noncommutative manifolds. Among the quotient varieties of the deformed preprojective algebras, we characterize those which are coadjoint orbits for the necklace Lie algebra. Deformed preprojective algebras are never Quillen-smooth but some of their approximations at specific dimension vectors are  $\alpha$ -smooth. The corresponding  $iss_\alpha \Pi_\theta$  are these coadjoint orbits. Presumably, families of these coadjoint orbits can be re-assembled to form nice infinite dimensional cellular spaces. We present the example of the adelic Grassmannian  $Gr^{ad}$  (important in the study of dynamical systems) which decomposes under the coadjoint action of a necklace Lie algebra in coadjoint orbits  $Gr^{ad}(n)$  each of which is the quotient variety of a deformed preprojective algebra. These quotient varieties also have a physical interpretation as the phase spaces of Calogero particles.



## Noncommutative Geometry

### 7.1. Formal structure.

Objects in noncommutative geometry  $\mathcal{A}_n$  are families of varieties  $(X_i)_i$  which are locally controlled by a set of noncommutative algebras  $\mathcal{A}$ . That is,  $X_i$  is locally the quotient variety of a representation variety  $rep_n A$  for some  $n$  and some  $\mathbb{C}$ -algebra  $A \in \mathcal{A}$ . In section 2.7 we have seen that the representation varieties form a somewhat mysterious subclass of the category of all (affine)  $GL_n$ -varieties. For this reason it is important to equip them with additional structures that may make them stand out among the  $GL_n$ -varieties. In this section we define the *formal structure* on representation varieties, extending in a natural way the formal structure introduced by M. Kapranov on smooth affine varieties. Let us give an illustrative example of this structure.

Consider the affine space  $\mathbb{A}^d$  with coordinate ring  $\mathbb{C}[x_1, \dots, x_d]$  and order the coordinate functions  $x_1 < x_2 < \dots < x_d$ . Let  $\mathfrak{f}_d$  be the free Lie algebra on  $\mathbb{C}x_1 \oplus \dots \oplus \mathbb{C}x_d$  which has an ordered basis  $B = \cup_{k \geq 1} B_k$  defined as follows.  $B_1$  is the ordered set  $\{x_1, \dots, x_d\}$  and  $B_2 = \{[x_i, x_j] \mid j < i\}$ , ordered such that  $B_1 < B_2$  and  $[x_i, x_j] < [x_k, x_l]$  iff  $j < l$  or  $j = l$  and  $i < k$ . Having constructed the ordered sets  $B_l$  for  $l < k$  we define

$$B_k = \{[t, w] \mid t = [u, v] \in B_l, w \in B_{k-l} \text{ such that } v \leq w < t \text{ for } l < k\}.$$

For  $l < k$  we let  $B_l < B_k$  and  $B_k$  is ordered by  $[t, w] < [t', w']$  iff  $w < w'$  or  $w = w'$  and  $t < t'$ .

It is well known that  $B$  is an ordered  $\mathbb{C}$ -basis of the Lie algebra  $\mathfrak{f}_d$  and that its enveloping algebra

$$U(\mathfrak{f}_d) = \mathbb{C}\langle x_1, \dots, x_d \rangle$$

is the free associative algebra on the  $x_i$ . We number the elements of  $\cup_{k \geq 2} B_k$  according to the order  $\{b_1, b_2, \dots\}$  and for  $b_i \in B_k$  we define  $ord(b_i) = k - 1$  (the number of brackets needed to define  $b_i$ ). Let  $\Lambda$  be the set of all functions with finite support  $\lambda : \cup_{k \geq 2} B_k \longrightarrow \mathbb{N}$  and define  $ord(\lambda) = \sum \lambda(b_i) ord(b_i)$ . Rephrasing the *Poincaré-Birkhoff-Witt* result for  $U(\mathfrak{f}_d)$  we have that any noncommutative polynomial  $p \in \mathbb{C}\langle x_1, \dots, x_d \rangle$  can be written uniquely as a finite sum

$$p = \sum_{\lambda \in \Lambda} [f_\lambda] M_\lambda$$

where  $[f_\lambda] \in \mathbb{C}[x_1, \dots, x_d] = S(B_1)$  and  $M_\lambda = \prod_i b_i^{\lambda(b_i)}$ . In particular, for every  $\lambda, \mu, \nu \in \Lambda$ , there is a unique bilinear differential operator with polynomial coefficients

$$C_{\lambda\mu}^\nu : \mathbb{C}[x_1, \dots, x_d] \otimes_{\mathbb{C}} \mathbb{C}[x_1, \dots, x_d] \longrightarrow \mathbb{C}[x_1, \dots, x_d]$$

defined by expressing the product  $\llbracket f \rrbracket M_\lambda \cdot \llbracket g \rrbracket M_\mu$  in  $\mathbb{C}\langle x_1, \dots, x_d \rangle$  uniquely as  $\sum_{\nu \in \Lambda} \llbracket C_{\lambda\mu}^\nu(f, g) \rrbracket M_\nu$ .

By associativity of  $\mathbb{C}\langle x_1, \dots, x_d \rangle$  the  $C_{\lambda\mu}^\nu$  satisfy the *associativity constraint*, that is, we have equality of the trilinear differential operators

$$\sum_{\mu_1} C_{\mu_1\lambda_3}^\nu \circ (C_{\lambda_1\lambda_2}^{\mu_1} \otimes id) = \sum_{\mu_2} C_{\lambda_1\mu_2}^\nu \circ (id \otimes C_{\lambda_2\lambda_3}^{\mu_2})$$

for all  $\lambda_1, \lambda_2, \lambda_3, \nu \in \Lambda$ . That is, one can define the algebra  $\mathbb{C}\langle x_1, \dots, x_d \rangle_{[[\text{ob}]]}$  to be the  $\mathbb{C}$ -vectorspace of possibly *infinite formal sums*  $\sum_{\lambda \in \Lambda} \llbracket f_\lambda \rrbracket M_\lambda$  with multiplication defined by the operators  $C_{\lambda\mu}^\nu$ .

Let  $A_d(\mathbb{C})$  be the  $d$ -th *Weyl algebra*, that is, the ring of differential operators with polynomial coefficients on  $\mathbb{A}^d$ . Let  $\mathcal{O}_{\mathbb{A}^d}$  be the structure sheaf on  $\mathbb{A}^d$  then it is well-known that the ring of sections  $\mathcal{O}_{\mathbb{A}^d}(U)$  on any Zariski open subset  $U \hookrightarrow \mathbb{A}^d$  is a left  $A_d(\mathbb{C})$ -module. Define a sheaf  $\mathcal{O}_{\mathbb{A}^d}^f$  of noncommutative algebras on  $\mathbb{A}^d$  by taking as its sections over  $U$  the algebra

$$\mathcal{O}_{\mathbb{A}^d}^f(U) = \mathbb{C}\langle x_1, \dots, x_d \rangle_{[[\text{ob}]]} \otimes_{\mathbb{C}[x_1, \dots, x_d]} \mathcal{O}_{\mathbb{A}^d}(U)$$

that is the  $\mathbb{C}$ -vectorspace of possibly infinite formal sums  $\sum_{\lambda \in \Lambda} \llbracket f_\lambda \rrbracket M_\lambda$  with  $f_\lambda \in \mathcal{O}_{\mathbb{A}^d}(U)$  and the multiplication is given as before by the action of the bilinear differential operators  $C_{\lambda\mu}^\nu$  on the left  $A_d(\mathbb{C})$ -module  $\mathcal{O}_{\mathbb{A}^d}(U)$ , that is, for all  $f, g \in \mathcal{O}_{\mathbb{A}^d}(U)$  we have

$$\llbracket f \rrbracket M_\lambda \cdot \llbracket g \rrbracket M_\mu = \sum_{\nu} \llbracket C_{\lambda\mu}^\nu(f, g) \rrbracket M_\nu$$

This sheaf of noncommutative algebras  $\mathcal{O}_{\mathbb{A}^d}^f$  is called *the formal structure* on  $\mathbb{A}^d$ .

We will now define such structures on arbitrary affine smooth varieties. Let  $R$  be an associative  $\mathbb{C}$ -algebra,  $R^{Lie}$  its Lie structure and  $R_m^{Lie}$  the subspace spanned by the expressions  $[r_1, [r_2, \dots, [r_{m-1}, r_m] \dots]]$  containing  $m-1$  instances of Lie brackets. The *commutator filtration* of  $R$  is the (increasing) filtration by ideals  $(F^k R)_{k \in \mathbb{Z}}$  with  $F^k R = R$  for  $k \in \mathbb{N}$  and

$$F^{-k} R = \sum_m \sum_{i_1 + \dots + i_m = k} RR_{i_1}^{Lie} R \dots RR_{i_m}^{Lie} R$$


Observe that all  $\mathbb{C}$ -algebra morphisms preserve the commutator filtration. The *associated graded algebra*  $gr_F R$  is a (negatively) graded commutative *Poisson algebra* with part of degree zero, the *abelianization*  $R_{ab} = \frac{R}{[R, R]}$ . If  $R = \mathbb{C}\langle x_1, \dots, x_d \rangle$ , then the commutator filtration has components

$$F^{-k} \mathbb{C}\langle x_1, \dots, x_d \rangle = \left\{ \sum_{\lambda} \llbracket f_\lambda \rrbracket M_\lambda, \forall \lambda : ord(\lambda) \geq k \right\}$$

**DEFINITION 7.1.** Denote with  $\text{nil}_k$  the category of associative  $\mathbb{C}$ -algebras  $R$  such that  $F^{-k} R = 0$  (in particular,  $\text{nil}_1 = \text{commalg}$  the category of commutative  $\mathbb{C}$ -algebras). An algebra  $A \in \text{Ob}(\text{nil}_k)$  is said to be *k-smooth* if and only if for all  $T \in \text{Ob}(\text{nil}_k)$ , all nilpotent twosided ideals  $I \triangleleft T$  and all  $\mathbb{C}$ -algebra morphisms



$A \xrightarrow{\phi} \frac{T}{I}$  there exist a lifted  $\mathbb{C}$ -algebra morphism



making the diagram commutative. Alternatively, an algebra is  $k$ -smooth if and only if it is  $\text{Nil}_k$ -smooth.

For example, the quotient  $\frac{\mathbb{C}\langle x_1, \dots, x_d \rangle}{F^{-k} \mathbb{C}\langle x_1, \dots, x_d \rangle}$  is  $k$ -smooth using the lifting property of free algebras and the fact that algebra morphisms preserve the commutator filtration. Generalizing this, if  $A$  is Quillen-smooth then the quotient

$$A^{(k)} = \frac{A}{F^{-k} A}$$

is  $k$ -smooth.

Kapranov proves [34, Thm 1.6.1] that an affine commutative *Grothendieck-smooth* algebra  $C$  has a *unique* (upto  $\mathbb{C}$ -algebra isomorphism identical on  $C$ )  $k$ -smooth *thickening*  $C^{(k)}$  with  $C_{ab}^{(k)} \simeq C$ . The inverse limit (connecting morphisms are given by the unicity result)

$$C^f = \varprojlim C^{(k)}$$

is then called the *formal completion* of  $C$ . Clearly, one has  $C_{ab}^f = C$ . For example,

$$\mathbb{C}[x_1, \dots, x_d]^f = \varprojlim \frac{\mathbb{C}\langle x_1, \dots, x_d \rangle}{F^{-k} \mathbb{C}\langle x_1, \dots, x_d \rangle} \simeq \mathbb{C}\langle x_1, \dots, x_d \rangle_{[[\text{cb}]]}.$$

If  $X$  is an affine smooth (commutative) variety, one can use the formal completion  $\mathbb{C}[X]^f$  to define a sheaf of noncommutative algebras  $\mathcal{O}_X^f$  defining the *formal structure* on  $X$ .

The fact that  $C$  is Grothendieck-smooth is essential to construct and prove uniqueness of the formal completion. At present, no sufficiently functorial extension of formal completion is known for arbitrary commutative  $\mathbb{C}$ -algebras. It is not true that any (non affine) smooth variety can be equipped with a formal structure. In fact, the obstruction gives important new invariants of a smooth variety related to *Atiyah classes*. We refer to [34, §4] for more details.

We recall briefly the algebraic construction of *microlocalization*. Let  $R$  be a filtered algebra with a *separated filtration*  $\{F_n\}_{n \in \mathbb{Z}}$  and let  $S$  be a multiplicatively closed subset of  $R$  containing 1 but not 0. For any  $r \in F_n - F_{n-1}$  we denote its *principal character*  $\sigma(r)$  to be the image of  $r$  in the associated graded algebra  $gr(R)$ . We assume that the set  $\sigma(S)$  is a multiplicatively closed subset of  $gr(R)$ . We define the *Rees ring*  $\tilde{R}$  to be the graded algebra

$$\tilde{R} = \bigoplus_{n \in \mathbb{Z}} F_n t^n \hookrightarrow R[t, t^{-1}]$$

where  $t$  is an extra central variable. If  $\sigma(s) \in gr(R)_n$  then we define the element  $\tilde{s} = st^n \in \tilde{R}_n$ . The set  $\tilde{S} = \{\tilde{s}, s \in S\}$  is a multiplicatively closed subset of homogeneous elements in  $\tilde{R}$ .

Assume that  $\sigma(S)$  is an Ore set in  $gr(R) = \frac{\tilde{R}}{(t)}$ , then for every  $n \in \mathbb{N}_0$  the image  $\pi_n(\tilde{S})$  is an Ore set in  $\frac{\tilde{R}}{(t^n)}$  where  $\tilde{R} \twoheadrightarrow \frac{\tilde{R}}{(t^n)}$  is the quotient morphism. Hence, we have an inverse system of graded localizations and can form the inverse limit in the graded sense

$$Q_S^\mu(\tilde{R}) = \varprojlim^g \pi_n(\tilde{S})^{-1} \frac{\tilde{R}}{(t^n)}$$

The element  $t$  acts torsionfree on this limit and hence we can form the filtered algebra

$$Q_S^\mu(R) = \frac{Q_S^\mu(\tilde{R})}{(t-1)Q_S^\mu(\tilde{R})}$$

which is the *micro-localization* of  $R$  at the multiplicatively closed subset  $S$ . We recall that the associated graded algebra of the microlocalization can be identified with the graded localization

$$gr(Q_S^\mu(R)) = \sigma(S)^{-1} gr(R).$$

Let  $R$  be a  $\mathbb{C}$ -algebra with  $R_{ab} = \frac{R}{[R,R]} = C$ . We assume that the commutator filtration  $(F^k)_{k \in \mathbb{Z}}$  is a separated filtration on  $R$ . Observe that this is not always the case (for example consider  $U(\mathfrak{g})$  for  $\mathfrak{g}$  a semi-simple Lie algebra) but often one can repeat the argument below replacing  $R$  with  $\frac{R}{\square F^n}$ .

Observe that  $gr(R)$  is a negatively graded commutative algebra with part of degree zero  $C$ . Take a multiplicatively closed subset  $S_c$  of  $C$ , then  $S = S_c + [R, R]$  is a multiplicatively closed subset of  $R$  with the property that  $\sigma(S) = S_c$  and clearly  $S_c$  is an Ore set in  $gr(R)$ . Therefore,  $\tilde{S}$  is a multiplicatively closed set of the Rees ring  $\tilde{R}$  consisting of homogeneous elements of degree zero. Observing that  $(t^n)_0 = F^{-n}t^n$  for all  $n \in \mathbb{N}_0$  we see that

$$Q_S^\mu(R) = \varprojlim^g \pi_n(S)^{-1} \frac{R}{F^{-n}}$$

where  $R \xrightarrow{\pi_n} \frac{R}{F^{-n}}$  is the quotient morphism and  $Q_S^\mu$  is filtered again by the commutator filtration and has as associated graded algebra

$$gr(Q_S^\mu(R)) = S_c^{-1} gr(R).$$

One can define a *microstructure sheaf*  $\mathcal{O}_R^\mu$  on the affine scheme  $X$  of  $C$  by taking as its sections over the affine Zariski open set  $X(f)$

$$\Gamma(X(f), \mathcal{O}_R^\mu) = Q_{S_f}^\mu(R)$$

where  $S = \{1, f, f^2, \dots\} + [R, R]$ . For  $C$  a Grothendieck-smooth affine commutative algebra this sheaf of noncommutative algebras is the *formal structure* on  $X$  introduced by M. Kapranov.

An important remark to make is that one really needs microlocalization to construct a sheaf of noncommutative algebras on  $X$ . If by some fluke we would have that all the  $S_f$  are already Ore sets in  $R$ , we might optimistically assume that taking as sections over  $X(f)$  the Ore localization  $S_f^{-1}R$  we would define a sheaf  $\mathcal{O}_R$  over  $X$ . This is in general *not* the case as the Ore set  $S_g$  need no longer be Ore in a localization  $S_f^{-1}R$ !

Still one can remedy this by defining a *noncommutative Zariski topology* on  $X$  using *words* in the Ore sets  $S_f$ , see [79, §1.3]. Whereas we do not need this to

define formal structures it seems to me inevitable that at a later stage in the development of noncommutative geometry we will need to resort to such *noncommutative Grothendieck topologies* on usual commutative schemes.

Having define a formal structure on affine smooth varieties, we will now extend it to arbitrary representation varieties. The starting point is that for every associative algebra  $A$  the functor

$$\mathbf{alg} \xrightarrow{Hom_{\mathbf{alg}}(A, M_n(-))} \mathbf{sets}$$

is *representable* in  $\mathbf{alg}$ . That is, there exists an associative  $\mathbb{C}$ -algebra  $\sqrt[n]{A}$  such that there is a natural equivalence between the functors

$$Hom_{\mathbf{alg}}(A, M_n(-)) \underset{n.e.}{\sim} Hom_{\mathbf{alg}}(\sqrt[n]{A}, -).$$

In other words, for every associative  $\mathbb{C}$ -algebra  $B$ , there is a functorial one-to-one correspondence between the sets

$$\begin{cases} \text{algebra maps } A \longrightarrow M_n(B) \\ \text{algebra maps } \sqrt[n]{A} \longrightarrow B \end{cases}$$

We call  $\sqrt[n]{A}$  the *n-th root algebra* of  $A$ .

EXAMPLE 7.2. If  $A = \mathbb{C}\langle x_1, \dots, x_d \rangle$ , then it is easy to see that  $\sqrt[n]{A}$  is the free algebra  $\mathbb{C}\langle x_{11,1}, \dots, x_{nn,d} \rangle$  on  $dn^2$  variables. For, given an algebra map  $A \xrightarrow{\phi} M_n(B)$  we obtain an algebra map  $\sqrt[n]{A} \longrightarrow B$  by sending the free variable  $x_{ij,k}$  to the  $(i, j)$ -entry of the matrix  $\phi(x_k) \in M_n(B)$ . Conversely, to an algebra map  $\sqrt[n]{A} \xrightarrow{\psi} B$  we assign the algebra map  $A \longrightarrow M_n(B)$  by sending  $x_k$  to the matrix  $(\psi(x_{ij,k}))_{i,j} \in M_n(B)$ . Clearly, these operations are each others inverses.

To define  $\sqrt[n]{A}$  in general, consider the free algebra product  $A * M_n(\mathbb{C})$  and consider the subalgebra

$$\sqrt[n]{A} = A * M_n(\mathbb{C})^{M_n(\mathbb{C})} = \{p \in A * M_n(\mathbb{C}) \mid p.(1 * m) = (1 * m).p \ \forall m \in M_n(\mathbb{C})\}$$

Before we can prove the universal property of  $\sqrt[n]{A}$  we need to recall a property that  $M_n(\mathbb{C})$  shares with any Azumaya algebra : if  $M_n(\mathbb{C}) \xrightarrow{\phi} R$  is an algebra morphism and if  $R^{M_n(\mathbb{C})} = \{r \in R \mid r.\phi(m) = \phi(m).r \ \forall m \in M_n(\mathbb{C})\}$ , then we have  $R \simeq M_n(\mathbb{C}) \otimes_{\mathbb{C}} R^{M_n(\mathbb{C})}$ .

In particular, if we apply this to  $R = A * M_n(\mathbb{C})$  and the canonical map  $M_n(\mathbb{C}) \xrightarrow{\phi} A * M_n(\mathbb{C})$  where  $\phi(m) = 1 * m$  we obtain that  $M_n(\sqrt[n]{A}) = M_n(\mathbb{C}) \otimes_{\mathbb{C}} \sqrt[n]{A} = A * M_n(\mathbb{C})$ .

Hence, if  $\sqrt[n]{A} \xrightarrow{f} B$  is an algebra map we can consider the composition

$$A \xrightarrow{id_A * 1} A * M_n(\mathbb{C}) \simeq M_n(\sqrt[n]{A}) \xrightarrow{M_n(f)} M_n(B)$$

to obtain an algebra map  $A \longrightarrow M_n(B)$ . Conversely, consider an algebra map  $A \xrightarrow{g} M_n(B)$  and the canonical map  $M_n(\mathbb{C}) \xrightarrow{i} M_n(B)$  which centralizes  $B$  in  $M_n(B)$ . Then, by the universal property of free algebra products we have an algebra map  $A * M_n(\mathbb{C}) \xrightarrow{g * i} M_n(B)$  and restricting to  $\sqrt[n]{A}$  we see that this maps

factors

$$\begin{array}{ccc}
 A * M_n(\mathbb{C}) & \xrightarrow{g^*i} & M_n(B) \\
 \uparrow & & \uparrow \\
 \downarrow & & \downarrow \\
 \sqrt[n]{A} & \cdots\cdots\cdots & B
 \end{array}$$

and one verifies that these two operations are each others inverses.

It follows from the functoriality of the  $\sqrt[n]{\cdot}$  construction that  $\mathbb{C}\langle x_1, \dots, x_d \rangle \longrightarrow A$  implies that  $\sqrt[n]{\mathbb{C}\langle x_1, \dots, x_d \rangle} \longrightarrow \sqrt[n]{A}$ . Therefore, if  $A$  is affine and generated by  $\leq d$  elements, then  $\sqrt[n]{A}$  is also affine and generated by  $\leq dn^2$  elements.

These properties allow us define a *formal completion* of  $\mathbb{C}[rep_n A]$  in a functorial way for any associative algebra  $A$ . Equip  $\sqrt[n]{A}$  with the commutator filtration

$$\dots \hookrightarrow F_{-2} \sqrt[n]{A} \hookrightarrow F_{-1} \sqrt[n]{A} \hookrightarrow \sqrt[n]{A} = \sqrt[n]{A} = \dots$$

Because algebra morphisms are commutator filtration preserving, it follows from the universal property of  $\sqrt[n]{A}$  that  $\frac{\sqrt[n]{A}}{F_{-k} \sqrt[n]{A}}$  is the object in  $\mathfrak{nil}_k$  representing the functor

$$\mathfrak{nil}_k \xrightarrow{Hom_{\mathfrak{alg}}(A, M_n(-))} \mathbf{sets}.$$

In particular, because the categories  $\mathfrak{commalg}$  and  $\mathfrak{nil}_1$  are naturally equivalent, we deduce that

$$\sqrt[n]{A}_{ab} = \frac{\sqrt[n]{A}}{[\sqrt[n]{A}, \sqrt[n]{A}]} = \frac{\sqrt[n]{A}}{F_{-1} \sqrt[n]{A}} \simeq \mathbb{C}[rep_n A]$$

because both algebras represent the same functor. We now define

$$\sqrt[n]{A}_{[[\mathfrak{ab}]]} = \lim_{\leftarrow} \frac{\sqrt[n]{A}}{F_{-k} \sqrt[n]{A}}.$$

Assume that  $A$  is Quillen-smooth, then so is  $\sqrt[n]{A}$  because we have seen before that

$$M_n(\sqrt[n]{A}) \simeq A * M_n(\mathbb{C})$$

and the class of Quillen-smooth algebras is easily seen to be closed under free products and matrix algebras.

As a consequence, we have for every  $k \in \mathbb{N}$  that the quotient  $\frac{\sqrt[n]{A}}{F_{-k} \sqrt[n]{A}}$  is  $k$ -smooth. Moreover, we have that

$$\left( \frac{\sqrt[n]{A}}{F_{-k} \sqrt[n]{A}} \right)_{ab} \simeq \frac{\sqrt[n]{A}}{[\sqrt[n]{A}, \sqrt[n]{A}]} \simeq \mathbb{C}[rep_n A].$$

Because  $\mathbb{C}[rep_n A]$  is an affine commutative Grothendieck-smooth algebra, we deduce from the uniqueness of  $k$ -smooth thickenings that

$$\mathbb{C}[rep_n A]^{(k)} \simeq \frac{\sqrt[n]{A}}{F_{-k} \sqrt[n]{A}}$$

and consequently that the formal completion of  $\mathbb{C}[rep_n A]$  can be identified with

$$\mathbb{C}[rep_n A]^f \simeq \sqrt[n]{A}_{[[\mathfrak{ab}]]}.$$

Therefore, if we define for an arbitrary  $\mathbb{C}$ -algebra  $A$  the *formal completion* of  $\mathbb{C}[rep_n A]$  to be  $\sqrt[n]{A}_{[[\text{ob}]}}$  we have a canonical extension of the formal structure on affine Grothendieck-smooth commutative algebras to the class of coordinate rings of representation spaces on which it is functorial in the algebras.

There is a natural action of  $GL_n$  by algebra automorphisms on  $\sqrt[n]{A}$ . Let  $u_A$  denote the universal morphism  $A \xrightarrow{u_A} M_n(\sqrt[n]{A})$  corresponding to the identity map on  $\sqrt[n]{A}$ . For  $g \in GL_n$  we can consider the composed algebra map

$$\begin{array}{ccc} A & \xrightarrow{u_A} & M_n(\sqrt[n]{A}) \\ & \searrow \psi_g & \downarrow g \cdot g^{-1} \\ & & M_n(\sqrt[n]{A}) \end{array}$$

Then  $g$  acts on  $\sqrt[n]{A}$  via the automorphism  $\sqrt[n]{A} \xrightarrow{\phi_g} \sqrt[n]{A}$  corresponding to the composition  $\psi_g$ . It is easy to verify that this defines indeed a  $GL_n$ -action on  $\sqrt[n]{A}$ .

The *formal structure sheaf*  $\mathcal{O}_{rep_n A}^f$  defined over  $rep_n A$  constructed from  $\sqrt[n]{A}$  will be denoted by  $\mathcal{O}_{\sqrt[n]{A}}^f$ . We see that it actually has a  $GL_n$ -structure which is compatible with the  $GL_n$ -action on  $rep_n A$ .

### 7.2. Semi invariants.

A major source of examples of noncommutative varieties are *moduli spaces* of  $\theta$ -semistable representations of quivers. Because the moduli space  $M_\alpha^{ss}(Q, \theta)$  is by definition the projective scheme of the graded *algebra of semi-invariants* of weight  $\chi_\theta^n$  for some  $n$

$$M_\alpha^{ss}(Q, \theta) = \text{proj } \bigoplus_{n=0}^\infty \mathbb{C}[rep_\alpha Q]^{GL(\alpha), \chi_\theta^n}$$

we need some control on these semi-invariants of quivers.

In this section we will give a generating set of semi-invariants. The strategy of proof should be clear by now. First, we will describe a large set of semi-invariants. Then we use classical invariant theory to describe all multilinear semi-invariants of  $GL(\alpha)$ , or equivalently, all multilinear invariants of  $SL(\alpha) = SL_{a_1} \times \dots \times SL_{a_k}$  and describe them in terms of these determinantal semi-invariants. Finally, we show by polarization and restitution that these semi-invariants do indeed generate all semi-invariants.

Let  $Q$  be a quiver on  $k$  vertices  $\{v_1, \dots, v_k\}$ . We introduce the *additive  $\mathbb{C}$ -category  $\text{add } Q$*  generated by the quiver. For every vertex  $v_i$  we introduce an indecomposable object which we denote by  $\textcircled{i}$ . An arbitrary object in  $\text{add } Q$  is then a sum of these

$$\textcircled{1}^{\oplus e_1} \oplus \dots \oplus \textcircled{k}^{\oplus e_k}$$

That is we can identify  $\text{add } Q$  with  $\mathbb{N}^k$ . Morphisms in the category  $\text{add } Q$  are defined by the rules

$$\begin{cases} \text{Hom}_{\text{add } Q}(\textcircled{i}, \textcircled{j}) = \textcircled{j} \xleftarrow{\text{dotted arrow}} \textcircled{i} \\ \text{Hom}_{\text{add } Q}(\textcircled{i}, \textcircled{i}) = \textcircled{i} \xleftarrow{\text{dotted arrow}} \textcircled{i} \end{cases}$$

where the right hand sides are the  $\mathbb{C}$ -vectorspaces spanned by all oriented paths from  $v_i$  to  $v_j$  in the quiver  $Q$ , including the idempotent (trivial) path  $e_i$  when  $i = j$ .

Clearly, for any  $k$ -tuples of positive integers  $\alpha = (u_1, \dots, u_k)$  and  $\beta = (v_1, \dots, v_k)$

$$Hom_{\mathbf{add} Q} (\bigoplus_{i=1}^k \textcircled{1}^{\oplus u_i} \oplus \dots \oplus \bigoplus_{i=1}^k \textcircled{k}^{\oplus u_i}, \bigoplus_{i=1}^k \textcircled{1}^{\oplus v_i} \oplus \dots \oplus \bigoplus_{i=1}^k \textcircled{k}^{\oplus v_i})$$

is defined by matrices and composition arises via matrix multiplication

$$\begin{bmatrix} M_{v_1 \times u_1}(\textcircled{1} \overset{\curvearrowright}{\leftarrow} \textcircled{1}) & \dots & M_{v_1 \times u_k}(\textcircled{1} \overset{\curvearrowright}{\leftarrow} \textcircled{k}) \\ \vdots & \ddots & \vdots \\ M_{v_k \times u_1}(\textcircled{k} \overset{\curvearrowright}{\leftarrow} \textcircled{1}) & \dots & M_{v_k \times u_k}(\textcircled{k} \overset{\curvearrowright}{\leftarrow} \textcircled{k}) \end{bmatrix}$$

Fix a dimension vector  $\alpha = (a_1, \dots, a_k)$  and a morphism  $\phi$  in  $\mathbf{add} Q$

$$\bigoplus_{i=1}^k \textcircled{1}^{\oplus u_i} \oplus \dots \oplus \bigoplus_{i=1}^k \textcircled{k}^{\oplus u_i} \xrightarrow{\phi} \bigoplus_{i=1}^k \textcircled{1}^{\oplus v_i} \oplus \dots \oplus \bigoplus_{i=1}^k \textcircled{k}^{\oplus v_i}$$

For any representation  $V \in rep_{\alpha} Q$  we can replace each occurrence of an arrow  $\textcircled{j} \xleftarrow{a} \textcircled{i}$  of  $Q$  in  $\phi$  by the  $a_j \times a_i$ -matrix  $V_a$ . This way we obtain a rectangular matrix

$$V(\phi) \in M_{\sum_{i=1}^k a_i v_i \times \sum_{i=1}^k a_i u_i}(\mathbb{C})$$

If we are in a situation where  $\sum a_i v_i = \sum a_i u_i$ , then we can define a *semi-invariant polynomial function* on  $rep_{\alpha} Q$  by

$$P_{\alpha, \phi}(V) = \det V(\phi)$$

We call such semi-invariants *determinantal semi-invariants*. One verifies that  $P_{\phi, \alpha}$  is a semi-invariant of weight  $\chi_{\theta}$  where  $\theta = (u_1 - v_1, \dots, u_k - v_k)$ . We will show that such determinantal semi-invariant together with traces along oriented cycles in the quiver  $Q$  generate all semi-invariants.

Because semi-invariants for the  $GL(\alpha)$ -action on  $rep_{\alpha} Q$  are the same as invariants for the restricted action of  $SL(\alpha) = SL_{a_1} \times \dots \times SL_{a_k}$ , we will describe the multilinear  $SL(\alpha)$ -invariants from classical invariant theory. Because

$$\begin{aligned} rep_{\alpha} Q &= \bigoplus_{\textcircled{j} \xleftarrow{a} \textcircled{i}} M_{a_j \times a_i}(\mathbb{C}) \\ &= \bigoplus_{\textcircled{j} \xleftarrow{a} \textcircled{i}} \mathbb{C}^{a_i} \otimes \mathbb{C}^{*a_j} \end{aligned}$$

we have to consider multilinear  $SL(\alpha)$ -invariants of

$$\bigotimes_{\textcircled{j} \xleftarrow{a} \textcircled{i}} \mathbb{C}^{a_i} \otimes \mathbb{C}^{*a_j} = \bigotimes_{\textcircled{i}} \left[ \bigotimes_{\textcircled{j} \xleftarrow{a} \textcircled{i}} \mathbb{C}^{a_i} \otimes \bigotimes_{\textcircled{i} \xleftarrow{a} \textcircled{j}} \mathbb{C}^{*a_i} \right]$$

Hence, any multilinear  $SL(\alpha)$ -invariant can be written as  $f = \prod_{i=1}^k f_i$  where  $f_i$  is a  $SL_{a_i}$ -invariant of

$$\begin{array}{ccc} \begin{array}{c} \otimes \\ \leftarrow \textcircled{i} \end{array} & \mathbb{C}^{a_i} \otimes & \begin{array}{c} \otimes \\ \textcircled{i} \leftarrow \end{array} & \mathbb{C}^{*a_i} \end{array}$$

To increase our cultural luggage let us recall the classical description of multilinear  $SL_n$ -invariants on  $M_n^{\oplus i} \oplus V_n^{\oplus j} \oplus V_n^{*\oplus z}$ , that is, the  $SL_n$ -invariant linear maps

$$\underbrace{M_n \otimes \dots \otimes M_n}_i \otimes \underbrace{V_n \otimes \dots \otimes V_n}_j \otimes \underbrace{V_n^* \otimes \dots \otimes V_n^*}_z \xrightarrow{f} \mathbb{C}$$

By the identification  $M_n = V_n \otimes V_n^*$  we have to determine the  $SL_n$ -invariant linear maps

$$V_n^{\otimes i+j} \otimes V_n^{*\otimes i+z} \xrightarrow{f} \mathbb{C}$$

The description of such invariants is given by classical invariant theory, see [81, II.5,Thm. 2.5.A].

**THEOREM 7.3.** *The multilinear  $SL_n$ -invariants  $f$  are linear combinations of invariants of one of the following two types*

- (1) For  $(i_1, \dots, i_n, h_1, \dots, h_n, \dots, t_1, \dots, t_n, s_1, \dots, s_r)$  a permutation of the  $i + j$  vector indices and  $(u_1, \dots, u_r)$  a permutation of the  $i + z$  covector indices, consider the  $SL_n$ -invariant

$$[v_{i_1}, \dots, v_{i_n}] [v_{h_1}, \dots, v_{h_n}] \dots [v_{t_1}, \dots, v_{t_n}] \phi_{u_1}(v_{s_1}) \dots \phi_{u_r}(v_{s_r})$$

where the brackets are the determinantal invariants

$$[v_{a_1}, \dots, v_{a_n}] = \det \begin{bmatrix} v_{a_1} & v_{a_2} & \dots & v_{a_n} \end{bmatrix}$$

- (2) For  $(i_1, \dots, i_n, h_1, \dots, h_n, \dots, t_1, \dots, t_n, s_1, \dots, s_r)$  a permutation of the  $i + z$  covector indices and  $(u_1, \dots, u_r)$  a permutation of the  $i + j$  vector indices, consider the  $SL_n$ -invariant

$$[\phi_{i_1}, \dots, \phi_{i_n}]^* [\phi_{h_1}, \dots, \phi_{h_n}]^* \dots [\phi_{t_1}, \dots, \phi_{t_n}]^* \phi_{u_1}(v_{s_1}) \dots \phi_{u_r}(v_{s_r})$$

where the cobrackets are the determinantal invariants

$$[\phi_{a_1}, \dots, \phi_{a_n}]^* = \det \begin{bmatrix} \phi_{a_1} \\ \vdots \\ \phi_{a_n} \end{bmatrix}$$

Observe that we do not have at the same time brackets and cobrackets, due to the relation

$$[v_1, \dots, v_n] [\phi_1, \dots, \phi_n] = \det \begin{bmatrix} \phi_1(v_1) & \dots & \phi_1(v_n) \\ \vdots & & \vdots \\ \phi_n(v_1) & \dots & \phi_n(v_n) \end{bmatrix}$$

We can give a matrix-interpretation of these basic invariants. Let us consider the case of a bracket of vectors (the case of cobrackets is similar)

$$[v_{i_1}, \dots, v_{i_n}]$$

If all the indices  $\{i_1, \dots, i_n\}$  are original vector-indices (and so do not come from the matrix-terms) we save this term and go to the next factor. Otherwise, if say  $i_1$  is one of the matrix indices,  $A_{i_1} = \phi_{i_1} \otimes v_{i_1}$ , then the covector  $\phi_{i_1}$  must be paired

up in a scalar product  $\phi_{i_1}(v_{u_1})$  with a vector  $v_{u_1}$ . Again, two cases can occur. If  $u_1$  is a vector index, we have that

$$\phi_{i_1}(v_{u_1})[v_{i_1}, \dots, v_{i_n}] = [A_{i_1} v_{u_1}, v_{i_2}, \dots, v_{i_n}] = [v'_{i_1}, v_{i_2}, \dots, v_{i_n}]$$

Otherwise, we can keep on matching the matrix indices and get an expression

$$\phi_{i_1}(v_{u_1}) \phi_{u_1}(v_{u_2}) \phi_{u_2}(v_{u_3}) \dots$$

until we finally hit again a vector index, say  $u_l$ , but then we have the expression

$$\phi_{i_1}(v_{u_1}) \phi_{u_1}(v_{z_1}) \dots \phi_{u_{l-1}}(v_{u_l}) [v_{i_1}, \dots, v_{i_n}] = [M v_{u_l}, v_{i_2}, \dots, v_{i_n}]$$


where  $M = A_{i_1} A_{u_1} \dots A_{u_{l-1}}$ . One repeats the same argument for all vectors in the brackets. As for the remaining scalar product terms, we have a similar procedure of matching up the matrix indices and one verifies that in doing so one obtains factors of the type

$$\phi(Mv) \text{ and } tr(M)$$

where  $M$  is a monomial in the matrices. As we mentioned, the case of covector-brackets is similar except that in matching the matrix indices with a covector  $\phi$ , one obtains a monomial in the transposed matrices.

Having found these interpretations of the basic  $SL_n$ -invariant linear terms, we can proceed by polarization and restitution processes to prove

**THEOREM 7.4.** *The  $SL_n$ -invariants of  $W = rep_\alpha Q'$  where  $Q'$  is the quiver*



are generated by the following four types of functions, where we write a typical element in  $W$  as

$$\underbrace{(A_1, \dots, A_k)}_k, \underbrace{(v_1, \dots, v_m)}_m, \underbrace{(\phi_1, \dots, \phi_p)}_p$$

with the  $A_i$  the matrices corresponding to the loops, the  $v_j$  making up the rows of the  $n \times m$  matrix and the  $\phi_j$  the columns of the  $p \times n$  matrix.

- $tr(M)$  where  $M$  is a monomial in the matrices  $A_i$ ,
- scalar products  $\phi_j(Mv_i)$  where  $M$  is a monomial in the matrices  $A_i$ ,
- brackets  $[M_1 v_{i_1}, M_2 v_{i_2}, \dots, M_n v_{i_n}]$  where the  $M_j$  are monomials in the matrices  $A_i$ ,
- cobrackets  $[M_1 \phi_{i_1}^T, \dots, M_n \phi_{i_n}^T]$  where the  $M_j$  are monomials in the matrices  $A_i$ ,

Returning to the special case under consideration, that is, of  $SL_m$ -invariants on  $\otimes_B \mathbb{C}^m \otimes \otimes_C \mathbb{C}^{*m}$ , it follows from this that the linear  $SL_m$ -invariants are determined by the following three sets

- *traces*, that is, for each pair  $(b, c)$  we have  $\mathbb{C}^m \otimes \mathbb{C}^{*m} = M_m(\mathbb{C}) \xrightarrow{Tr} \mathbb{C}$ .
- *brackets*, that is, for each  $m$ -tuple  $(b_1, \dots, b_m)$  we have an invariant  $\otimes_{b_j} \mathbb{C}^m \longrightarrow \mathbb{C}$  defined by

$$v_{b_1} \otimes \dots \otimes v_{b_m} \mapsto det [v_{b_1} \quad \dots \quad v_{b_m}]$$



- *cobrackets*, that is, for each  $m$ -tuple  $(c_1, \dots, c_m)$  we have an invariant  $\otimes_{c_i} \mathbb{C}^{*m} \longrightarrow \mathbb{C}$  defined by

$$\phi_{c_1} \otimes \dots \otimes \phi_{c_m} \mapsto \det \begin{bmatrix} \phi_{c_1} \\ \vdots \\ \phi_{c_m} \end{bmatrix}$$

Multilinear  $SL_m$ -invariants of  $\otimes_B \mathbb{C}^m \otimes \otimes_C \mathbb{C}^{*m}$  are then spanned by invariants constructed from the following data. Take three disjoint index-sets  $I, J$  and  $K$  and consider surjective maps

$$\begin{cases} B & \xrightarrow{\mu} I \sqcup K \\ C & \xrightarrow{\nu} J \sqcup K \end{cases}$$

subject to the following conditions

$$\begin{cases} \# \mu^{-1}(k) = 1 = \# \nu^{-1}(k) & \text{for all } k \in K. \\ \# \mu^{-1}(i) = m = \# \nu^{-1}(j) & \text{for all } i \in I \text{ and } j \in J. \end{cases}$$

To this data  $\gamma = (\mu, \nu, I, J, K)$  we can associate a multilinear  $SL_m$ -invariant  $f_\gamma(\otimes_B v_b \otimes \otimes_C \phi_c)$  defined by

$$\prod_{k \in K} \phi_{\nu^{-1}(k)}(v_{\mu^{-1}(k)}) \prod_{i \in I} \det [v_{b_1} \dots v_{b_m}] \prod_{j \in J} \det \begin{bmatrix} \phi_{c_1} \\ \vdots \\ \phi_{c_m} \end{bmatrix}$$

where  $\mu^{-1}(i) = \{b_1, \dots, b_m\}$  and  $\nu^{-1}(j) = \{c_1, \dots, c_m\}$ . Observe that  $f_\gamma$  is determined only up to a sign by the data  $\gamma$ .

But then, we also have a spanning set for the multilinear  $SL(\alpha)$ -invariants on

$$rep_\alpha Q = \bigotimes_{\textcircled{v}} \left[ \bigotimes_{\textcircled{v} \leftarrow \textcircled{v}} \mathbb{C}^{a_v} \otimes \bigotimes_{\textcircled{v} \leftarrow \textcircled{v}} \mathbb{C}^{*a_v} \right]$$

determined by quintuples  $\Gamma = (\mu, \nu, I, J, K)$  where we have disjoint index-sets partitioned over the vertices  $v \in \{v_1, \dots, v_k\}$  of  $Q$

$$\begin{cases} I & = \bigsqcup_v I_v \\ J & = \bigsqcup_v J_v \\ K & = \bigsqcup_v K_v \end{cases}$$

together with surjective maps from the set of all arrows  $A$  of  $Q$

$$\begin{cases} A & \xrightarrow{\mu} I \sqcup K \\ A & \xrightarrow{\nu} J \sqcup K \end{cases}$$

where we have for every arrow  $\textcircled{w} \xleftarrow{a} \textcircled{v}$  that

$$\begin{cases} \mu(a) & \in I_v \sqcup K_v \\ \nu(a) & \in J_w \sqcup K_w \end{cases}$$

and these maps  $\mu$  and  $\nu$  are subject to the numerical restrictions

$$\begin{cases} \# \mu^{-1}(k) = 1 = \# \nu^{-1}(k) & \text{for all } k \in K. \\ \# \mu^{-1}(i) = a_v = \# \nu^{-1}(j) & \text{for all } i \in I_v \text{ and all } j \in J_v. \end{cases}$$

Such a quintuple  $\Gamma = (\mu, \nu, I, J, K)$  determines for every vertex  $v$  a quintuple

$$\gamma_v = (\mu_v = \mu \mid \{ \textcircled{w} \xleftarrow{a} \textcircled{v} \}, \nu_v = \nu \mid \{ \textcircled{v} \xleftarrow{a} \textcircled{w} \}, I_v, J_v, K_v)$$

satisfying the necessary numerical restrictions to define the  $SL_{a_v}$ -invariant  $f_{\gamma_v}$  described before. Then, the multilinear  $SL(\alpha)$ -invariant on  $rep_\alpha Q$  determined by  $\Gamma$  is defined to be

$$f_\Gamma = \prod_v f_{\gamma_v}$$

and we have to show that these semi-invariants lie in the linear span of the determinantal semi-invariants.

First, consider the case where the index set  $K$  is empty. If we denote the total number of arrows in  $Q$  by  $n$ , then the numerical restrictions imposed give us two expressions for  $n$

$$\sum_v a_v \cdot \# I_v = n = \sum_v a_v \cdot \# J_v$$

Every arrow  $\textcircled{w} \xleftarrow{a} \textcircled{v}$  determines a pair of indices  $\mu(a) \in I_v$  and  $\nu(a) \in J_w$ . To the quintuple  $\Gamma$  we assign a map  $\Phi_\Gamma$  in  $\mathbf{add} Q$


$$\textcircled{1}^{\oplus I_1} \oplus \dots \oplus \textcircled{k}^{\oplus I_k} \xrightarrow{\Phi_\Gamma} \textcircled{1}^{\oplus J_1} \oplus \dots \oplus \textcircled{k}^{\oplus J_k}$$

which decomposes as a block-matrix in blocks  $M_{v,w} \in Hom(\textcircled{v}^{\oplus I_v}, \textcircled{w}^{\oplus J_w})$  of which the  $(i, j)$  entry is given by the sum of arrows

$$\sum_{\substack{\mu(a)=i \\ \nu(a)=j}} \textcircled{w} \xleftarrow{a} \textcircled{v}$$

For a representation  $V \in rep_\alpha Q$ ,  $V(\Phi_\Gamma)$  is an  $n \times n$  matrix and the determinant defines the determinantal semi-invariant  $P_{\Phi_\Gamma, \Gamma}$  which we claim to be equal to the basic invariant  $f_\Gamma$  possibly up to a sign.

We introduce a new quiver situation. Let  $Q'$  be the quiver with vertices the elements of  $I \sqcup J$  and with arrows the set  $A$  of arrows of  $Q$ , but this time we take the starting point of the arrow  $\textcircled{w} \xleftarrow{a} \textcircled{v}$  in  $Q$  to be  $\mu(a) \in I$  and the terminating vertex to be  $\nu(a) \in J$ . That is,  $Q'$  is a bipartite quiver



On  $Q'$  we have the quintuple  $\Gamma' = (\mu', \nu', I', J', K')$  where  $K' = \emptyset$ ,

$$I' = \bigsqcup_{i \in I} I'_i = \bigsqcup_{i \in I} \{i\} \quad J' = \bigsqcup_{j \in J} J'_j = \bigsqcup_{j \in J} \{j\}$$

and  $\mu' = \mu, \nu' = \nu$ . We define an additive functor  $\text{add } Q' \xrightarrow{s} \text{add } Q$  by

$$\textcircled{i} \xrightarrow{s} \textcircled{v} \quad \textcircled{j} \xrightarrow{s} \textcircled{w} \quad \textcircled{v} \xleftarrow{a} \textcircled{w} \xrightarrow{s} \textcircled{v} \xleftarrow{a} \textcircled{w}$$

for all  $i \in I_v$  and all  $j \in J_w$ . The functor  $s$  induces a functor  $\text{rep } Q \xrightarrow{s} \text{rep } Q'$  defined by  $V \xrightarrow{s} V \circ s$ . If  $V \in \text{rep}_\alpha Q$  then  $s(V) \in \text{rep}_{\alpha'} Q'$  where

$$\alpha' = (\underbrace{c_1, \dots, c_p}_{\# I}, \underbrace{d_1, \dots, d_q}_{\# J}) \quad \text{with} \quad \begin{cases} c_i = a_v & \text{if } i \in I_v \\ d_j = a_w & \text{if } j \in J_w \end{cases}$$

That is, the characteristic feature of  $Q'$  is that every vertex  $i \in I$  is the source of exactly  $c_i$  arrows (follows from the numerical condition on  $\mu$ ) and that every vertex  $j \in J$  is the sink of exactly  $d_j$  arrows in  $Q'$ . That is, locally  $Q'$  has the following form

$$\textcircled{c} \longrightarrow \boxed{c} \longrightarrow \quad \text{or} \quad \longrightarrow \boxed{d} \longrightarrow \textcircled{d}$$

There are induced maps

$$\text{rep}_\alpha Q \xrightarrow{s} \text{rep}_{\alpha'} Q' \quad GL(\alpha) \xrightarrow{s} GL(\alpha')$$

where the latter follows from functoriality by considering  $GL(\alpha)$  as the automorphism group of the trivial representation in  $\text{rep}_\alpha Q$ . These maps are compatible with the actions as one checks that  $s(g.V) = s(g).s(V)$ . Also  $s$  induces a map on the coordinate rings  $\mathbb{C}[\text{rep}_\alpha Q] \xrightarrow{s} \mathbb{C}[\text{rep}_{\alpha'} Q']$  by  $s(f) = f \circ s$ . In particular, for the determinantal semi-invariants we have

$$s(P_{\alpha', \phi'}) = P_{\alpha, s(\phi')}$$

and from the compatibility of the action it follows that when  $f$  is a semi-invariant the  $GL(\alpha')$  action on  $\text{rep}_{\alpha'} Q'$  with character  $\chi'$ , then  $s(f)$  is a semi-invariant for the  $GL(\alpha)$ -action on  $\text{rep}_\alpha Q$  with character  $s(\chi) = \chi' \circ s$ . In particular we have that

$$s(P_{\alpha', \Phi_{\Gamma'}}) = P_{\alpha, s(\Phi_{\Gamma'})} = P_{\alpha, \Phi_\Gamma} \quad \text{and} \quad s(f_{\Gamma'}) = f_\Gamma$$

Hence in order to prove our claim, we may replace the triple  $(Q, \alpha, \Gamma)$  by the triple  $(Q', \alpha', \Gamma')$ . We will do this and forget the dashes from here on.

In order to verify that  $f_\Gamma = \pm P_{\alpha, \Phi_\Gamma}$  it suffices to check this equality on the image of

$$W = \bigoplus_{\textcircled{i} \xrightarrow{a} \textcircled{j}} \mathbb{C}^{c_i} \oplus \mathbb{C}^{*d_j} \quad \text{in} \quad \bigotimes_{\textcircled{i} \xrightarrow{a} \textcircled{j}} \mathbb{C}^{c_i} \otimes \mathbb{C}^{*d_j}$$

One verifies that both  $f_\Gamma$  and  $P_{\alpha, \Phi_\Gamma}$  are  $GL(\alpha)$ -semi-invariants on  $W$  of weight  $\chi_\theta$  where

$$\theta = (\underbrace{1, \dots, 1}_{\# I}, \underbrace{-1, \dots, -1}_{\# J})$$

Using the characteristic local form of  $Q = Q'$ , we see that  $W$  is isomorphic to the  $GL(\alpha)$ -module

$$W \simeq \bigoplus_{i \in I} (\underbrace{\mathbb{C}^{c_i} \oplus \dots \oplus \mathbb{C}^{c_i}}_{c_i}) \oplus \bigoplus_{j \in J} (\underbrace{\mathbb{C}^{*d_j} \oplus \dots \oplus \mathbb{C}^{*d_j}}_{d_j}) \simeq \bigoplus_{i \in I} M_{c_i}(\mathbb{C}) \oplus \bigoplus_{j \in J} M_{d_j}(\mathbb{C})$$

and the  $i$  factors of  $GL(\alpha)$  act by inverse right-multiplication on the component  $M_{c_i}$  (and trivially on all others) and the  $j$  factors act by left-multiplication on the


component  $M_{d_j}$  (and trivially on the others). That is,  $GL(\alpha)$  acts on  $W$  with an open orbit, say that of the element

$$w = (\mathbb{1}_{c_1}, \dots, \mathbb{1}_{c_p}, \mathbb{1}_{d_1}, \dots, \mathbb{1}_{d_q}) \in W$$

One verifies immediately from the definitions that that both  $f_\Gamma$  and  $P_{\alpha, \Phi_\Gamma}$  evaluate to  $\pm 1$  in  $w$ . Hence, indeed,  $f_\Gamma$  can be expressed as a determinantal semi-invariant.

Remains to consider the case when  $K$  is non-empty. For  $k \in K$  two situations can occur

- $\mu^{-1}(k) = a$  and  $\nu^{-1}(k) = b$  are distinct, then  $k$  corresponds to replacing the arrows  $a$  and  $b$  by their concatenation




- $\mu^{-1}(k) = a = \nu^{-1}(k)$  then  $a$  is a loop in  $Q$  and  $k$  corresponds



to taking the trace of  $a$ .


This time we construct a new quiver  $Q''$  with vertices  $\{w_1, \dots, w_n\}$  corresponding to the set  $A$  of arrows in  $Q$ . The arrows in  $Q''$  will correspond to elements of  $K$ , that is if  $k \in K$  we have the arrow (or loop) in  $Q''$  with notations as before



We consider the connected components of  $Q''$ . They are of the following three types

- (oriented cycle) : To an oriented cycle  $C$  in  $Q''$  corresponds an oriented cycle  $C'_C$  in the original quiver  $Q$ . We associate to it the trace  $tr(C'_C)$  of this cycle.
- (open paths) : An open path  $P$  in  $Q''$  corresponds to an oriented path  $P'_P$  in  $Q$  which may be a cycle. To  $P$  we associate the corresponding path  $P'_P$  in  $Q$ .
- (isolated points) : They correspond to arrows in  $Q$ .

We will now construct a new quiver  $Q'$  having the same vertex set  $\{v_1, \dots, v_k\}$  as  $Q$  but with arrows corresponding to the set of paths  $P'_P$  described above. The starting and ending vertex of the arrow corresponding to  $P'_P$  are of course the starting and ending vertex of the path  $P_P$  in  $Q$ . Again, we define an additive functor  $\text{add } Q' \xrightarrow{s} \text{add } Q$  by the rules



If the path  $P'_P$  is the concatenation of the arrows  $a_d \circ \dots \circ a_1$  in  $Q$ , we define the maps

$$\begin{cases} \mu'(P'_P) = \mu(a_1) \\ \nu'(P'_P) = \nu(a_d) \end{cases} \quad \text{whence} \quad \begin{cases} \{P'_P\} \xrightarrow{\mu} I' \\ \{P'_P\} \xrightarrow{\nu} J' \end{cases}$$




FIGURE 1. Left-right bipartite quiver.

that is, a quintuple  $\Gamma' = (\mu', \nu', I', J', K' = \emptyset)$  for the quiver  $Q'$ . One then verifies that

$$\begin{aligned} f_{\Gamma} &= s(f_{\Gamma'}) \prod_C tr(C'_C) = s(P_{\alpha, \Phi_{\Gamma'}}) \prod_C tr(C'_C) \\ &= P_{\alpha, s(\Phi_{\Gamma'})} \prod_C tr(C'_C) \end{aligned}$$

finishing the proof of the fact that multilinear semi-invariants lie in the linear span of determinantal semi-invariants (and traces of oriented cycles).

The arguments above can be reformulated in a more combinatorial form which is often useful in constructing semi-invariants of a specific weight, as is necessary in the study of the moduli spaces  $M_{\alpha}^{ss}(Q, \theta)$ . Let  $Q$  be a quiver on the vertices  $\{v_1, \dots, v_k\}$ , fix a dimension vector  $\alpha = (a_1, \dots, a_k)$  and a character  $\chi_{\theta}$  where  $\theta = (t_1, \dots, t_k)$  such that  $\theta(\alpha) = 0$ . We will call a bipartite quiver  $Q'$  as in figure 1 on left vertex-set  $L = \{l_1, \dots, l_p\}$  and right vertex-set  $R = \{r_1, \dots, r_q\}$  and a dimension vector  $\beta = (c_1, \dots, c_p; d_1, \dots, d_q)$  to be of type  $(Q, \alpha, \theta)$  if the following conditions are met

- All left and right vertices correspond to vertices of  $Q$ , that is, there are maps

$$\begin{cases} L & \xrightarrow{l} \{v_1, \dots, v_k\} \\ R & \xrightarrow{r} \{v_1, \dots, v_k\} \end{cases}$$

possibly occurring with multiplicities, that is there is a map

$$L \cup R \xrightarrow{m} \mathbb{N}_+$$

such that  $c_i = m(l_i)a_z$  if  $l(l_i) = v_z$  and  $d_j = m(r_j)a_z$  if  $r(r_j) = v_z$ .

- There can only be an arrow  $(l_i) \longrightarrow (r_j)$  if for  $v_k = l(l_i)$  and  $v_l = r(r_j)$  there is an oriented path



in  $Q$  allowing the trivial path and loops if  $v_k = v_l$ .

- Every left vertex  $l_i$  is the source of exactly  $c_i$  arrows in  $Q'$  and every right-vertex  $r_j$  is the sink of precisely  $d_j$  arrows in  $Q'$ .

- Consider the  $u \times u$  matrix where  $u = \sum_i c_i = \sum_j d_j$  (both numbers are equal to the total number of arrows in  $Q'$ ) where the  $i$ -th row contains the entries of the  $i$ -th arrow in  $Q'$  with respect to the obvious left and right bases. Observe that this is a  $GL(\beta)$  semi-invariant on  $rep_\beta Q'$  with weight determined by the integral  $k + l$ -tuple  $(-1, \dots, -1; 1, \dots, 1)$ . If we fix for every arrow  $a$  from  $l_i$  to  $r_j$  in  $Q'$  an  $m(r_j) \times m(l_i)$  matrix  $p_a$  of linear combinations of paths in  $Q$  from  $l(l_i)$  to  $r(r_j)$ , we obtain a morphism

$$rep_\alpha Q \longrightarrow rep_\beta Q'$$

sending a representation  $V \in rep_\alpha Q$  to the representation  $W$  of  $Q'$  defined by  $W_a = p_a(V)$ . Composing this map with the above semi-invariant we obtain a  $GL(\alpha)$  semi-invariant of  $rep_\alpha Q$  with weight determined by the  $k$ -tuple  $\theta = (t_1, \dots, t_k)$  where

$$t_i = \sum_{j \in r^{-1}(v_i)} m(r_j) - \sum_{j \in l^{-1}(v_i)} m(l_j)$$

We call such semi-invariants *standard determinantal*. Summarizing the arguments of this section we have proved after applying polarization and restitution processes

**THEOREM 7.5.** *The semi-invariants of the  $GL(\alpha)$ -action on  $rep_\alpha Q$  are generated by traces of oriented cycles and by standard determinantal semi-invariants.*


**7.3. Universal localization.**

In order to prove that the moduli spaces  $M_\alpha^{ss}(Q, \theta)$  are locally controlled by Quillen-smooth algebras, we need to recall the notion of *universal localization*. We refer to the monograph by A. Schofield [69] for full details.

Let  $A$  be a  $\mathbb{C}$ -algebra and  $\text{projmod } A$  the category of finitely generated projective left  $A$ -modules. Let  $\Sigma$  be some class of maps in this category (that is some left  $A$ -module morphisms between certain projective modules). Then, there exists an algebra map  $A \xrightarrow{j_\Sigma} A_\Sigma$  with *the universal property* that the maps  $A_\Sigma \otimes_A \sigma$  have an inverse for all  $\sigma \in \Sigma$ .  $A_\Sigma$  is called the *universal localization* of  $A$  with respect to the set of maps  $\Sigma$ .


**PROPOSITION 7.6.** *When  $A$  is Quillen-smooth, then so is  $A_\Sigma$ .*

**PROOF.** Consider a test-object  $(T, I)$  in  $\mathbf{alg}$ , then we have the following diagram



where  $\psi$  exists by Quillen-smoothness of  $A$ . By *Nakayama's lemma* all maps  $\sigma \in \Sigma$  become isomorphisms under tensoring with  $\psi$ . Then,  $\phi$  exists by the universal property of  $A_\Sigma$ . □

Consider the special case when  $A$  is the path algebra  $\mathbb{C}Q$  of a quiver on  $k$  vertices. Then, we can identify the isomorphism classes in  $\text{projmod } \mathbb{C}Q$  with the opposite category of  $\text{add } Q$  introduced in the foregoing section. To each vertex  $v_i$  corresponds an *indecomposable projective* left  $\mathbb{C}Q$ -ideal  $P_i = \mathbb{C}Qe_i$  having as  $\mathbb{C}$ -vector-space basis all paths in  $Q$  starting at  $v_i$ . For the homomorphisms we have

$$\text{Hom}_{\mathbb{C}Q}(P_i, P_j) = \bigoplus_{\text{paths } p} \mathbb{C}p = \text{Hom}_{\text{add } Q}(\textcircled{j}, \textcircled{i})$$


where  $p$  is an oriented path in  $Q$  starting at  $v_j$  and ending at  $v_i$ . Therefore, any  $A$ -module morphism  $\sigma$  between two projective left modules

$$P_{i_1} \oplus \dots \oplus P_{i_u} \xrightarrow{\sigma} P_{j_1} \oplus \dots \oplus P_{j_v}$$

can be represented by an  $u \times v$  matrix  $M_\sigma$  whose  $(p, q)$ -entry  $m_{pq}$  is a linear combination of oriented paths in  $Q$  starting at  $v_{j_q}$  and ending at  $v_{i_p}$ .

Now, form an  $v \times u$  matrix  $N_\sigma$  of free variables  $y_{pq}$  and consider the algebra  $\mathbb{C}Q_\sigma$  which is the quotient of the free product  $\mathbb{C}Q * \mathbb{C}\langle y_{11}, \dots, y_{uv} \rangle$  modulo the ideal of relations determined by the matrix equations

$$M_\sigma \cdot N_\sigma = \begin{bmatrix} v_{i_1} & & 0 \\ & \ddots & \\ 0 & & v_{i_u} \end{bmatrix} \quad N_\sigma \cdot M_\sigma = \begin{bmatrix} v_{j_1} & & 0 \\ & \ddots & \\ 0 & & v_{j_v} \end{bmatrix}$$

Equivalently,  $\mathbb{C}Q_\sigma$  is the path algebra of a quiver *with relations* where the quiver is  $Q$  extended with arrows  $y_{pq}$  from  $v_{i_p}$  to  $v_{j_q}$  for all  $1 \leq p \leq u$  and  $1 \leq q \leq v$  and the relations are the above matrix entry relations.

Repeating this procedure for every  $\sigma \in \Sigma$  we obtain the universal localization  $\mathbb{C}Q_\Sigma$ . This proves

**PROPOSITION 7.7.** *If  $\Sigma$  is a finite set of maps, then the universal localization  $\mathbb{C}Q_\Sigma$  is an affine  $\mathbb{C}$ -algebra.*

It is easy to verify that the representation space  $\text{rep}_n \mathbb{C}Q_\sigma$  is an affine Zariski open subscheme (but possibly empty) of  $\text{rep}_n \mathbb{C}Q$ . Indeed, if  $V = (V_a)_a \in \text{rep}_\alpha Q$ , then  $V$  determines a point in  $\text{rep}_n \mathbb{C}Q_\Sigma$  if and only if the matrices  $M_\sigma(V)$  in which the arrows are all replaced by the matrices  $V_a$  are invertible for all  $\sigma \in \Sigma$ .

In particular, this induces numerical conditions on the dimension vectors  $\alpha$  such that  $\text{rep}_\alpha Q_\Sigma \neq \emptyset$ . Let  $\alpha = (a_1, \dots, a_k)$  be a dimension vector such that  $\sum a_i = n$  then every  $\sigma \in \Sigma$  say with

$$P_1^{\oplus e_1} \oplus \dots \oplus P_k^{\oplus e_k} \xrightarrow{\sigma} P_1^{\oplus f_1} \oplus \dots \oplus P_k^{\oplus f_k}$$

gives the numerical condition

$$e_1 a_1 + \dots + e_k a_k = f_1 a_1 + \dots + f_k a_k.$$

These numerical restrictions will be used to relate  $\theta$ -stable representations of  $Q$  to simple representations of universal localizations of  $\mathbb{C}Q$ .

Fix a character  $\theta = (t_1, \dots, t_k) \in \mathbb{Z}^k$  and divide the set of indices  $1 \leq i \leq k$  into the *left set*  $L = \{i_1, \dots, i_u\}$  consisting of those  $i$  such that  $t_i \leq 0$  and the *right set*  $R = \{j_1, \dots, j_v\}$  consisting of those  $j$  such that  $t_j \geq 0$ . Consider a dimension

vector  $\alpha$  such that  $\theta.\alpha = 0$ , then  $\theta$  determines the *character*

$$GL(\alpha) \xrightarrow{\chi_\theta} \mathbb{C}^* \quad (g_1, \dots, g_k) \mapsto \prod_i \det(g_i)^{t_i}$$

Next, consider the sets of morphisms

$$\Sigma_\theta = \bigcap_{z \in \mathbb{N}_+} \Sigma_\theta(z)$$

where  $\Sigma_\theta(z)$  is the set of all morphisms

$$P_{i_1}^{\oplus -zt_{i_1}} \oplus \dots \oplus P_{i_u}^{\oplus -zt_{i_u}} \xrightarrow{\sigma} P_{j_1}^{\oplus zt_{j_1}} \oplus \dots \oplus P_{j_v}^{\oplus zt_{j_v}}$$

With notation as before, it follows that

$$d_\sigma(V) = \det M_\sigma(V) \quad V \in \text{rep}_\alpha Q$$

is a semi-invariant on  $\text{rep}_\alpha Q$  of weight  $z\chi_\theta$ . This semi-invariant determines the Zariski open subset of  $\text{rep}_\alpha Q$

$$X_\sigma(\alpha) = \{V \in \text{rep}_\alpha Q \mid d_\sigma(V) \neq 0\}$$

It is clear from the results of section 6.3 that  $X_\sigma(\alpha)$  consists of  $\theta$ -semistable representations. We can characterize the  $\theta$ -stable representations in this open set.

LEMMA 7.8. *For  $V \in X_\sigma(\alpha)$  the following are equivalent*

- (1)  *$V$  is a  $\theta$ -stable representation.*
- (2)  *$V$  is a simple  $\alpha$ -dimensional representation of the universal localization  $\mathbb{C}Q_\sigma$ .*

PROOF. Let  $W$  be a  $\beta$ -dimensional subrepresentation of  $V$  with  $\beta = (b_1, \dots, b_k)$ , then for  $W$  to be a  $\beta$ -dimensional representation of the universal localization  $\mathbb{C}Q_\sigma$  it must satisfy the numerical restriction

$$-t_{i_1}b_{i_1} - \dots - t_{i_u}b_{i_u} = t_{j_1}b_{j_1} + \dots + t_{j_v}b_{j_v} \quad \text{that is} \quad \theta.\beta = 0$$

Hence, if  $V$  is  $\theta$ -stable, there are no proper subrepresentations of  $V$  as a  $\mathbb{C}Q_\sigma$ -representation. Conversely, if  $V$  is an  $\alpha$ -dimensional subrepresentation of  $\mathbb{C}Q_\sigma$  we must have that  $d_\sigma(V) \neq 0$ . But then, if  $W$  is a  $\beta$ -dimensional  $Q$ -subrepresentation of  $V$  we must have that  $\sum_a -t_{i_a}b_{i_a} \leq \sum_b t_{j_b}b_{j_b}$  (if not,  $\sigma(V)$  would have a kernel) whence  $\theta.\beta \geq 0$ . If  $W$  is a subrepresentation such that  $\theta.\beta = 0$ , then  $W$  would be a proper  $\mathbb{C}Q_\sigma$  subrepresentation of  $V$ , a contradiction. Therefore,  $V$  is  $\theta$ -stable.  $\square$

THEOREM 7.9. *The moduli space of  $\theta$ -semistable representations of the quiver  $Q$*

$$M_\alpha^{ss}(Q, \theta)$$

*is locally controlled by the set of Quillen-smooth algebras  $\{\mathbb{C}Q_\sigma \mid \sigma \in \Sigma_\theta\}$ .*

PROOF. By the results of the foregoing section we know that the quotient varieties of the Zariski open affine subsets  $X_\sigma(\alpha)$  cover the moduli space  $M_\alpha^{ss}(Q, \theta)$ . Further, by lemma 7.8 we have a canonical isomorphism

$$X_\sigma(\alpha)/GL(\alpha) \simeq \text{iss}_\alpha \mathbb{C}Q_\sigma$$

Finally, because

$$\text{rep}_n \mathbb{C}Q_\sigma = \sqcup_\alpha GL_n \times^{GL(\alpha)} \text{rep}_\alpha \mathbb{C}Q_\sigma$$

where the disjoint union is taken over all  $\alpha = (a_1, \dots, a_k)$  such that  $\sum_i a_i = n$ , we have that  $\text{iss}_\alpha \mathbb{C}Q_\sigma$  is an irreducible component of  $\text{iss}_n \mathbb{C}Q_\sigma$  finishing the proof.  $\square$



Lemma 7.8 also allows us to study the moduli spaces  $M_\alpha^{ss}(Q, \theta)$  locally by the local quiver settings associated to semi-simple representations. That is, let  $\xi \in M_\alpha^{ss}(Q, \theta)$  be the point corresponding to

$$M_\xi = S_1^{\oplus e_1} \oplus \dots \oplus S_z^{\oplus e_z}$$

where  $S_i$  is a  $\theta$ -stable representation of dimension vector  $\beta_i$  occurring in  $M_\xi$  with multiplicity  $e_i$ .

**THEOREM 7.10.** *With notations as above, the étale local structure of the moduli space  $M_\alpha^{ss}(Q, \theta)$  near  $\xi$  is that of the quotient variety  $\text{iss}_\beta Q_\xi$  where  $\beta = (e_1, \dots, e_z)$  and  $Q_\xi$  is the quiver on  $z$  vertices such that*

$$\begin{cases} \# \textcircled{i} \xleftarrow{a} \textcircled{j} & = -\chi_Q(\beta_i, \beta_j) \\ \# \textcircled{i} & = 1 - \chi_Q(\beta_i, \beta_i) \end{cases}$$

near the trivial representation.


**PROOF.** In view of the above results and the slice theorems, we only have to compute the ext-spaces  $\text{Ext}_{\mathbb{C}Q_\sigma}^1(S_i, S_j)$ . From [69, Thm. 4.7] we recall that the category of  $\mathbb{C}Q_\sigma$  representations is closed under extensions in the category of representations of  $Q$ . Therefore, we have for all  $\mathbb{C}Q_\sigma$ -representations  $V$  and  $W$  that

$$\text{Ext}_{\mathbb{C}Q}^1(V, W) \simeq \text{Ext}_{\mathbb{C}Q_\sigma}^1(V, W)$$

from which the result follows using theorem 3.29. □

In the following section we will give some applications of this result. Universal localizations can also be used to determine the *formal structure* on representation spaces of quivers.

Let  $Q$  be a quiver on  $k$  vertices and consider the extended quiver  $Q^{(n)}$



That is, we add to the vertices and arrows of  $Q$  one extra vertex  $v_0$  and for every vertex  $v_i$  in  $Q$  we add  $n$  directed arrows from  $v_0$  to  $v_i$ . We will denote the  $j$ -th arrow  $1 \leq j \leq n$  from  $v_0$  to  $v_i$  by  $x_{ij}$ .

Consider the morphism between projective left  $\mathbb{C}Q^{(n)}$ -modules

$$P_1 \oplus P_2 \oplus \dots \oplus P_k \xrightarrow{\sigma} \underbrace{P_0 \oplus \dots \oplus P_0}_n$$

determined by the matrix

$$M_\sigma = \begin{bmatrix} x_{11} & \dots & \dots & x_{1n} \\ \vdots & & & \vdots \\ x_{k1} & \dots & \dots & x_{kn} \end{bmatrix}.$$

We consider the universal localization  $\mathbb{C}Q_\sigma^{(n)}$ , that is, we add for each vertex  $v_i$  in  $Q$  another  $n$  arrows  $y_{ij}$  with  $1 \leq j \leq n$  from  $v_i$  to  $v_0$ .


With these arrows  $y_{ij}$  one forms the  $n \times k$  matrix

$$N_\sigma = \begin{bmatrix} y_{11} & \cdots & y_{k1} \\ \vdots & & \vdots \\ \vdots & & \vdots \\ y_{1n} & \cdots & y_{kn} \end{bmatrix}$$

and the universal localization  $\mathbb{C}Q_\sigma^{(n)}$  is described by the relations

$$M_\sigma.N_\sigma = \begin{bmatrix} v_1 & & 0 \\ & \ddots & \\ 0 & & v_k \end{bmatrix} \quad \text{and} \quad N_\sigma.M_\sigma = \begin{bmatrix} v_0 & & & 0 \\ & \ddots & & \\ & & \ddots & \\ 0 & & & v_1 \end{bmatrix}.$$

We will depict this quiver with relations by the picture  $Q_\sigma^{(n)}$



From the discussion above it follows that there is a canonical isomorphism

$$rep_m \sqrt[n]{\mathbb{C}Q} \simeq rep_m \mathbb{C}Q_\sigma^{(n)}.$$

In fact we can even identify

$$\sqrt[n]{\mathbb{C}Q} = v_0 \mathbb{C}Q_\sigma^{(n)} v_0.$$

Indeed, the right hand side is generated by all the oriented cycles in  $Q_\sigma^{(n)}$  starting and ending at  $v_0$  and is therefore generated by the  $y_{ip}x_{iq}$  and the  $y_{ip}ax_{jq}$  where  $a$  is an arrow in  $Q$  starting in  $v_j$  and ending in  $v_i$ . If we have an algebra morphism

$$\mathbb{C}Q \xrightarrow{\phi} M_n(B)$$

then we have an associated algebra morphism


$$v_0 \mathbb{C}Q_\sigma^{(n)} v_0 \xrightarrow{\psi} B$$

defined by sending  $y_{ip}ax_{jq}$  to the  $(p, q)$ -entry of the  $n \times n$  matrix  $\phi(a)$  and  $y_{ip}x_{iq}$  to the  $(p, q)$ -entry of  $\phi(v_i)$ . The defining relations among the  $x_{ip}$  and  $y_{iq}$  introduced before imply that  $\psi$  is indeed an algebra morphism.

EXAMPLE 7.11. Let  $A = \mathbb{C}\langle a, b \rangle$ , that is  $A$  is the path algebra of the quiver



In order to describe  $\sqrt[n]{A}$  we consider the quiver with relations



We see that the algebra of oriented cycles in  $v_0$  in this quiver with relations is isomorphic to the free algebra in  $2n^2$  free variables


$$\mathbb{C}\langle y_1 a x_1, \dots, y_n a x_n, y_1 b x_1, \dots, y_n b x_n \rangle$$

which coincides with our knowledge of  $\sqrt[n]{\mathbb{C}\langle a, b \rangle}$ .

There is some elementary calculus among the  $n$ -th roots of algebras. For example, it follows from the universal property of  $\sqrt[n]{A}$  that there is a natural morphism

$$\sqrt[k_1]{\sqrt[k_2]{\dots \sqrt[k_z]{A}}} \longleftarrow \sqrt[k]{A}$$

where  $k = \prod k_i$ . When  $A = \mathbb{C}Q$  we can represent this morphism graphically by the picture



where the map is given by composing paths from  $v_0$  to  $v_i$ . Also observe that we used the isomorphisms in the rightmost part of the left quiver to remove additional arrows from the extra vertices to  $v_i$  at each stage.

Probably more important are the *connecting morphisms*

$$\sqrt[k_1]{A} * \sqrt[k_2]{A} * \dots * \sqrt[k_z]{A} \xleftarrow{c(k_1, \dots, k_z)} \sqrt[k]{A}$$

with  $k = \sum k_i$  obtained from the universal property of  $\sqrt[n]{A}$  by composing algebra morphisms  $A \xrightarrow{\phi_i} M_{k_i}(B)$  to an algebra morphism

$$A \xrightarrow{\begin{bmatrix} \phi_1 & & 0 \\ & \ddots & \\ 0 & & \phi_z \end{bmatrix}} M_k(B).$$

Observing that the ordering of the factors is important (but only up to isomorphism of the representations).




FIGURE 2. Free product of quivers.

We need to have a quiver interpretation of the *free product*  $\mathbb{C}Q_1 * \mathbb{C}Q_2$  of two path algebras (at least as far as finite dimensional representations are concerned). Let  $Q_1$  be a quiver on  $k$  vertices  $\{v_1, \dots, v_k\}$  and  $Q_2$  a quiver on  $p$  vertices  $\{w_1, \dots, w_p\}$  and consider the extended quiver  $Q_1 * Q_2$  of figure 2. That is, we add one extra arrow from each vertex of  $Q_1$  to each arrow of  $Q_2$ .

Let  $\{P_1, \dots, P_k\}$  be the projective left  $\mathbb{C}Q_1 * \mathbb{C}Q_2$ -modules corresponding to the vertices of  $Q_1$  and  $\{P'_1, \dots, P'_p\}$  those corresponding to the vertices of  $Q_2$  and consider the morphism

$$P'_1 \oplus \dots \oplus P'_p \xrightarrow{\sigma} P_1 \oplus \dots \oplus P_k$$

determined by the  $p \times k$  matrix

$$M_\sigma = \begin{bmatrix} x_{11} & \dots & x_{1k} \\ \vdots & & \vdots \\ x_{p1} & \dots & x_{pk} \end{bmatrix}$$


where  $x_{ij}$  denotes the extra arrow from vertex  $v_j$  to vertex  $w_i$ .

Let  $Q_1 * Q_{2,\sigma}$  denote the quiver with relations one obtains by inverting this map (as above). Then, it is fairly easy to see that

$$rep_n \mathbb{C}Q_1 * Q_2 \simeq rep_n Q_1 * Q_{2,\sigma}$$

where the right-hand side denote the subscheme of  $n$ -dimensional representations of the quiver  $Q_1$  times the  $n$ -dimensional representations of  $Q_2$  where the extra arrows determine an isomorphism of the representations.

Using this interpretation of the free product one can now give a graphical interpretation of the connecting morphisms in the case of the two loop quiver (the general case is similar).



obtained by 'grafting' the bottom tree. Observe that again we used the isomorphisms given by the  $k_i$  bundles to eliminate adding extra arrows in the free products.

**7.4. Compact manifolds.**

$\mathbf{geo} @_{\mathbf{n}}$  is the study of *families* of algebraic varieties (with specified connecting morphisms) which are local controlled by a set of noncommutative algebras. If this set of algebras consists of Quillen-smooth algebras we say that the family of varieties is a *noncommutative manifold*. If all varieties in the family are in addition projective (possibly with singularities) we say that the family is a *compact noncommutative manifold*.

So far, we have not specified the properties of the connecting morphisms. In this section we present a first class of examples, the *sum families*. In the next chapter we will encounter another possibility coming from the theory of completely integrable dynamical systems.

DEFINITION 7.12. A *sum family* is an object  $(X_n)_n$  in  $\mathbf{geo} @_{\mathbf{n}}$  indexed over the positive integers such that for each  $n$  there is a  $GL_n$ -variety  $Y_n$  and a quotient morphism

$$Y_n \twoheadrightarrow Y_n/GL_n \simeq X_n$$

and  $Y_n$  is locally of the form  $rep_n A$  for an affine  $\mathbb{C}$ -algebra  $A$  belonging to a set  $\mathcal{A}$  of algebras. Moreover, there are *equivariant connecting sum-maps*

$$Y_m \times Y_n \xrightarrow{\oplus} Y_{m+n}$$

for all  $m, n \in \mathbb{N}_+$  where equivariance means with respect to the group  $GL_m \times GL_n$  embedded diagonally in  $GL_{m+n}$ . If the set  $\mathcal{A}$  consists of Quillen-smooth algebras, we call  $(X_n)_n$  a *sum manifold*.

THEOREM 7.13. For a quiver  $Q$  on  $k$  vertices and a fixed character  $\theta \in \mathbb{Z}^k$ , the family of varieties

$$\left( \bigsqcup_{\substack{\alpha=(a_1, \dots, a_k) \\ \sum_i a_i = n}} M_{\alpha}^{ss}(Q, \theta) \right)_n$$

is a sum manifold in  $\mathbf{geo} @_{\mathbf{n}}$ . If  $Q$  has no oriented cycles, then this family is a compact sum manifold.

PROOF. In view of theorem 7.10 we only need to construct equivariant-sum maps. They are induced from the direct sums of representations

$$rep_{\alpha} Q \times rep_{\beta} Q \xrightarrow{\oplus} rep_{\alpha+\beta} Q \quad (V, W) \mapsto V \oplus W$$

and the required properties are clearly satisfied. □

EXAMPLE 7.14. Let  $M_{\mathbb{P}^2}(n; 0, n)$  be the moduli space of semi-stable vectorbundles of rank  $n$  over the projective plane  $\mathbb{P}^2$  with Chern numbers  $c_1 = 0$  and  $c_2 = n$ . Using results of K. Hulek [31] one can identify this moduli space with

$$M_{\mathbb{P}^2}(n; 0, n) \simeq M_{(n,n)}^{ss}(Q, \theta)$$

where  $Q$  and  $\theta$  are the following quiver-setting



Therefore, the family of moduli spaces  $(M_{\mathbb{P}^2}(n; 0, n))_n$  is a compact sum manifold in  $\mathbf{geo} @_{\mathbf{n}}$ .

Let  $C$  be a smooth projective curve of genus  $g$  and let  $M_C(n, 0)$  be the moduli space of semi-stable vectorbundles of rank  $n$  and degree 0 over  $C$ . We expect that the family of moduli spaces  $(M_C(n, 0))_n$  is a compact sum manifold.

In this section we will investigate another class of examples : representations of the *torus knot groups* . Consider a slid cylinder  $C$  with  $m$  line segments on its curved face, equally spaced and parallel to the axis. If the ends of  $C$  are identified with a twist of  $2\pi\frac{n}{m}$  where  $n$  is an integer relatively prime to  $m$ , we obtain a single curve  $K_{m,n}$  on the surface of a solid torus  $T$ . If we assume that the torus  $T$  lies in  $\mathbb{R}^3$  in the standard way, the curve  $K_{m,n}$  is called the  $(m, n)$  *torus knot* .

Computing the *fundamental group* of the complement  $\mathbb{R}^3 - K_{m,n}$  one obtains the  $(m, n)$ -*torus knot group*


$$\pi_1(\mathbb{R}^3 - K_{m,n}) = G_{m,n} \simeq \langle a, b \mid a^m = b^n \rangle$$

An important example is the *three string braid group*.

EXAMPLE 7.15. Consider Artin's braid group  $B_3$  on three strings.  $B_3$  has the presentation

$$B_3 \simeq \langle L, R \mid LR^{-1}L = R^{-1}LR^{-1} \rangle$$

where  $L$  and  $R$  are the fundamental 3-braids



If we let  $S = LR^{-1}L$  and  $T = R^{-1}L$ , an algebraic manipulation shows that

$$B_3 = \langle S, T \mid T^3 = S^2 \rangle$$

is an equivalent presentation for  $B_3$ . The center of  $B_3$  is the infinite cyclic group generated by the braid

$$Z = S^2 = (LR^{-1}L)^2 = (R^{-1}L)^3 = T^3$$

It follows from the second presentation of  $B_3$  that the quotient group modulo the center is isomorphic to

$$\frac{B_3}{\langle Z \rangle} \simeq \langle s, t \mid s^2 = 1 = t^3 \rangle \simeq \mathbb{Z}_2 * \mathbb{Z}_3$$


the free product of the cyclic group of order 2 (with generator  $s$ ) and the cyclic group of order 3 (with generator  $t$ ). This group is isomorphic to the modular group  $PSL_2(\mathbb{Z})$  via

$$\bar{L} \longrightarrow \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \quad \text{and} \quad \bar{R} \longrightarrow \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$$

It is well known that the modular group  $PSL_2(\mathbb{Z})$  acts on the upper half-plane  $H^2$  by left multiplication in the usual way, that is

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} : H^2 \longrightarrow H^2 \quad \text{given by} \quad z \longrightarrow \frac{az + b}{cz + d}$$

The fundamental domain  $H^2/PSL_2(\mathbb{Z})$  for this action is the hyperbolic triangle






EXAMPLE 7.16. Consider the modular group  $PSL_2(\mathbb{Z}) \simeq \mathbb{Z}_2 * \mathbb{Z}_3$ , the free product of the cyclic groups of order two and three with generators  $\sigma$  resp.  $\tau$ . Let  $S$  be an  $n$ -dimensional simple representation of  $PSL_2(\mathbb{Z})$ . Let  $\xi$  be a 3-rd root of unity, then restricting  $S$  to these finite Abelian subgroups we have

$$\begin{cases} S \downarrow_{\mathbb{Z}_2} \simeq S_1^{\oplus a_1} \oplus S_{-1}^{\oplus a_2} \\ S \downarrow_{\mathbb{Z}_3} \simeq T_1^{\oplus b_1} \oplus T_{\xi}^{\oplus b_2} \oplus T_{\xi^2}^{\oplus b_3} \end{cases}$$

where  $S_x$  resp.  $T_x$  are the one-dimensional representations on which  $\sigma$  resp.  $\tau$  acts via multiplication with  $x$ . Observe that  $a_1 + a_2 = b_1 + b_2 + b_3 = n$  and we associate to  $S$  a representation  $V$  of the quiver situation



with  $V_{1i} = S_i^{\oplus a_i}$  and  $V_{2j} = T_j^{\oplus b_j}$  and where the linear map corresponding to an arrow

$$\textcircled{b_j} \xrightarrow{a_{ij}} \textcircled{a_i}$$

$$V_{a_{ij}} : S_i^{\oplus a_i} \hookrightarrow S \downarrow_{\mathbb{Z}_2} = V \downarrow_{\mathbb{Z}_3} \twoheadrightarrow T_j^{\oplus b_j}$$

of the canonical injections and projections. If  $\alpha = (a_1, a_2, b_1, b_2, b_3)$  then we take as  $\theta = (-1, -1, +1, +1, +1)$ . Observe that  $\oplus_{i,j} V_{a_{ij}} : \mathbb{C}^n \longrightarrow \mathbb{C}^n$  is a linear isomorphism. If  $W \hookrightarrow V$  is a subrepresentation, then  $\theta(W) \geq 0$ . Indeed, if the dimension vector of  $W$  is  $\beta = (c_1, c_2, d_1, d_2, d_3)$  and assume that  $\theta(W) < 0$ , then  $k = c_1 + c_2 > l = d_1 + d_2 + d_3$ , but then the restriction of  $\oplus V_{a_{ij}}$  to  $W$  gives a linear map  $\mathbb{C}^k \longrightarrow \mathbb{C}^l$  having a kernel which is impossible. Hence,  $V$  is a  $\theta$ -semistable representation of the quiver. In fact,  $V$  is even  $\theta$ -stable, for consider a subrepresentation  $W \hookrightarrow V$  with dimension vector  $\beta$  as before and  $\theta(W) = 0$ , that is,  $c_1 + c_2 = d_1 + d_2 + d_3 = m$ , then the isomorphism  $\oplus_{i,j} V_{a_{ij}} \upharpoonright W$  and the decomposition into eigenspaces of  $\mathbb{C}^m$  with respect to the  $\mathbb{Z}_2$  and  $\mathbb{Z}_3$ -action, makes  $\mathbb{C}^m$  into an  $m$ -dimensional representation of  $PSL_2(\mathbb{Z})$  which is a subrepresentation of  $S$ .  $S$  being simple then implies that  $W = V$  or  $W = 0$ , whence  $V$  is  $\theta$ -stable. The underlying reason is that the group algebra  $\mathbb{C}PSL_2(\mathbb{Z})$  is a universal localization of the path algebra  $\mathbb{C}Q$  of the above quiver.

As irreducible  $\overline{G}_{m,n}$ -representations correspond to  $\theta$ -stable representations of the quiver  $Q_{m,n}$  we need to determine the dimension vectors  $\alpha$  of  $\theta$ -stables. In section 6.3 we have given an inductive algorithm to determine them. However, using the fact that the moduli spaces are locally controlled and hence are determined locally by local quivers we can apply the easier classification of simple roots given in section 3.4 so solve this problem.

EXAMPLE 7.17. With  $S_{ij}$  we denote the simple 1-dimensional representation of  $PSL_2(\mathbb{Z})$  determined by


$$S_{ij} \downarrow_{\mathbb{Z}_2} = S_i \quad \text{and} \quad S_{ij} \downarrow_{\mathbb{Z}_3}$$

Let  $n = x_1 + \dots + x_6$  and we aim to study the local structure of  $rep_n \mathbb{C}PSL_2(\mathbb{Z})$  in a neighborhood of the semi-simple  $n$ -dimensional representation

$$V_{\xi} = S_{11}^{\oplus x_1} \oplus S_{12}^{\oplus x_2} \oplus S_{13}^{\oplus x_3} \oplus S_{21}^{\oplus x_4} \oplus S_{22}^{\oplus x_5} \oplus S_{23}^{\oplus x_6}$$



To determine the structure of  $Q_\xi$  we have to compute  $\dim Ext^1(S_{ij}, S_{kl})$ . To do this we view the  $S_{ij}$  as representations of the quiver  $Q_{2,3}$  in the example above. For example  $S_{12}$  is the representation



of dimension vector  $(1, 0; 0, 1, 0)$ . For representations of  $Q_{2,3}$ , the dimensions of  $Hom$  and  $Ext$ -groups are determined by the bilinear form

$$\chi_Q = \begin{bmatrix} 1 & 0 & -1 & -1 & -1 \\ 0 & 1 & -1 & -1 & -1 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$


If  $V \in rep_\alpha Q$  and  $W \in rep_\beta Q$  where  $\alpha = (a_1, a_2; b_1, b_2, b_3)$  with  $a_1 + a_2 = b_1 + b_2 + b_3 = k$  and  $\beta = (c_1, c_2; d_1, d_2, d_3)$  with  $c_1 + c_2 = d_1 + d_2 + d_3 = l$  we have

$$\dim Hom(V, W) - \dim Ext^1(V, W) = \chi_Q(\alpha, \beta) = kl - (a_1c_1 + a_2c_2 + b_1d_1 + b_2d_2 + b_3d_3)$$

As  $Hom(S_{ij}, S_{kl}) = \mathbb{C}^{\oplus \delta_{ik} \delta_{jl}}$  we have that

$$\dim Ext^1(S_{ij}, S_{kl}) = \begin{cases} 1 & \text{if } i \neq k \text{ and } j \neq l \\ 0 & \text{otherwise} \end{cases}$$

But then, the local quiver setting  $(Q_\xi, \alpha_\xi)$  is



We want to determine whether the irreducible component of  $rep_n \mathbb{C}PSL_2(\mathbb{Z})$  containing  $V_\xi$  contains simple  $PSL_2(\mathbb{Z})$ -representations, or equivalently, whether  $\alpha_\xi$  is the dimension vector of a simple representation of  $Q_\xi$ , that is,

$$\chi_{Q_\xi}(\alpha_\xi, \epsilon_j) \leq 0 \quad \text{and} \quad \chi_{Q_\xi}(\epsilon_j, \alpha_\xi) \quad \text{for all } 1 \leq j \leq 6$$

The Euler-form of  $Q_\xi$  is determined by the matrix where we number the vertices cyclicly

$$\chi_{Q_\xi}^\bullet = \begin{bmatrix} 1 & -1 & 0 & 0 & 0 & -1 \\ -1 & 1 & -1 & 0 & 0 & 0 \\ 0 & -1 & 1 & -1 & 0 & 0 \\ 0 & 0 & -1 & 1 & -1 & 0 \\ 0 & 0 & 0 & -1 & 1 & -1 \\ -1 & 0 & 0 & 0 & -1 & 1 \end{bmatrix}$$


leading to the following set of inequalities

$$\begin{cases} x_1 \leq x_5 + x_6 \\ x_2 \leq x_4 + x_6 \\ x_3 \leq x_4 + x_5 \end{cases} \quad \begin{cases} x_4 \leq x_2 + x_3 \\ x_5 \leq x_1 + x_3 \\ x_6 \leq x_1 + x_2 \end{cases}$$

Finally, observe that  $V_\xi$  corresponds to a  $Q_{2,3}$ -representation of dimension vector  $(x_1 + x_2 + x_3, x_4 + x_5 + x_6; x_1 + x_4, x_2 + x_5, x_3 + x_6)$ . If we write this dimension vector as  $(a_1, a_2; b_1, b_2, b_3)$  then the inequalities are equivalent to the conditions

$$a_i \geq b_j \quad \text{for all } 1 \leq i \leq 2 \text{ and } 1 \leq j \leq 3$$


which gives us the desired restriction on the quintuples



at least when  $a_i \geq 3$  and  $b_j \geq 2$ . The remaining cases are handled similarly.

Observe that we can use a similar strategy to determine the restrictions on irreducible representations of any torus knot group quotient  $\bar{G}_{m,n} \simeq \mathbb{Z}_m * \mathbb{Z}_n$ . Having the classification of the dimension vectors  $\alpha$  of  $\theta$ -semistable representations of  $Q_{m,n}$  we can use the local quiver settings to study these projective varieties  $M_\alpha^{ss}(Q_{m,n}, \theta)$ , in particular to determine the  $\alpha$  for which this moduli space is a projective smooth variety.

EXAMPLE 7.18. For example,  $iss_4 PSL_2(\mathbb{Z})$  has several components of dimension 3 and 2. For one of the three 3-dimensional components, the one corresponding to  $\alpha = (2, 2; 2, 1, 1)$ , the different types of semi-simples  $M_\xi$  and corresponding local quivers  $Q_\xi$  are listed in figure 3. To verify whether  $iss_n PSL_2(\mathbb{Z})$  is smooth in  $\xi$  it suffices to prove that the traces along oriented cycle for the quiver-setting  $(Q_\xi, \alpha_\xi)$  generate a polynomial algebra. For example, consider a point  $\xi \in iss_4 PSL_2(\mathbb{Z})$  of type



Then, the traces along oriented cycles in  $Q_\xi$  are generated by the following three algebraic independent polynomials

$$\begin{cases} x = ac + bd \\ y = eg + fh \\ z = (cg + dh)(ea + fb) \end{cases}$$

and hence  $iss_4 PSL_2(\mathbb{Z})$  is smooth in  $\xi$ . The other cases being easier, we see that this component of  $iss_4 PSL_2(\mathbb{Z})$  is a smooth compact manifold.

A further application of our local quiver-settings  $(Q_\xi, \alpha_\xi)$  is that one can often describe large families of irreducible  $G_{m,n}$ -representations, starting from knowing only rather trivial ones.





FIGURE 3. Local quiver settings for  $M_{\alpha}^{ss}(Q_{2,3}, \theta)$  for  $\alpha = (2, 2; 2, 1, 1)$ .

EXAMPLE 7.19. Consider the semisimple  $PSL_2(\mathbb{Z})$ -representation  $\xi$  of type



Then,  $M_{\xi}$  is determined by the following matrices

$$\left( \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{bmatrix}, \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \zeta^2 & 0 & 0 \\ 0 & 0 & \zeta & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \right)$$

The quiver-setting  $(Q_{\xi}, \alpha_{\xi})$  implies that any nearby orbit is determined by a matrix-couple

$$\left( \begin{bmatrix} 1 & b_1 & 0 & 0 \\ a_1 & -1 & d_1 & 0 \\ 0 & c_1 & 1 & f_1 \\ 0 & 0 & e_1 & -1 \end{bmatrix}, \begin{bmatrix} 1 & b_2 & 0 & 0 \\ a_2 & \zeta^2 & d_2 & 0 \\ 0 & c_2 & \zeta & f_2 \\ 0 & 0 & e_2 & 1 \end{bmatrix} \right)$$

and as there is just one arrow in each direction these entries must satisfy

$$0 = a_1 a_2 = b_1 b_2 = c_1 c_2 = d_1 d_2 = e_1 e_2 = f_1 f_2$$

As the square of the first matrix must be the identity matrix  $\mathbb{1}_4$ , we have in addition that

$$0 = a_1 b_1 = c_1 d_1 = e_1 f_1$$

Hence, we get several sheets of 3-dimensional families of representations (possibly, matrix-couples lying on different sheets give isomorphic  $PSL_2(\mathbb{Z})$ -representations, as the isomorphism holds in the étale topology and not necessarily in the Zariski topology). One of the sheets has representatives

$$\left( \begin{bmatrix} 1 & 0 & 0 & 0 \\ a & -1 & d & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & e & -1 \end{bmatrix}, \begin{bmatrix} 1 & b & 0 & 0 \\ 0 & \zeta^2 & 0 & 0 \\ 0 & c & \zeta & f \\ 0 & 0 & 0 & 1 \end{bmatrix} \right)$$

From the description of dimension vectors of semi-simple quiver representations it follows that such a representation is simple if and only if

$$ab \neq 0 \quad cd \neq 0 \quad \text{and} \quad ef \neq 0$$

Moreover, these simples are not-isomorphic unless their traces  $ab, cd$  and  $ef$  evaluate to the same numbers.

Finally, one can use the local quiver-settings  $(Q_\xi, \alpha_\xi)$  to determine the isomorphism classes of  $G_{m,n}$ -representations having a specified Jordan-Hölder sequence. For this we apply the theory on nullcones developed in the foregoing chapter.

**EXAMPLE 7.20.** In the above example, this nullcone problem is quite trivial. A representation has  $M_\xi$  as Jordan-Hölder sum if and only if all traces vanish, that is,

$$ab = cd = ef = 0$$

Under the action of the group  $GL(\alpha_\xi) = \mathbb{C}^* \times \mathbb{C}^* \times \mathbb{C}^* \times \mathbb{C}^*$ , these orbits are easily seen to be classified by the arrays

$$\begin{array}{|c|c|c|} \hline a & c & e \\ \hline b & d & f \\ \hline \end{array}$$

filled with zeroes and ones subject to the rule that no column can have two 1's, giving  $27 = 3^3$ -orbits.

### 7.5. Differential forms.

In this section we will define the complex of *noncommutative differential forms* of an arbitrary  $\mathbb{C}$ -algebra  $A$  and deduce some extra features in case  $A$  is Quillen-smooth. In the following section we will compute the *noncommutative deRham cohomology* spaces which will be of crucial importance in the final chapter.

Let us recall briefly the classical (commutative) case. When  $A$  is a commutative  $\mathbb{C}$ -algebra, the  $A$ -module of *Kähler differentials*  $\Omega_A^1$  is generated by the  $\mathbb{C}$ -linear symbols  $da$  for  $a \in A$  satisfying the relations

$$d(ab) = adb + bda \quad \forall a, b \in A$$

and the map  $A \xrightarrow{d} \Omega_A^1$  is the universal derivation. By convention we define

$$\begin{cases} \Omega_A^0 = A \\ \Omega_A^n = \wedge_A^n \Omega_A^1 \end{cases}$$

where the *exterior product* is taken over  $A$  (not over  $\mathbb{C}$ ). Observe that it is spanned by the elements  $a_0 da_1 \wedge \dots \wedge da_n$  that we usually write  $a_0 da_1 \dots da_n$ .

The *exterior differential operator*

$$\Omega_A^n \xrightarrow{d} \Omega_A^{n+1}$$

is defined by

$$d(a_0 da_1 \dots da_n) = da_0 da_1 \dots da_n$$

and gives rise to a sequence

$$A = \Omega_A^0 \xrightarrow{d} \Omega_A^1 \xrightarrow{d} \dots \xrightarrow{d} \Omega_A^n \xrightarrow{d} \Omega_A^{n+1} \xrightarrow{d} \dots$$

which is a complex (that is,  $d \circ d = 0$ ) called the *deRham complex*. The *homology groups* of this complex

$$H_{dR}^n A = \frac{\text{Ker } \Omega_A^n \xrightarrow{d} \Omega_A^{n+1}}{\text{Im } \Omega_A^{n-1} \xrightarrow{d} \Omega_A^n}$$

are called the *de Rham cohomology groups* of  $A$  (over  $\mathbb{C}$ ).

We will extend this to noncommutative  $\mathbb{C}$ -algebras. We denote by  $\text{dgalg}$  the category of *differential graded  $\mathbb{C}$ -algebras*, that is, an object  $R \in \text{dgalg}$  is a  $\mathbb{Z}$ -graded  $\mathbb{C}$ -algebra

$$R = \bigoplus_{i \in \mathbb{Z}} R_i$$

endowed with a differential  $d$  of degree one

$$\dots \xrightarrow{d} R_{i-1} \xrightarrow{d} R_i \xrightarrow{d} R_{i+1} \xrightarrow{d} \dots$$

such that  $d \circ d = 0$  and for all  $r \in R_i$  and  $s \in R$  we have

$$d(rs) = (dr)s + (-1)^i r(ds).$$

Clearly, morphisms in  $\text{dgalg}$  are  $\mathbb{C}$ -algebra morphisms  $R \xrightarrow{\phi} S$  which are graded and commute with the differentials.

To a  $\mathbb{C}$ -algebra  $A$  we will now associate the differential graded algebra  $\Omega A$  of *noncommutative differential forms*. Denote the quotient vector space  $A/\mathbb{C}$  with  $\bar{A}$  and define

$$\Omega^n A = A \otimes \underbrace{\bar{A} \otimes \dots \otimes \bar{A}}_n$$

for  $n \geq 0$  and  $\Omega^n A = 0$  for  $n < 0$ . For  $a_i \in A$  we denote the image of  $a_0 \otimes a_1 \otimes \dots \otimes a_n$  in  $\Omega^n A$  by

$$(a_0, \dots, a_n).$$

Consider the vectorspace  $\Omega A = \bigoplus_{n \in \mathbb{Z}} \Omega^n A$  and define a product on it by

$$(a_0, \dots, a_n)(a_{n+1}, \dots, a_m) = \sum_{i=0}^n (-1)^{n-i} (a_0, \dots, a_{i-1}, a_i a_{i+1}, a_{i+2}, \dots, a_m).$$

Further, define an operator  $d$  of degree one

$$\dots \xrightarrow{d} \Omega^{n-1} A \xrightarrow{d} \Omega^n A \xrightarrow{d} \Omega^{n+1} A \xrightarrow{d} \dots$$

by the rule

$$d(a_0, \dots, a_n) = (1, a_0, \dots, a_n).$$

**THEOREM 7.21.** *These formulas define the unique  $\text{dgalg}$  structure on  $\Omega A$  such that*

$$a_0 da_1 \dots da_n = (a_0, a_1, \dots, a_n).$$

PROOF. In any  $R = \oplus_i R_i \in \mathbf{dgalg}$  containing  $A$  as an even degree subalgebra we have the following identities

$$\begin{aligned} d(a_0 da_1 \dots da_n) &= da_0 da_1 \dots da_n \\ (a_0 da_1 \dots da_n)(a_{n+1} da_{n+2} \dots da_m) &= (-1)^n a_0 a_1 da_2 \dots da_m \\ &\quad + \sum_{i=1}^n (-1)^{n-i} a_0 da_1 \dots d(a_i a_{i+1}) \dots da_m \end{aligned}$$

which proves uniqueness.

To prove existence, we define  $d$  on  $\Omega A$  as above making the  $\mathbb{Z}$ -graded  $\mathbb{C}$ -vectorspace  $\Omega A$  into a complex as  $d \circ d = 0$ . Consider the *graded endomorphism ring* of the complex

$$\mathbf{End} = \oplus_{n \in \mathbb{Z}} \mathbf{End}_n = \oplus_{n \in \mathbb{Z}} \mathbf{Hom}_{\text{complex}}(\Omega^\bullet A, \Omega^{\bullet+n} A).$$

With the composition as multiplication,  $\mathbf{End}$  is a  $\mathbb{Z}$ -graded  $\mathbb{C}$ -algebra and we make it into an object in  $\mathbf{dgalg}$  by defining a differential

$$\dots \xrightarrow{D} \mathbf{End}_{n-1} \xrightarrow{D} \mathbf{End}_n \xrightarrow{D} \mathbf{End}_{n+1} \xrightarrow{D} \dots$$

by the formula on any homogeneous  $\phi$

$$D\phi = d \circ \phi - (-1)^{\deg \phi} \phi \circ d.$$

Now define the morphism  $A \xrightarrow{l} \mathbf{End}_0$  which assigns to  $a \in A$  the left multiplication operator

$$la(a_0, \dots, a_n) = (aa_0, \dots, a_n)$$

and extend it to a map

$$\Omega A \xrightarrow{l_*} \mathbf{End} \quad \text{by} \quad l_*(a_0, \dots, a_n) = la_0 \circ D la_1 \circ \dots \circ D la_n.$$

Applying the general formulae given at the beginning of the proof to the subalgebra  $l(A) \hookrightarrow \mathbf{End}$  we see that the image of  $l_*$  is a differential graded subalgebra of  $\mathbf{End}$  and is the differential graded subalgebra generated by  $l(A)$ .

Define an evaluation map  $\mathbf{End} \xrightarrow{ev} \Omega A$  by  $ev(\phi) = \phi(1)$ . Because

$$\begin{aligned} D la_i(1, a_{i+1}, \dots, a_n) &= d(a_i, a_{i-1}, \dots, a_n) - la_i d(1, a_{i+1}, \dots, a_n) \\ &= (1, a_i, \dots, a_n) \end{aligned}$$

we have that

$$ev(la_0 \circ D la_1 \circ \dots \circ D la_n) = (a_0, \dots, a_n)$$

showing that  $ev$  is a left inverse for  $l_*$  whence  $l_*$  is injective.

Hence we can use the isomorphism  $\Omega A \simeq \text{Im}(l_*)$  to transport the  $\mathbf{dgalg}$  structure to  $\Omega A$  finishing the proof.  $\square$

EXAMPLE 7.22. Noncommutative differential forms of  $\mathbb{C} \times \mathbb{C}$ .

Let  $A = \mathbb{C} \times \mathbb{C}$  and  $e$  and  $f$  the idempotents corresponding to the two factors. The quotient space  $\bar{A} = A/\mathbb{C}1$  can be identified with  $\mathbb{C}\bar{e}$  and therefore

$$\Omega^n \mathbb{C} \times \mathbb{C} = (\mathbb{C} \times \mathbb{C}) \otimes \mathbb{C}\bar{e}^{\otimes n} = (\mathbb{C} \times \mathbb{C}) de^n.$$

The differential  $d$  is defined by the formula

$$d((\alpha e + \beta f) de^n) = (\alpha - \beta) de^{n+1}$$

and the product of  $\Omega \mathbb{C} \times \mathbb{C}$  is defined by the rule

$$(\alpha e + \beta f) de^n (\gamma e + \delta f) de^m = \begin{cases} (\alpha \gamma e + \beta \delta f) de^{n+m} & \text{when } n \text{ is even} \\ (\alpha \delta e + \beta \gamma f) de^{n+m} & \text{when } n \text{ is odd} \end{cases}$$

We will relate the algebra structure of  $\Omega A$  to that of  $A$ . The trick is to define *another* multiplication on  $\Omega A$  making it only into a *filtered* algebra. We then prove that this filtered algebra is isomorphic to the  $I$ -adic filtration of an algebra constructed from  $A$  and we recover the **dgalg** multiplication on  $\Omega A$  by taking the associated graded algebra.

We introduce the universal algebra  $\mathbb{L}_A$  with respect to *based linear maps* from  $A$  to  $\mathbb{C}$ -algebras. A based linear map is a  $\mathbb{C}$ -linear map

$$A \xrightarrow{\rho} R$$

where  $R$  is a  $\mathbb{C}$ -algebra and  $\rho(1) = 1$ . The *curvature* of  $\rho$  is then defined to be the bilinear map  $A \times A \xrightarrow{\omega} R$  defined by

$$\omega(a, a') = \rho(aa') - \rho(a)\rho(a')$$

that is, it is a measure for the failure of  $\rho$  to be an algebra map. Observe that  $\omega$  vanishes if either  $a$  or  $a'$  is 1 so it can be viewed as a linear map

$$\overline{A} \otimes \overline{A} \xrightarrow{\omega} R.$$

Let  $T(A) = \bigoplus_{n \geq 0} A^{\otimes n}$  be the *tensor algebra* of the vectorspace  $A$  and define

$$\mathbb{L}_A = \frac{T(A)}{T(A)(1 - 1_A)T(A)}$$

where  $1_A$  is the identity of  $A$  consider as a 1-tensor in  $T(A)$ , then we have a based linear map

$$A \xrightarrow{\rho^{un}} \mathbb{L}_A \quad a \mapsto \bar{a}$$

where  $\bar{a}$  is the image in  $\mathbb{L}_A$  of the 1-tensor  $a$  in  $T(A)$ . The map  $\rho^{un}$  is universal for based linear maps  $A \xrightarrow{\rho} R$ , that is, there is a unique *algebra morphism*  $\mathbb{L}_A \xrightarrow{\phi_\rho} R$  making the diagram commute

$$\begin{array}{ccc} & & \mathbb{L}_A \\ & \nearrow \rho^{un} & \downarrow \exists \phi_\rho \\ A & \xrightarrow{\rho} & R \end{array}$$

In particular, there is a canonical algebra map  $\mathbb{L}_A \xrightarrow{\phi_{id}} A$  corresponding to the identity map on  $A$ . We define

$$\mathbb{I}_A = \text{Ker } \phi_{id} \triangleleft \mathbb{L}_A$$

and equip  $\mathbb{L}_A$  with the  $\mathbb{I}_A$ -adic filtration.

For an arbitrary  $R \in \mathbf{dgalg}$  we define the *Fedosov product* on  $R$  to be the one induced by defining on homogeneous  $r, s \in R$  the product

$$r.s = rs - (-1)^{\text{deg } r} drds$$

One easily checks that the Fedosov product is associative. Observe that if we decompose  $R = R^{ev} \oplus R^{odd}$  into its homogeneous components of even (resp. odd) degree, then this new multiplication is compatible with this decomposition and makes  $R$  into a  $\mathbb{Z}/2\mathbb{Z}$ -graded algebra.

We will now investigate the Fedosov product on  $\Omega A$ . Let  $\omega^{un}$  be the curvature of the universal based linear map  $A \xrightarrow{\rho^{un}} \mathbb{L}_A$ .

THEOREM 7.23. *There is an isomorphism of algebras*

$$\mathbb{L}_A \simeq (\Omega^{ev} A, \cdot)$$

between  $\mathbb{L}_A$  and the even forms  $\Omega^{ev} A$  equipped with the Fedosov product given by

$$\rho^{un}(a_0)\omega^{un}(a_1, a_2) \dots \omega^{un}(a_{2n-1}, a_{2n}) \longrightarrow a_0 da_1 \dots da_{2n}$$

Under this isomorphism we have the correspondence

$$\mathbb{I}_A^n \simeq \bigoplus_{k \geq n} \Omega^{2k} A$$

The associated graded algebra gives an isomorphism

$$\mathrm{gr}_{\mathbb{I}_A} \mathbb{L}_A = \bigoplus_{\mathbb{I}_A^{n+1}}^{\mathbb{I}_A^n} \simeq \Omega^{ev} A$$

with even forms equipped with the **dgalg** structure.

PROOF. Consider the based linear map  $A \xrightarrow{\rho} \Omega^{ev} A$  given by inclusion, then its curvature is given by

$$\omega(a, a') = aa' - a.a' = dada'.$$

By the universal property of  $\mathbb{L}_A$  there is an algebra morphism

$$\mathbb{L}_A \xrightarrow{\phi} (\Omega^{ev} A, \cdot)$$

such that  $\phi(\rho^{un}(a)) = a$  and  $\phi(\omega^{un}(a, a')) = dada'$ . Observe that the Fedosov product coincides with the usual **dgalg** product when one of the terms is *closed*, that is  $d r = 0$ . Therefore, we have

$$\phi(\rho^{un}(a_0)\omega^{un}(a_1, a_2) \dots \omega^{un}(a_{2n-1}, a_{2n})) = a_0 da_1 \dots da_{2n}$$

On the other hand, as  $\Omega^{2n} A = A \otimes \overline{A}^{\otimes 2n}$  we have a well defined map  $\Omega^{ev} A \xrightarrow{\psi} \mathbb{L}_A$  given by

$$\psi(a_0 da_1 \dots da_{2n}) = \rho^{un}(a_0)\omega^{un}(a_1, a_2) \dots \rho^{un}(a_{2n-1}, a_{2n})$$

and it remains to prove that this map is surjective. The image of  $\psi$  is closed under left multiplication as it is closed under left multiplication by elements  $\rho^{un}(a)$  (and they generate  $\mathbb{L}_A$ ) as

$$\begin{aligned} & \rho^{un}(a) \cdot \rho^{un}(a_0)\omega^{un}(a_1, a_2) \dots \omega^{un}(a_{2n-1}, a_{2n}) \\ = & \rho^{un}(aa_0)\omega^{un}(a_1, a_2) \dots \omega^{un}(a_{2n-1}, a_{2n}) - \omega^{un}(a, a_0)\omega^{un}(a_1, a_2) \dots \omega^{un}(a_{2n-1}, a_{2n}) \end{aligned}$$

Because the image contains 1 this proves the claim and the isomorphism.

Identify via this isomorphism  $\mathbb{L}_A$  with  $\Omega^{ev} A$ . Because  $dada' \in \mathbb{I}_A$  we have  $\Omega^{2k} A \hookrightarrow \mathbb{I}_A^n$  for all  $k \geq n$ . Thus,  $F_n = \bigoplus_{k \geq n} \Omega^{2k} A \hookrightarrow \mathbb{I}_A$ . Conversely,  $\mathbb{I}_A = F_1$  and hence

$$\mathbb{I}_A^n = (F_1)^n \hookrightarrow F_n$$

by the definition of the Fedosov product. Therefore,  $\mathbb{I}_A^n = F_n$  and the claim over the associated graded follows.  $\square$

EXAMPLE 7.24. Even differential forms of  $\mathbb{C} \times \mathbb{C}$ .

As before, let  $e$  and  $f$  be the idempotents of  $A = \mathbb{C} \times \mathbb{C}$  corresponding to the two components. By definition,

$$\mathbb{L}_{\mathbb{C} \times \mathbb{C}} = \frac{T(\mathbb{C}e + \mathbb{C}f)}{(1 - e - f)} = \frac{\mathbb{C}\langle E, F \rangle}{(1 - E - F)} \simeq \mathbb{C}\langle E \rangle$$



The universal based linear map is given by

$$\mathbb{C} \times \mathbb{C} \xrightarrow{\rho^{un}} \mathbb{C}[E] \quad \begin{cases} e & \mapsto E \\ f & \mapsto 1 - E \end{cases}$$

and the curvature on  $\bar{A} = \mathbb{C}\bar{e}$  is given by

$$\omega^{un}(e, e) = E - E^2$$

Therefore the isomorphism between  $\Omega^{ev} A$  and  $\mathbb{L}_A = \mathbb{C}[E]$  is given by

$$(\alpha e + \beta f)de^{2n} \xrightarrow{\psi} (\alpha E + \beta(1 - E))(E - E^2)^n.$$

The Fedosov product on  $\Omega^{ev} A$  is given by the formula (using the multiplication formulas we found above)

$$(\alpha e + \beta f)de^{2n} \cdot (\gamma e + \delta f)de^{2m} = (\alpha\gamma e + \beta\delta f)de^{2n+2m} - (\alpha - \beta)(\gamma - \delta)de^{2n+2m+2}$$

In order to check that  $\psi$  is indeed an algebra morphism we need to verify that in  $\mathbb{C}[E]$  we have the equality

$$\begin{aligned} & (\alpha E + \beta(1 - E))(E - E^2)^n (\gamma E + \delta(1 - E))(E - E^2)^m \\ = & (\alpha\gamma E + \beta\delta(1 - E))(E - E^2)^{n+m} - (\alpha - \beta)(\gamma - \delta)(E - E^2)^{n+m+1} \end{aligned}$$

which is indeed the case.

Further,  $\mathbb{L}_A = \mathbb{C}[E](E - E^2)$  and indeed  $\frac{\mathbb{C}[E]}{(E - E^2)} \simeq \mathbb{C} \times \mathbb{C}$ . Finally, under the identification  $\psi$  we obtain the usual multiplication of noncommutative differential forms from

$$\Omega^{2n} A \times \Omega^{2m} A = \frac{(E - E^2)^n}{(E - E^2)^{n+1}} \times \frac{(E - E^2)^m}{(E - E^2)^{m+1}} \longrightarrow \frac{(E - E^2)^{n+m}}{(E - E^2)^{n+m+1}} = \Omega^{2n+2m} A.$$


We now turn to *all* noncommutative differential forms  $\Omega A$ . Observe that this algebra has an involution  $\sigma$  which is the identity on even forms and is minus the identity on odd forms.  $\sigma$  is an algebra automorphism both for the usual  $\mathbf{dgalg}$ -algebra structure as for the Fedosov product. Algebras with an involution are called *super-algebras*.

We want to construct an algebra universal for algebra morphisms from  $A$  to a super-algebra. Consider the *free product*  $A * A$  which is defined as follows. Let  $\mathcal{B}_1$  be a vectorspace basis for  $A - \mathbb{C} \cdot 1$  and  $\mathcal{B}_2$  a duplicate of it. As a  $\mathbb{C}$ -vectorspace  $A * A$  has a basis consisting of words

$$w = a_1 b_1 a_2 b_2 \dots a_k b_k \quad \text{or} \quad w = a_1 b_1 a_2 b_2 \dots a_k$$


for some  $k$  where the  $a_i$ 's all belong to  $\mathcal{B}_1$  or all to  $\mathcal{B}_2$  and the  $b_j$ 's all belong to the other base set. On this vectorspace one defines a  $\mathbb{C}$ -algebra structure in the obvious way, that is by concatenating words and if necessary (if the end term of the first word lies in the same base-set as the beginning term of the second) use the multiplication table in  $A$  to reduce to a linear combination of allowed words.

The algebra  $A * A$  is universal with respect to *pairs of algebra maps*  $A \begin{matrix} \xrightarrow{f} \\ \xrightarrow{g} \end{matrix} R$  from  $A$  to  $R$ . That is, there is a unique algebra map  $\gamma$



making the diagram commute. Here,  $i_1$  is the inclusion of  $A$  in  $A * A$  using only syllables in  $\mathcal{B}_1$  and  $i_2$  is defined similarly. The construction of  $\gamma$  clearly is induced by sending  $a \in \mathcal{B}_1$  to  $f(a)$  and  $b \in \mathcal{B}_2$  to  $g(b)$ .

Further, interchanging the bases  $\mathcal{B}_1 \xrightarrow{\tau} \mathcal{B}_2$  equips  $A * A$  with an involution, or if you prefer, makes  $A * A$  a super-algebra. Now, let  $S$  be a super-algebra with involution  $\sigma_S$  and let  $A \xrightarrow{f} S$  be an algebra morphism, then there is a unique morphism of super-algebras  $\psi$  making the diagram commute



$\psi$  is the universal map corresponding to the pair of algebra maps  $A \begin{matrix} \xrightarrow{f} \\ \xrightarrow{\sigma_S \circ f} \end{matrix} S$ .

For any  $a \in A$  we define the elements in  $A * A$  :

$$\begin{cases} p(a) &= \frac{1}{2}(i_1(a) + i_2(a)) \\ q(a) &= \frac{1}{2}(i_1(a) - i_2(a)) \end{cases}$$

and we define  $\mathbb{Q}_A \triangleleft A * A$  to be the ideal of  $A * A$  generated by the elements  $q(a)$  for  $a \in A$ , then clearly

$$A \simeq \frac{A * A}{\mathbb{Q}_A}$$

We now have an analog of the previous theorem for all differential forms.

**THEOREM 7.25.** *There is an isomorphism of super-algebras*

$$A * A \simeq (\Omega A, \cdot)$$

between  $A * A$  and the noncommutative differential forms  $\Omega A$  equipped with the Fedosov product given by

$$p(a_0)q(a_1) \dots q(a_n) \longrightarrow a_0 da_1 \dots da_n$$

Under this isomorphism we have the correspondence

$$\mathbb{Q}_A^n \simeq \oplus_{k \geq n} \Omega^k A$$

and the associated graded algebra is isomorphic to  $\Omega A$  with the usual dgalg structure.

PROOF. We have an algebra map  $A \xrightarrow{u} \Omega A$  equipped with the Fedosov product given by  $a \mapsto a + da$  because

$$\begin{aligned} (a + da).(a' + da') &= aa' - dada' + ada' + daa' + dada' \\ &= aa' + d(aa') \end{aligned}$$

By the universal property of  $A * A$  there is a super-algebra morphism

$$A * A \xrightarrow{\psi} \Omega A \quad \psi(p(a)) = a \quad \text{and} \quad \psi(q(a)) = da$$

But then using that the Fedosov product coincides with the usual product when one of the forms is closed we have

$$\psi(p(a_0)q(a_1) \dots q(a_n)) = a_0 da_1 \dots da_n$$

Conversely, we have a section to  $\psi$  defined by

$$\Omega A \xrightarrow{\phi} A * A \quad a_0 da_1 \dots da_n \mapsto p(a_0)q(a_1) \dots q(a_n)$$

and we only have to prove that  $\phi$  is surjective. The image  $\text{Im } \phi$  is closed under left multiplication by  $p(a)$  and  $q(a)$  as  $p(1) = 1$  and


$$\begin{cases} p(a)p(a_0)q(a_1) \dots q(a_n) = p(aa_0)q(a_1) \dots q(a_n) - q(a)q(a_0)q(a_1) \dots q(a_n) \\ q(a)p(a_0)q(a_1) \dots q(a_n) = q(aa_0)q(a_1) \dots q(a_n) - p(a)q(a_0)q(a_1) \dots q(a_n) \end{cases}$$

Because the elements  $p(a)$  and  $q(a)$  generate  $A * A$ , the image  $\text{Im } \phi$  is a left ideal containing 1, whence  $\psi$  is surjective.

The claims about the ideals  $\mathbb{Q}_A^n$  and about the associated graded algebra follow as in the proof for even forms.  $\square$

EXAMPLE 7.26. Noncommutative differential forms of  $\mathbb{C}\langle x, y \rangle$ .

The noncommutative free algebra in two variables  $\mathbb{C}\langle x, y \rangle$  is the path algebra of the quiver



Clearly we have  $\mathbb{C}\langle x, y \rangle * \mathbb{C}\langle x, y \rangle = \mathbb{C}\langle x_1, y_1, x_2, y_2 \rangle$  and the maps

$$\begin{cases} p(x) = \frac{1}{2}(x_1 + x_2) & q(x) = \frac{1}{2}(x_1 - x_2) \\ p(y) = \frac{1}{2}(y_1 + y_2) & q(y) = \frac{1}{2}(y_1 - y_2) \end{cases}$$

It is easy to compute the maps  $p$  and  $q$  on any monomial in  $x$  and  $y$  using the formulae holding in any  $A * A$

$$\begin{cases} p(aa') = p(a)p(a') + q(a)q(a') \\ q(aa') = p(a)q(a') + q(a)p(a') \end{cases}$$

Further note that it follows from this that  $\mathbb{Q}_{\mathbb{C}\langle x, y \rangle} = (x_1 - x_2, y_1 - y_2)$  and we have all the required tools to calculate (in principle) with  $\Omega \mathbb{C}\langle x, y \rangle$ .

EXAMPLE 7.27. Noncommutative differential forms of  $\mathbb{C} \times \mathbb{C}$ .

The infinite dihedral group  $D_\infty$  is the group with presentation

$$D_\infty = \langle a, b \mid a^2 = 1 = b^2 \rangle$$

that is, an arbitrary element in  $D_\infty$  is a word of the form

$$a^i babab \dots abab^j$$

where  $i, j = 0$  or  $1$ . Multiplication is given by concatenation of words, using the relations  $a^2 = 1 = b^2$  when necessary.

The *group algebra*  $\mathbb{C}[D_\infty]$  is the vectorspace with basis  $D_\infty$  and with multiplication induced by the groupmultiplication in  $D_\infty$ . We now claim that

$$(\mathbb{C} \times \mathbb{C}) * (\mathbb{C} \times \mathbb{C}) \simeq \mathbb{C}[D_\infty]$$

Indeed,  $\mathbb{C} \times \mathbb{C} \simeq \mathbb{C}[\mathbb{Z}_2]$  the group algebra of the cyclic group of order two, that is  $\mathbb{C}[\mathbb{Z}_2] = \mathbb{C}[x]/(x^2 - 1)$ , the isomorphism being given by

$$e \longrightarrow \frac{1}{2}(1 + x) \quad f \longrightarrow \frac{1}{2}(1 - x)$$

One also has the obvious notion of a free product in the category of groups and from the definition it is clear that

$$\mathbb{Z}_2 * \mathbb{Z}_2 \simeq D_\infty$$

and therefore also on the level of group algebras

$$\mathbb{C}[\mathbb{Z}_2] * \mathbb{C}[\mathbb{Z}_2] \simeq \mathbb{C}[D_\infty]$$

The relevant maps  $\mathbb{C} \times \mathbb{C} \begin{matrix} \xrightarrow{p} \\ \xrightarrow{q} \end{matrix} \mathbb{C}[D_\infty]$  are given by

$$\begin{cases} p(e) = \frac{1}{2} + \frac{1}{4}(a + b) & q(e) = \frac{1}{4}(a - b) \\ p(f) = \frac{1}{2} - \frac{1}{4}(a + b) & q(f) = -\frac{1}{4}(a - b) \end{cases}$$

and so  $\mathbb{Q}_{\mathbb{C} \times \mathbb{C}} = (a - b) \triangleleft \mathbb{C}[D_\infty]$ . Again, this information allows us to calculate with  $\Omega \mathbb{C} \times \mathbb{C}$  by referring all computations to the more familiar group algebra  $\mathbb{C}[D_\infty]$ .

The above definitions and results are valid for every  $\mathbb{C}$ -algebra  $A$ . We will indicate a few extra properties provided the algebra  $A$  is Quillen-smooth.

We have the universal lifting algebra  $\mathbb{L}_A$  for based *linear* maps from  $A$  to  $\mathbb{C}$ -algebras and the ideal  $\mathbb{I}_A$  such that

$$A \xleftarrow[\simeq]{\overline{\phi_{id}}} \frac{\mathbb{L}_A}{\mathbb{I}_A}.$$

The  $\mathbb{I}_A$ -adic completion of  $\mathbb{L}_A$  is by definition the inverse limit


$$\hat{\mathbb{L}}_A = \varprojlim_n \frac{\mathbb{L}_A}{\mathbb{I}_A}$$

Assume that  $A$  is formally smooth, then for every  $k$  we have an algebra map lifting  $\overline{\phi_{id}}^{-1}$

$$\begin{array}{ccc} & & \frac{\mathbb{L}_A}{\mathbb{I}_A^k} \\ & \nearrow \vartheta^* & \downarrow \\ A & \xrightarrow{\overline{\phi_{id}}^{-1}} & \frac{\mathbb{L}_A}{\mathbb{I}_A} \end{array}$$

These compatible lifts define an algebra lift  $A \xrightarrow{I^{un}} \hat{\mathbb{L}}_A$ . This map can be used to construct algebra lifts modulo nilpotent ideals in a systematic way. Assume  $I \triangleleft R$  is such that  $I^k = 0$  and there is an algebra map  $A \xrightarrow{\mu} \frac{R}{I}$ . We can lift  $\mu$  to  $R$  as

a based linear map, say  $\rho$ . Now we have the following situation



Here,  $\phi_\rho$  is the algebra map coming from the universal lifting property of  $\mathbb{L}_A$  and  $\hat{\phi}_\rho$  is its extension to the completion. But then,  $\tilde{\mu} = \hat{\phi}_\rho \circ l^{un}$  is an algebra lift of  $\mu$ . That is,

PROPOSITION 7.28. *A is formally smooth if and only if there is an algebra section  $A \rightarrow \hat{\mathbb{L}}_A$  to the projection  $\hat{\mathbb{L}}_A \rightarrow A$  defined by mapping out  $\mathbb{I}_A$ .*

We will give an explicit construction of the embedding  $A \xrightarrow{l^{un}} \hat{\mathbb{L}}_A$ . By formal smoothness we have an algebra lift

$$\begin{array}{ccc}
 & A \oplus \Omega^2 A = & \frac{\mathbb{L}_A}{\mathbb{I}_A^2} \\
 \nearrow \nu & & \downarrow \\
 A \xrightarrow{id} A & = & \frac{\mathbb{L}_A}{\mathbb{I}_A}
 \end{array}$$

which is of the form  $l_2(a) = a - \phi(a)$  for a linear map  $A \xrightarrow{\phi} \Omega^2 A$ . As  $\mathbb{L}_A$  is freely generated by the  $a \in A - \mathbb{C}1$ , we can define a derivation on  $\mathbb{L}_A$  defined by

$$\mathbb{L}_A \xrightarrow{D} \mathbb{L}_A \quad D(a) = \phi(a) \quad \forall a \in A.$$

This derivation is called the *Yang-Mills derivation* of  $A$ .

Clearly  $D(\mathbb{L}_A) \hookrightarrow \mathbb{I}_A$  and we have

$$\begin{aligned}
 D(dada') &= D(aa' - a.a') \\
 &= D(aa') - D(a).a' - a.D(a') \\
 &= \phi(aa') - \phi(a).a' - a.\phi(a') \\
 &\equiv aa' - a.a' \pmod{\mathbb{I}_A^2} \\
 &\equiv dada' \pmod{\mathbb{I}_A^2}
 \end{aligned}$$

the next to last equality coming from the fact that  $l_2$  is an algebra map. Hence,  $D = id$  on  $\frac{\mathbb{L}_A}{\mathbb{I}_A^2} = \Omega^2 A$ .

Further,  $D(\mathbb{I}_A^n) \hookrightarrow \mathbb{I}_A^n$  and so  $D$  induces a derivation on the associated graded  $gr_{\mathbb{I}_A} \mathbb{L}_A$ . As this derivation is zero on  $A = \frac{\mathbb{L}_A}{\mathbb{I}_A}$  and one on  $\frac{\mathbb{I}_A^n}{\mathbb{I}_A^{n+1}}$  it is  $n$  on  $\frac{\mathbb{I}_A^n}{\mathbb{I}_A^{n+1}}$ . But

then we have by induction

$$(D - n)\dots(D - 1)D(\mathbb{L}_A) \hookrightarrow \mathbb{I}_A^{n+1}$$

Therefore,  $\frac{\mathbb{L}_A}{\mathbb{I}_A^{n+1}}$  decomposes into eigenspaces of  $D$  corresponding to the eigenvalues  $0, 1, \dots, n$  and because  $D$  is a derivation this decomposition defines a grading compatible with the product.

Hence, we obtain an isomorphism of  $\frac{\mathbb{L}_A}{\mathbb{I}_A^{n+1}}$  with its associated graded algebra by lifting  $\frac{\mathbb{I}_A^k}{\mathbb{I}_A^{k+1}}$  to the eigenspace of  $D$  on  $\frac{\mathbb{I}_A^k}{\mathbb{I}_A^{k+1}}$  corresponding to the eigenvalue  $k$ .

Taking the inverse limit as  $n \rightarrow \infty$  we obtain an algebra isomorphism of  $\hat{\mathbb{L}}_A$  with the completion of its associated graded algebra, that is,

$$\Omega^{\text{ev}} A = \prod_n \Omega^{2n} A \simeq \hat{\mathbb{L}}_A$$

In particular, the kernel of  $D$  is a subalgebra of  $\hat{\mathbb{L}}_A$  mapped isomorphically onto  $A$  by the canonical surjection  $\hat{\mathbb{L}}_A \rightarrow A$ . Hence, this subalgebra gives the desired universal lift  $A \xrightarrow{l^{un}} \hat{\mathbb{L}}_A$ .

We can even give an explicit formula for  $l^{un}$ . Let  $L$  be the degree two operator on  $\Omega^{\text{ev}} A$  defined by

$$L(a_0 da_1 \dots da_{2n}) = \phi(a_0) da_1 \dots da_{2n} + \sum_{j=1}^{2n} a_0 da_1 \dots da_{j-1} d\phi(a_j) da_{j+1} \dots da_{2n}$$

and let  $H$  denote the degree zero operator on even forms which is multiplication by  $n$  on  $\Omega^{2n} A$ . Then, we have the relations

$$[H, L] = L \quad \text{and} \quad D = H + L$$

whence we have on  $\hat{\Omega}^{\text{ev}} A$  that

$$e^{-L} H e^L = H + e^{-L} [H, e^L] = H + \int_0^1 e^{-tL} [H, L] e^{tL} dt = D$$

Therefore, the universal lift for all  $a \in A$  is given by

$$l^{un}(a) = e^{-L} a = a - \phi(a) + \frac{1}{2} L\phi(a) - \dots$$

**EXAMPLE 7.29.** The universal lift for  $\mathbb{C} \times \mathbb{C}$ .

Recall the correspondence between  $\Omega^{\text{ev}} \mathbb{C} \times \mathbb{C}$  and  $\mathbb{L}_{\mathbb{C} \times \mathbb{C}} = \mathbb{C}[E]$  given by

$$(\alpha e + \beta f) d e^{2n} \longrightarrow (\alpha E + \beta(1 - E))(E - E^2)^n$$

Lifting  $e$  to  $\frac{1}{2}$  we have to compute

$$(2 - E)^2 E^2 = E + (2E - 1)(E - E^2) + (E - E^2)^2$$

whence  $\phi(e) = (1 - 2E)(E - E^2)$  and as  $f = 1 - e$  we have  $\phi(f) = (2E - 1)(E - E^2)$ . The Yang-Mills derivation  $D$  on  $\mathbb{C}[E]$  is hence the one determined by

$$\mathbb{C}[E] \xrightarrow{D} \mathbb{C}[E] \quad D(E) = (1 - 2E)(E - E^2).$$

To determine the universal lift of  $e$  we have to compute

$$l^{un}(e) = e - Le + \frac{1}{2} L^2 e - \frac{1}{6} L^3 e + \dots$$

and we have

$$\begin{aligned} L(e) &= \phi(e) = (f - e)de^2 \\ L^2(e) &= L(f - e)de^2 = -6(f - e)de^4 \\ L^3(e) &= -6L(f - e)de^4 = 60(f - e)de^6 \\ L^4(e) &= \dots \end{aligned}$$

and therefore

$$l^{un}(e) = E + (2E - 1)(E - E^2) + 3(2E - 1)(E - E^2)^2 + 10(2E - 1)(E - E^2)^3 + \dots$$

Another characteristic feature of formally smooth algebras is the existence of *connections* on  $\Omega^1 A$ . If  $E$  is an  $A$ -bimodule, then a connection on  $E$  consists of two operators

- A *right connection* :  $E \xrightarrow{\nabla_r} E \otimes_A \Omega^1 A$  satisfying
 
$$\nabla_r(aea') = a(\nabla_r e)a' + aeda',$$
- A *left connection* :  $E \xrightarrow{\nabla_l} \Omega^1 A \otimes_A E$  satisfying
 
$$\nabla_l(aea') = a(\nabla_l e)a' + daea'$$

Given a right connection  $\nabla_r$  there is a bimodule splitting  $s_r$  of the right multiplication map  $m_r$

$$E \otimes_A A \begin{array}{c} \xrightarrow{m_r} \\ \xleftarrow{s_r} \end{array} E$$

given by the formula

$$s_r(e) = e \otimes 1 - j(\nabla_r e) \quad \text{where} \quad j(e \otimes da) = ea \otimes 1 - e \otimes a$$

Similarly, a left connection gives a bimodule splitting  $s_l$  to the left multiplication map. Consequently, if a connection exists on  $E$ , then  $E$  must be a *projective* bimodule.

Consider the special bimodule of noncommutative 1-forms  $\Omega^1 A$ , then as  $\Omega^1 A \otimes_A \Omega^1 A = \Omega^2 A$  a connection on  $\Omega^1 A$  is the datum of three maps

$$\Omega^1 A \begin{array}{c} \xrightarrow{\nabla_l} \\ \xrightarrow{d} \\ \xrightarrow{\nabla_r} \end{array} \Omega^2 A$$

satisfying the following properties

$$\begin{aligned} \nabla_l(aea') &= a\nabla_l(e)a' & + (da)ea' \\ d(aea') &= a(de)a' & + (da)ea' & - ae(da') \\ \nabla_r(aea') &= a\nabla_r(e)a' & & + ae(da') \end{aligned}$$

Hence, if  $\nabla_r$  is a right connection then  $d + \nabla_r$  is a left connection and if  $\nabla_l$  is a left connection then  $\nabla_l - d$  is a right connection. Therefore, on-sided connections exist on  $\Omega^1 A$  if and only if connections exist and hence if and only if  $\Omega^1 A$  is a projective bimodule.

But then we have an  $A$ -bimodule splitting of the exact sequence

$$0 \longrightarrow \Omega^2 A \xrightarrow{j} \Omega^1 A \otimes A \xrightarrow{m} \Omega^1 A \longrightarrow 0$$

where  $j(\omega da) = \omega a \otimes 1 - \omega \otimes a$  and  $m(\omega \otimes a) = \omega a$ .

**PROPOSITION 7.30.** *A connection exists on  $\Omega^1 A$  if and only if  $A$  is formally smooth.*

PROOF. A bimodule splitting of the above map is determined by a retraction bimodule map  $p$  for  $j$ . As  $\Omega^1 A \otimes A \simeq A \otimes \bar{A} \otimes A$ , a bimodule map  $p$

$$\Omega^1 A \otimes A \xrightarrow{p} \Omega^2 A$$

is equivalent to a map  $\bar{A} \xrightarrow{\phi} \Omega^2 A$  via  $p(a_0 da_1 \otimes a_2) = a_0 \phi(a_1 a_2)$ . But then we have

$$\begin{aligned} pj(da_1 da_2) &= p((da_1)a_2 \otimes 1 - da_1 \otimes da_2) \\ &= p(d(a_1 a_2) \otimes 1 - a_1(da_2) \otimes 1 - da_1 \otimes a_2) \\ &= \phi(a_1 a_2) - a_1 \phi(a_2) - \phi(a_1) a_2 \end{aligned}$$

and splitting of the map means  $pj = id$  that is that  $\phi$  satisfies

$$\phi(aa') = a\phi(a') + \phi(a)a' + dada'$$

which is equivalent to an algebra lift

$$A \xrightarrow{\phi^*} \frac{\mathbb{L}_A}{\mathbb{I}_A} = A \oplus \Omega^2 A$$

Now, assume we have an algebra morphism

$$A \xrightarrow{f} \frac{R}{I} \quad \text{with} \quad I^2 = 0$$

and lift  $f$  to a based linear map  $A \xrightarrow{\rho} R$ . By the universal property of  $\mathbb{L}_A$  we have an algebra lift

$$\mathbb{L}_A \xrightarrow{\rho^*} R$$

living over  $f$ . Therefore  $\rho^*(\mathbb{I}_A) \subset I$  and therefore  $\rho^*$  is zero on  $\mathbb{I}_A^2$  giving an algebra morphism

$$\frac{\mathbb{L}_A}{\mathbb{I}_A^2} \xrightarrow{f^*} R$$

living over  $f$ . But then the existence of an algebra map  $\phi^*$  as above gives a desired lifting  $f^* \circ \phi^*$  of  $f$ , finishing the proof.  $\square$

For a map  $\bar{A} \xrightarrow{\phi} \Omega^2 A$  as above, a connection is given by the formulae

$$\nabla_r(ada') = a\phi(a') \quad \text{and} \quad \nabla_r(ada') = a\phi(a') + dada'$$

EXAMPLE 7.31. Connection on  $\mathbb{C}\langle x, y \rangle$ .

Clearly we have  $\Omega^1 \mathbb{C}\langle x, y \rangle = \mathbb{C}\langle x, y \rangle \otimes \mathbb{C}x + \mathbb{C}y \otimes \mathbb{C}\langle x, y \rangle$  which is the free bimodule generated by  $dx$  and  $dy$ . There is a canonical connection with

$$\begin{cases} \phi(x) = 0 & \text{and} & \nabla_l(dx) = \nabla_r(dx) = 0 \\ \phi(y) = 0 & \text{and} & \nabla_l(dy) = \nabla_r(dy) = 0 \end{cases}$$

The image of  $\phi$  on any word  $z_1 \dots z_n$  with  $z_i = x$  or  $y$  is given by the formula

$$\begin{aligned} \phi(z_1 \dots z_n) &= \nabla_r d(z_1 \dots z_n) \\ &= \nabla_r \left( \sum_{i=1}^n z_1 \dots z_{i-1} (dz_i) z_{i+1} \dots z_n \right) \\ &= \sum_{i=1}^{n-1} z_1 \dots z_{i-1} (dz_i) d(z_{i+1} \dots z_n) \end{aligned}$$



EXAMPLE 7.32. Connection on  $\mathbb{C} \times \mathbb{C}$ .

We have calculated above that the lifting map  $\phi$  is determined by

$$\phi(e) = (1 - 2E)(E - E^2) = (f - e)de^2$$

Therefore the corresponding left and right connections are given by

$$\begin{cases} \nabla_r((\alpha e + \beta f)de) = (\beta f - \alpha e)de^2 \\ \nabla_l((\alpha e + \beta f)de) = (\alpha f - \beta e)de^2 \end{cases}$$

### 7.6. deRham cohomology.

In this section we will compute various sorts of *noncommutative deRham cohomology*. We have for an arbitrary  $\mathbb{C}$ -algebra  $A$  the complex of *noncommutative differential forms*

$$A = \Omega^0 A \xrightarrow{d} \Omega^1 A \xrightarrow{d} \dots \xrightarrow{d} \Omega^n A \xrightarrow{d} \Omega^{n+1} A \xrightarrow{d} \dots$$

A first attempt to define noncommutative de Rham cohomology is to take the homology groups of this complex, we call these the *big* noncommutative de Rham cohomology

$$H_{big}^n A = \frac{Ker \Omega^n A \xrightarrow{d} \Omega^{n+1} A}{Im \Omega^{n-1} A \xrightarrow{d} \Omega^n A}$$

EXAMPLE 7.33. Big de Rham cohomology of  $\mathbb{C} \times \mathbb{C}$ .

We have seen before that  $\Omega^n \mathbb{C} \times \mathbb{C} = (\mathbb{C} \times \mathbb{C})de^n$  and that the differential is given by

$$\begin{array}{ccc} \Omega^n \mathbb{C} \times \mathbb{C} & \xrightarrow{d} & \Omega^{n+1} \mathbb{C} \times \mathbb{C} \\ (\alpha e + \beta f)de^n & \mapsto & (\alpha - \beta)de^{n+1} \end{array}$$

From which it is immediately clear that

$$\begin{cases} H_{big}^0 \mathbb{C} \times \mathbb{C} = \mathbb{C} \\ H_{big}^n \mathbb{C} \times \mathbb{C} = 0 \end{cases}$$

for all  $n \geq 1$ . This is not quite the answer  $H^0 \mathbb{C} \times \mathbb{C} = \mathbb{C} \oplus \mathbb{C}$  we would expect from the commutative case.

For a general  $\mathbb{C}$ -algebra  $A$  it is usually very difficult to compute these cohomology groups. In case of free algebras we can use the graded structure of the complex together with the *Euler derivation* to compute them, a trick we will use later in greater generality.

EXAMPLE 7.34. Big de Rham cohomology of  $\mathbb{C}\langle x, y \rangle$ .

Define the *Euler derivation*  $E$  on  $\mathbb{C}\langle x, y \rangle$  by

$$E(x) = x \quad \text{and} \quad E(y) = y$$

Observe that if  $w$  is a *word* in  $x$  and  $y$  of degree  $k$ , then we have the Eulerian property that

$$E(w) = kw$$

as one easily verifies.

We can define a degree preserving *derivation*  $L_E$  on the differentially graded algebra  $\Omega \mathbb{C}\langle x, y \rangle$  by the rules

$$L_E(a) = E(a) \quad \text{and} \quad L_E(da) = dE(a) \quad \forall a \in \mathbb{C}\langle x, y \rangle$$


Further we introduce the degree  $-1$  *contraction operator*  $i_E$  which is the *super-derivation* on  $\Omega \mathbb{C}\langle x, y \rangle$ , that is,

$$i_E(\omega\omega') = i_E(\omega)\omega' + (-1)^i \omega i_E(\omega') \quad \text{for } \omega \in \Omega^i \mathbb{C}\langle x, y \rangle$$

defined by the rules

$$i_E(a) = 0 \quad i_E(da) = E(a) \quad \forall a \in \mathbb{C}\langle x, y \rangle.$$

That is, we have the following situation



These operators satisfy the equation

$$L_E = i_E \circ d + d \circ i_E$$

as both sides are *derivations* on  $\Omega \mathbb{C}\langle x, y \rangle$  and coincide on the generators  $a$  and  $da$  for  $a \in \mathbb{C}\langle x, y \rangle$  of this differentially graded algebra.

We claim that  $L_E$  is a total degree preserving linear automorphism on

$$\Omega^n \mathbb{C}\langle x, y \rangle \quad \text{for } n \geq 1.$$

For if  $w_i$  for  $0 \leq i \leq n$  are words in  $x$  and  $y$  of degree  $k_i$  with  $k_i \geq 1$  for  $i \geq 1$ , then we have

$$L_E(w_0 dw_1 \dots dw_n) = (k_0 + \dots + k_n)w_0 dw_1 \dots dw_n.$$

Using the words in  $x$  and  $y$  as a basis for  $\bar{A}$  we see that the kernel and image of the differential  $d$  must be homogeneous. But then, if  $\omega$  is a multi-homogeneous element in  $\Omega^n \mathbb{C}\langle x, y \rangle$  and in  $\text{Ker } d$  we have for some integer  $k \neq 0$  that

$$k\omega = L_E(\omega) = (i_E \circ d + d \circ i_E)\omega = d(i_E \omega)$$

and hence  $\omega$  lies in  $\text{Im } d$ . Therefore, we have proved


$$\begin{cases} H_{big}^0 \mathbb{C}\langle x, y \rangle = \mathbb{C} \\ H_{big}^n \mathbb{C}\langle x, y \rangle = 0 \end{cases}$$

for all  $n \geq 1$ .

The examples show that the differentially graded algebra  $\Omega A$  is *formal* for  $A = \mathbb{C} \times \mathbb{C}$  or  $\mathbb{C}\langle x, y \rangle$ . Recall that for an arbitrary  $A_\infty$ -algebra  $\Omega$  (in particular for  $\Omega \in \mathbf{dga}$ ), the homology algebra  $H^* \Omega$  has a *canonical*  $A_\infty$ -structure. That is, we have  $m_1 = 0$ ,  $m_2$  is induced by the 'multiplication'  $m_2$  on  $\Omega$  and there is a quasi-isomorphism of  $A_\infty$ -algebras  $H^* \Omega \longrightarrow \Omega$  lifting the identity of  $H^* \Omega$ .

The  $A_\infty$ -algebra  $\Omega$  is said to be *formal* if the canonical structure makes  $H^* \Omega$  into an ordinary associative graded algebra (that is, such that all  $m_n = 0$  for  $n \geq 3$ ). In particular, if  $\Omega = \Omega A$  and if the big deRham cohomology is concentrated in degree zero, then the degree properties of  $m_n$  imply that  $m_n = 0$  for  $n \geq 3$  and hence that  $\Omega A$  is formal.

Let  $A$  be an arbitrary  $\mathbb{C}$ -algebra and  $\theta \in \text{Der}_{\mathbb{C}} A$ , the Lie algebra of  $\mathbb{C}$ -algebra *derivations* of  $A$ , then we define a degree preserving derivation  $L_\theta$  and a degree  $-1$  super-derivation  $i_\theta$  on  $\Omega A$



defined by the rules

$$\begin{cases} L_\theta(a) = \theta(a) & L_\theta(da) = d \theta(a) \\ i_\theta(a) = 0 & i_\theta(da) = \theta(a) \end{cases}$$

for all  $a \in A$ . In this generality we again have the fundamental identity

$$L_\theta = i_\theta \circ d + d \circ i_\theta$$

as both sides are degree preserving derivations on  $\Omega A$  and they agree on all the generators  $a$  and  $da$  for  $a \in A$ .

LEMMA 7.35. *Let  $\theta, \gamma \in \text{Der}_{\mathbb{C}} A$ , then we have on  $\Omega A$  the following identities of operators*

$$\begin{cases} L_\theta \circ i_\gamma - i_\gamma \circ L_\theta = [L_\theta, i_\gamma] & = i_{[\theta, \gamma]} = i_{\theta \circ \gamma - \gamma \circ \theta} \\ L_\theta \circ L_\gamma - L_\gamma \circ L_\theta = [L_\theta, L_\gamma] & = L_{[\theta, \gamma]} = L_{\theta \circ \gamma - \gamma \circ \theta} \end{cases}$$

PROOF. Consider the first identity. By definition both sides are degree  $-1$  super-derivations on  $\Omega A$  so it suffices to check that they agree on generators. Clearly, both sides give 0 when evaluated on  $a \in A$  and for  $da$  we have

$$(L_\theta \circ i_\gamma - i_\gamma \circ L_\theta)da = L_\theta \gamma(a) - i_\gamma d \theta(a) = \theta \gamma(a) - \gamma \theta(a) = i_{[\theta, \gamma]}(da)$$

A similar argument proves the second identity.  $\square$

Let  $Q$  be a quiver on  $k$  vertices  $\{v_1, \dots, v_k\}$ , then we can define an *Euler derivation*  $E$  on  $\mathbb{C}Q$  by the rules that

$$E(v_i) = 0 \quad \forall 1 \leq i \leq k \quad \text{and} \quad E(a) = a \quad \forall a \in Q_a$$

By induction on the length  $l(p)$  of an oriented path  $p$  in the quiver  $Q$  one easily verifies that  $E(p) = l(p)p$ . By the lemma above we have all the necessary ingredients to redo the argument in example 7.34.

THEOREM 7.36. *For a quiver  $Q$  on  $k$  vertices, the noncommutative differential forms  $\Omega \mathbb{C}Q$  is formal. In fact, we have for the big deRham cohomology*

$$\begin{cases} H_{big}^0 \mathbb{C}Q & \simeq \mathbb{C} \times \dots \times \mathbb{C} \quad (k \text{ factors}) \\ H_{big}^n \mathbb{C}Q & \simeq 0 \quad \forall n \geq 1 \end{cases}$$

For  $\omega \in \Omega^i A$  and  $\omega' \in \Omega^j A$  we define the *super-commutator* to be

$$[\omega, \omega'] = \omega\omega' - (-1)^{ij}\omega'\omega$$

That is, it is the usual commutator unless both  $i$  and  $j$  are odd in which case it is the sum  $\omega\omega' + \omega'\omega$ .

As the differential  $d$  is a super-derivation on  $\Omega A$  we have that

$$d([\omega, \omega']) = [d\omega, \omega'] + (-1)^i[\omega, d\omega']$$

and therefore the differential maps the *subspaces* of super-commutators to subspaces of super-commutators. Therefore, if we define

$$\text{DR}^n A = \frac{\Omega^n A}{\sum_{i=0}^n [\Omega^i A, \Omega^{n-i} A]}$$

Then the  $\text{dgalg}$ -structure on  $\Omega A$  induces one on the complex

$$\text{DR}^0 A \xrightarrow{d} \text{DR}^1 A \xrightarrow{d} \text{DR}^2 A \xrightarrow{d} \dots$$

which is called the *Karoubi complex* of  $A$ .

We define the *noncommutative de Rham cohomology groups* of  $A$  to be the homology of the Karoubi complex, that is

$$H_{dR}^n A = \frac{\text{Ker } DR^n A \xrightarrow{d} DR^{n+1} A}{\text{Im } DR^{n-1} A \xrightarrow{d} DR^n A}$$

EXAMPLE 7.37. Noncommutative de Rham cohomology of  $\mathbb{C} \times \mathbb{C}$ . Recall that the product on  $\Omega \mathbb{C} \times \mathbb{C}$  is given by the formula

$$(\alpha e + \beta f)de^n(\gamma e + \delta f)de^m = \begin{cases} (\alpha\gamma e + \beta\delta f)de^{n+m} & \text{when } n \text{ is even} \\ (\alpha\delta e + \beta\gamma f)de^{n+m} & \text{when } n \text{ is odd} \end{cases}$$

If  $m$  is odd, then we deduce from this that the commutator

$$[\alpha e + \beta f, (\gamma e + \delta f)de^m] = (\alpha - \beta)(\gamma e - \delta f)de^m$$

and hence we can write any element of  $\Omega^m \mathbb{C} \times \mathbb{C} = (\mathbb{C} \times \mathbb{C})de^m$  as a (super) commutator, whence

$$DR^m \mathbb{C} \times \mathbb{C} = 0 \quad \text{when } m \text{ is odd.}$$

On the other hand, if  $m$  is even then any commutator with  $k$  even

$$[(\alpha e + \beta f)de^k, (\gamma e + \delta f)de^{m-k}] = 0$$

whereas if  $k$  is odd we have

$$[(\alpha e + \beta f)de^k, (\gamma e + \delta f)de^{m-k}] = (\alpha\delta + \beta\gamma)de^m$$

As a consequence the space of super-commutators in  $\Omega^m \mathbb{C} \times \mathbb{C}$  is one dimensional and therefore

$$DR^m \mathbb{C} \times \mathbb{C} = \mathbb{C} \quad \text{when } m \text{ is even and } > 0.$$

Thus, the Karoubi complex of  $\mathbb{C} \times \mathbb{C}$  has the following form


$$\mathbb{C} \times \mathbb{C} \xrightarrow{d} 0 \xrightarrow{d} \mathbb{C} \xrightarrow{d} 0 \xrightarrow{d} \mathbb{C} \xrightarrow{d} 0 \xrightarrow{d} \dots$$

and therefore we have for the noncommutative de Rham cohomology groups

$$H_{dR}^n \mathbb{C} \times \mathbb{C} = \begin{cases} \mathbb{C} \times \mathbb{C} & \text{when } n = 0 \\ 0 & \text{when } n \text{ is odd} \\ \mathbb{C} & \text{when } n \text{ is even and } > 0. \end{cases}$$

EXAMPLE 7.38. Noncommutative de Rham cohomology of  $\mathbb{C}\langle x, y \rangle$ .

Consider again the Eulerian derivation  $E$  on  $\mathbb{C}\langle x, y \rangle$  and the operators  $L_E$  and  $i_E$  on  $\Omega \mathbb{C}\langle x, y \rangle$ . Repeating the above argument that  $d$  is compatible with the subspaces of super-commutators for  $i_E$  and  $L_E$  we see that we have induced operations



We have again that  $L_E$  is an isomorphism on  $DR^n \mathbb{C}\langle x, y \rangle$  whenever  $n \geq 1$  and again we deduce from the equality  $L_E = i_E \circ d + d \circ i_E$  that

$$H_{dR}^n \mathbb{C}\langle x, y \rangle = \begin{cases} \mathbb{C} & \text{when } n = 0, \\ 0 & \text{when } n \geq 1. \end{cases}$$

THEOREM 7.39. Let  $Q$  be a quiver on  $k$  vertices, then the Karoubi complex of  $\mathbb{C}Q$  is acyclic. In particular,

$$\begin{cases} H_{dR}^0 \mathbb{C}Q \simeq \mathbb{C} \times \dots \times \mathbb{C} \text{ (} k \text{ factors)} \\ H_{dR}^n \mathbb{C}Q \simeq 0 \quad \forall n \geq 1 \end{cases}$$

So far we have considered differential forms with respect to the basefield  $\mathbb{C}$ . Sometimes it is useful to consider only the *relative differential forms* on  $A$  with respect to a subalgebra  $B$ . These can be defined as follows.

Let  $\bar{A}_B$  be the cokernel of the inclusion  $B \hookrightarrow A$  in the category  $B$ -bimod of bimodules over  $B$ . We define the space of relative differential forms of degree  $n$  with respect to  $B$  to be

$$\Omega_B^n A = A \otimes_B \underbrace{\bar{A}_B \otimes_B \dots \otimes_B \bar{A}_B}_n$$

By definition  $\Omega_B^n A$  is the quotient space of  $\Omega^n A$  by the relations

$$\begin{aligned} (a_0, \dots, a_{i-1}b, a_i, \dots, a_n) &= (a_0, \dots, a_{i-1}, ba_i, \dots, a_n) \\ (a_0, \dots, a_{i-1}, s, a_{i+1}, \dots, a_n) &= 0 \end{aligned}$$

for all  $b \in B$  and  $1 \leq i \leq n$ . One verifies that the multiplication and differential defined on  $\Omega A$  are compatible with these relations, making  $\Omega_B A$  an object in  $\mathbf{dgalg}$ . Moreover, there is a canonical epimorphism

$$\Omega A \twoheadrightarrow \Omega_B A \quad \text{in } \mathbf{dgalg}.$$

We will now determine the kernel. First we give the universal property for  $\Omega_B A$ . Given  $\Gamma = \oplus \Gamma^n$  in  $\mathbf{dgalg}$  and an algebra map  $A \xrightarrow{f} \Gamma^0$  such that  $d(f B) = 0$ , then there is a unique morphism in  $\mathbf{dgalg}$  making the diagram commute

$$\begin{array}{ccc} \Omega_B A & \xrightarrow{\exists f_*} & \Gamma \\ \uparrow & & \uparrow \\ A & \xrightarrow{f} & \Gamma^0 \end{array}$$

Indeed, by the universal property of  $\Omega A$  there is a unique morphism  $\Omega A \xrightarrow{f_*} \Gamma$  in  $\mathbf{dgalg}$  extending  $f$  given by

$$f_*(a_0 da_1 \dots da_n) = f(a_0) d(f(a_1)) \dots d(f(a_n)).$$

If  $d(f B) = 0$  then one verifies that  $f_*$  is compatible with the relations defining  $\Omega_B A$ , proving the universal property.

PROPOSITION 7.40. *For a subalgebra  $B$  of  $A$  we have an isomorphism in  $\mathbf{dgalg}$*

$$\Omega_B A = \frac{\Omega A}{\Omega A d(B) \Omega A}$$

PROOF. The ideal generated by  $d(B)$  is closed under  $d$  and therefore the quotient is an object in  $\mathbf{dgalg}$  with the same universal property as  $\Omega_B A$ .  $\square$

An important special case is when  $B = \mathbb{C} \times \dots \times \mathbb{C}$  is the subalgebra of  $\mathbb{C}Q$  generated by the vertex-idempotents. In this case we will denote

$$\Omega_{rel} \mathbb{C}Q = \Omega_B \mathbb{C}Q$$

and call it the relative differential forms on  $Q$ .

LEMMA 7.41. *Let  $Q$  be a quiver on  $k$  vertices, then a basis for  $\Omega_{rel}^n \mathbb{C}Q$  is given by the elements*

$$p_0 dp_1 \dots dp_n$$

where  $p_i$  is an oriented path in the quiver such that length  $p_0 \geq 0$  and length  $p_i \geq 1$  for  $1 \leq i \leq n$  and such that the starting point of  $p_i$  is the endpoint of  $p_{i+1}$  for all  $1 \leq i \leq n - 1$ .

PROOF. Clearly  $l(p_i) \geq 1$  when  $i \geq 1$  or  $p_i$  would be a vertex-idempotent whence in  $B$ . Let  $v$  be the starting point of  $p_i$  and  $w$  the end point of  $p_{i+1}$  and assume that  $v \neq w$ , then

$$p_i \otimes_B p_{i+1} = p_i v \otimes_B w p_{i+1} = p_i v w \otimes_B p_{i+1} = 0$$

from which the assertion follows. □

We define the *big relative de Rham* cohomology groups of  $A$  with respect to  $B$  to be the cohomology of the complex


$$\Omega_B^0 A \xrightarrow{d} \Omega_B^1 A \xrightarrow{d} \Omega_B^2 A \xrightarrow{d} \dots$$

that is,

$$H_B^n A = \frac{Ker \Omega^n A \xrightarrow{d} \Omega^{n+1} A}{Im \Omega^{n-1} A \xrightarrow{d} \Omega^n A}$$

In the case of path algebras of quivers, we can use the grading by length on paths and the Eulerian derivation to compute these relative de Rham groups.

EXAMPLE 7.42. Big relative de Rham cohomology.  
Let  $\mathbb{M}$  (resp.  $\mathbb{C} \times \mathbb{C}$ ) be the path algebras of the quivers



The Eulerian derivation  $E$  on  $\mathbb{M}$  is defined by

$$E(e) = E(f) = 0 \quad E(x) = x \quad E(y) = y \quad E(u) = u \quad \text{and} \quad E(v) = v.$$

Observe that  $E$  respects all relation holding in  $\mathbb{M}$  and so is indeed a  $\mathbb{C} \times \mathbb{C}$ - derivation of  $\mathbb{M}$ .

As before we define a degree preserving derivation  $L_E$  and a degree  $-1$  super-derivation  $i_E$  on  $\Omega_{rel} \mathbb{M} = \Omega_{\mathbb{C} \times \mathbb{C}} \mathbb{M}$  by the rules

$$\begin{cases} L_E(a) = E(a) & L_E(da) = dE(a) \\ i_E(a) = 0 & i_E(da) = E(a) \end{cases}$$

for all  $a \in \mathbb{M}$ . We have the equality

$$L_E = i_E \circ d + d \circ i_E$$

and arguing as before we obtain that

$$H_{rel}^n \mathbb{M} = \begin{cases} \mathbb{C} \times \mathbb{C} & \text{when } n = 0, \\ 0 & \text{when } n \geq 1. \end{cases}$$

THEOREM 7.43. *Let  $Q$  be a quiver on  $k$  vertices, then the relative differential forms  $\Omega_{rel} \mathbb{C}Q$  is a formal differentially graded algebra. In fact,*

$$\begin{cases} H_{rel}^0 \mathbb{C}Q \simeq \mathbb{C} \times \dots \times \mathbb{C} \text{ (} k \text{ factors)} \\ H_{rel}^n \mathbb{C}Q \simeq 0 \quad \forall n \geq 1 \end{cases}$$

We can repeat the construction of the Karoubi complex verbatim for relative differential operators and define a *relative Karoubi complex*

$$DR_B^0 A \xrightarrow{d} DR_B^1 A \xrightarrow{d} DR_B^2 A \xrightarrow{d} \dots$$

where

$$DR_B^n A = \frac{\Omega_B^n A}{\sum_{i=0}^n [\Omega_B^i A, \Omega_B^{n-i} A]}$$

Clearly, we then define the *noncommutative relative de Rham cohomology groups* of  $A$  with respect to  $B$  to be the homology of this complex

$$H_{B,dR}^n A = \frac{Ker DR_B^n A \xrightarrow{d} DR_B^{n+1} A}{Im DR_B^{n-1} A \xrightarrow{d} DR_B^n A}$$

Let  $\theta \in Der_B A$ , that is  $\theta$  is a  $\mathbb{C}$ -derivation on  $A$  such that  $\theta(b) = 0$  for every  $b \in B$ . Then, as

$$L_\theta(db) = d\theta(b) = 0 \quad \text{and} \quad i_\theta(db) = \theta(b) = 0$$

we see that the operators  $L_\theta$  and  $i_\theta$  can be defined on the relative forms

$$\Omega_B A = \frac{\Omega A}{\Omega A dB \Omega A}$$

and also on the relative Karoubi complex. Again, these induced operators satisfy the identities of lemma 7.35. In the special case of the Eulerian derivation  $E$  on the path algebra  $\mathbb{C}Q$  we see that  $E \in Der_B \mathbb{C}Q$  and hence we have the following result.

THEOREM 7.44. *Let  $Q$  be a quiver on  $k$  vertices. Then, the relative Karoubi complex is acyclic. That is,*

$$\begin{cases} H_{rel,dR}^0 \mathbb{C}Q \simeq \mathbb{C} \times \dots \times \mathbb{C} \text{ (} k \text{ factors)} \\ H_{rel,dR}^n \mathbb{C}Q \simeq 0 \quad \forall n \geq 1 \end{cases}$$

### 7.7. Symplectic structure.

Let  $Q$  be a quiver on  $k$  vertices  $\{v_1, \dots, v_k\}$ . We will determine the first terms in the relative Karoubi complex. Define

$$dR_{rel}^n \mathbb{C}Q = \frac{\Omega_{rel}^n \mathbb{C}Q}{\sum_{i=0}^n [\Omega_{rel}^i \mathbb{C}Q, \Omega_{rel}^{n-i} \mathbb{C}Q]}$$

In the commutative case,  $dR^0$  are the functions on the manifold and  $dR^1$  the 1-forms. We will characterize the *noncommutative functions* and *noncommutative 1-forms* in the case of quivers.

Recall that a *necklace word*  $w$  in the quiver  $Q$  is an equivalence class of an oriented cycle  $c = a_1 \dots a_l$  of length  $l \geq 0$  in  $Q$ , where  $c \sim c'$  if  $c'$  is obtained from  $c$  by cyclicly permuting the composing arrows  $a_i$ .

LEMMA 7.45. *A  $\mathbb{C}$ -basis for the noncommutative functions*

$$dR_{rel}^0 \mathbb{C}Q \simeq \frac{\mathbb{C}Q}{[\mathbb{C}Q, \mathbb{C}Q]}$$

are the necklace words in the quiver  $Q$ .

PROOF. Let  $\mathbb{W}$  be the  $\mathbb{C}$ -space spanned by all necklace words  $w$  in  $Q$  and define a linear map

$$\mathbb{C}Q \xrightarrow{n} \mathbb{W} \quad \begin{cases} p \mapsto w_p & \text{if } p \text{ is a cycle} \\ p \mapsto 0 & \text{if } p \text{ is not} \end{cases}$$

for all oriented paths  $p$  in the quiver  $Q$ , where  $w_p$  is the necklace word in  $Q$  determined by the oriented cycle  $p$ . Because  $w_{p_1 p_2} = w_{p_2 p_1}$  it follows that the commutator subspace  $[\mathbb{C}Q, \mathbb{C}Q]$  belongs to the kernel of this map. Conversely, let

$$x = x_0 + x_1 + \dots + x_m$$

be in the kernel where  $x_0$  is a linear combination of non-cyclic paths and  $x_i$  for  $1 \leq i \leq m$  is a linear combination of cyclic paths mapping to the same necklace word  $w_i$ , then  $n(x_i) = 0$  for all  $i \geq 0$ . Clearly,  $x_0 \in [\mathbb{C}Q, \mathbb{C}Q]$  as we can write every noncyclic path  $p = a.p' = a.p' - p'.a$  as a commutator. If  $x_i = a_1 p_1 + a_2 p_2 + \dots + a_l p_l$  with  $n(p_i) = w_i$ , then  $p_1 = q.q'$  and  $p_2 = q'.q$  for some paths  $q, q'$  whence  $p_1 - p_2$  is a commutator. But then,  $x_i = a_1(p_1 - p_2) + (a_2 - a_1)p_2 + \dots + a_l p_l$  is a sum of a commutator and a linear combination of strictly fewer elements. By induction, this shows that  $x_i \in [\mathbb{C}Q, \mathbb{C}Q]$ .  $\square$

If we fix a dimension vector  $\alpha$ , then taking traces defines a map

$$dR^0 \mathbb{C}Q \xrightarrow{tr} \mathbb{C}[rep_\alpha Q]$$

whence noncommutative functions determine  $GL(\alpha)$ -invariant commutative functions on the representation space  $rep_\alpha Q$  and hence commutative functions on the quotient varieties  $iss_\alpha Q$ . In fact, we have seen that the image  $tr(dR^0 \mathbb{C}Q)$  generates the ring of polynomial invariants  $\mathbb{C}[rep_\alpha Q]^{GL(\alpha)} = \mathbb{C}[iss_\alpha Q]$ .

LEMMA 7.46.  *$dR_{rel}^1 \mathbb{C}Q$  is isomorphic as  $\mathbb{C}$ -space to*

$$\bigoplus_{\begin{array}{c} \textcircled{j} \xleftarrow{a} \textcircled{i} \end{array}} v_i \cdot \mathbb{C}Q \cdot v_j \cdot da = \bigoplus_{\begin{array}{c} \textcircled{j} \xleftarrow{a} \textcircled{i} \end{array}} \textcircled{i} \xleftarrow{\quad} \textcircled{j} \xleftarrow{a} \textcircled{i}$$

PROOF. If  $p.q$  is not a cycle, then  $pdq = [p, dq]$  and so vanishes in  $dR_{rel}^1 \mathbb{C}Q$  so we only have to consider terms  $pdq$  with  $p.q$  an oriented cycle in  $Q$ . For any three paths  $p, q$  and  $r$  in  $Q$  we have the equality


$$[p.qdr] = pqdr - qd(rp) + qrdp$$

whence in  $dR_{rel}^1 \mathbb{C}Q$  we have relations allowing to reduce the length of the differential part

$$qd(rp) = pqdr + qrdp$$



so  $dR_{rel}^1 \mathbb{C}Q$  is spanned by terms of the form  $pda$  with  $a \in Q_a$  and  $p.a$  an oriented cycle in  $Q$ . Therefore, we have a surjection

$$\Omega_{rel}^1 \mathbb{C}Q \longrightarrow \bigoplus_{\text{cycles } a} v_i \cdot \mathbb{C}Q \cdot v_j \cdot da$$



By construction, it is clear that  $[\Omega_{rel}^0 \mathbb{C}Q, \Omega_{rel}^1 \mathbb{C}Q]$  lies in the kernel of this map and using an argument as in the lemma above one shows also the converse inclusion.  $\square$

EXAMPLE 7.47.  $dR_{rel}^i \mathbb{M}$ .

Take the path algebra  $\mathbb{M}$  of the quiver of example 7.42. Noncommutative functions on  $\mathbb{M}$  are the 0-forms, which is by definition the quotient space

$$dR_{rel}^0 \mathbb{M} = \frac{\mathbb{M}}{[\mathbb{M}, \mathbb{M}]}$$

If  $p$  is an oriented path of length  $\geq 1$  in the quiver with different begin- and endpoint, then we can write  $p$  as a concatenation  $p = p_1 p_2$  with  $p_i$  an oriented path of length  $\geq 0$  such that  $p_2 p_1 = 0$  in  $\mathbb{M}$ . As  $[p_1, p_2] = p_1 p_2 - p_2 p_1 = 0$  in  $dR_{rel}^0 \mathbb{M}$  we deduce that the space of noncommutative functions on  $\mathbb{M}$  has as  $\mathbb{C}$ -basis the necklace words  $w$



where each bead is this time one of the elements

$$\blacksquare = x \quad \square = y \quad \text{and} \quad \blacktriangledown = uv$$

together with the necklace words of length zero  $e$  and  $f$ . Each necklace word  $w$  corresponds to the equivalence class of the words in  $\mathbb{M}$  obtained from multiplying the beads in the indicated orientation and two words in  $\{x, y, u, v\}$  in  $\mathbb{M}$  are said to be equivalent if they are identical up to cyclic permutation of the terms.

Substituting each bead with the  $n \times n$  matrices specified before and taking traces we get a map

$$dR_{rel}^0 \mathbb{M} = \frac{\mathbb{M}}{[\mathbb{M}, \mathbb{M}]} \xrightarrow{tr} \mathbb{C}[rep_\alpha \mathbb{M}]$$

Hence, noncommutative functions on  $\mathbb{M}$  induce ordinary functions on all the representation spaces  $rep_\alpha \mathbb{M}$  and these functions are  $GL(\alpha)$ -invariant. Moreover, the image of this map generates the ring of polynomial invariants as we mentioned before.

Next, we consider noncommutative 1-forms on  $\mathbb{M}$  which are by definition elements of the space

$$dR_{rel}^1 \mathbb{M} = \frac{\Omega_{rel}^1 \mathbb{M}}{[\mathbb{M}, \Omega_{rel}^1 \mathbb{M}]}$$

Recall that  $\Omega_{rel}^1 \mathbb{M}$  is spanned by the expressions  $p_0 dp_1$  with  $p_0$  resp.  $p_1$  oriented paths in the quiver of length  $\geq 0$  resp.  $\geq 1$  and such that the starting point of  $p_0$  is the end point of  $p_1$ . To form  $dR_{rel}^1 \mathbb{M}$  we have to divide out expressions such as

$$[p, p_0 dp_1] = pp_0 dp_1 + p_0 p_1 dp - p_0 d(p_1 p)$$

That is, if we have connecting oriented paths  $p_2$  and  $p_1$  both of length  $\geq 1$  we have in  $dR_{rel}^1 \mathbb{M}$

$$p_0 d(p_1 p_2) = p_2 p_0 dp_1 + p_0 p_1 dp_2$$

and by iterating this procedure whenever the differential term is a path of length  $\geq 2$  we can represent each class in  $dR_{rel}^1 \mathbb{M}$  as a combination from

$$\mathbb{M}e \, dx + \mathbb{M}e \, dy + \mathbb{M}e \, du + \mathbb{M}f \, dv$$

Now,  $\mathbb{M}e = e\mathbb{M}e + f\mathbb{M}e$  and let  $p \in f\mathbb{M}e$ . Then, we have in  $dR_{rel}^1 \mathbb{M}$

$$d(xp) = p \, dx + x \, dp$$


but by our description of  $\Omega^1 \mathbb{M}$  the left hand term is zero as is the second term on the right, whence  $p \, dx = 0$ . A similar argument holds replacing  $x$  by  $y$ . As for  $u$ , let  $p \in e\mathbb{M}e$ , then we have in  $dR_{rel}^1 \mathbb{M}$

$$d(up) = p \, du + u \, dp$$

and again the left-hand and the second term on the right are zero whence  $p \, du = 0$ . An analogous result holds for  $v$  and  $p \in f\mathbb{M}f$ . Therefore, we have the description of noncommutative 1-forms on  $\mathbb{M}$

$$dR_{rel}^1 \mathbb{M} = e\mathbb{M}e \, dx + e\mathbb{M}e \, dy + f\mathbb{M}e \, du + e\mathbb{M}f \, dv$$

That is, in graphical terms




Using the above descriptions of  $dR_{rel}^i \mathbb{C}Q$  for  $i = 0, 1$  and the differential  $dR_{rel}^0 \mathbb{C}Q \xrightarrow{d} dR_{rel}^1 \mathbb{C}Q$  we can define *partial differential operators* associated to any arrow  $(j \xleftarrow{a} i)$  in  $Q$ .

$$\frac{\partial}{\partial a} : dR_{rel}^0 \mathbb{C}Q \longrightarrow v_i \mathbb{C}Q v_j \quad \text{by} \quad df = \sum_{a \in Q_a} \frac{\partial f}{\partial a} da$$

To take the partial derivative of a necklace word  $w$  with respect to an arrow  $a$ , we run through  $w$  and each time we encounter  $a$  we open the necklace by removing that occurrence of  $a$  and then take the sum of all the paths obtained.

EXAMPLE 7.48. For the path algebra  $\mathbb{M}$  we have the partial differential operators



Recall that a *symplectic structure* on a (commutative) manifold  $M$  is given by a closed differential 2-form. The non-degenerate 2-form  $\omega$  gives a canonical isomorphism

$$T M \simeq T^* M$$

that is, between vector fields on  $M$  and differential 1-forms. Further, there is a unique  $\mathbb{C}$ -linear map from functions  $f$  on  $M$  to vectorfields  $\xi_f$  by the requirement that  $-df = i_{\xi_f}\omega$  where  $i_\xi$  is the contraction of  $n$ -forms to  $n - 1$ -forms using the vectorfield  $\xi$ . We can make the functions on  $M$  into a *Poisson algebra* by defining

$$\{f, g\} = \omega(\xi_f, \xi_g)$$

and one verifies that this bracket satisfies the Jacobi and Leibnitz identities.

The *Lie derivative*  $L_\xi$  with respect to  $\xi$  is defined by the Cartan homotopy formula

$$L_\xi \varphi = i_\xi d\varphi + di_\xi \varphi$$

for any differential form  $\varphi$ . A vectorfield  $\xi$  is said to be *symplectic* if it preserves the symplectic form, that is,  $L_\xi \omega = 0$ . In particular, for any function  $f$  on  $M$  we have that  $\xi_f$  is symplectic. Moreover the assignment

$$f \longrightarrow \xi_f$$

defines a Lie algebra morphism from the functions  $\mathcal{O}(M)$  on  $M$  equipped with the Poisson bracket to the Lie algebra of symplectic vectorfields,  $Vect_\omega M$ . Moreover, this map fits into the exact sequence

$$0 \longrightarrow \mathbb{C} \longrightarrow \mathcal{O}(M) \longrightarrow Vect_\omega M \longrightarrow H_{dR}^1 M \longrightarrow 0$$

Recall the definition of the double quiver  $Q^d$  of a quiver  $Q$  given in section 5.6 by assigning to every arrow  $a \in Q_a$  an arrow  $a^*$  in  $Q^d$  in the opposite direction.

DEFINITION 7.49. The canonical *noncommutative symplectic structure* on the double quiver  $Q^d$  is given by the element

$$\omega = \sum_{a \in Q_a} dada^* \in \mathbf{dR}_{rel}^2 \mathbb{C}Q^d$$

We will use  $\omega$  to define a correspondence between the noncommutative 1-forms  $\mathbf{dR}_{rel}^1 \mathbb{C}Q^d$  and the *noncommutative vectorfields* which are defined to be  $B = \mathbb{C}Q^v$ -derivations of the path algebra  $\mathbb{C}Q^d$ . Recall that if  $\theta \in Der_B \mathbb{C}Q^d$  we define operators  $L_\theta$  and  $i_\theta$  on  $\Omega \mathbb{C}Q^d$  and on  $\mathbf{dR} \mathbb{C}Q^d$  by the rules

$$\begin{cases} L_\theta(a) = \theta(a) & L_\theta(da) = d\theta(a) \\ i_\theta(a) = 0 & i_\theta(da) = \theta(a) \end{cases}$$

and that the following identities are satisfied for all  $\theta, \gamma \in Der_B \mathbb{C}Q^d$

$$[L_\theta, L_\gamma] = L_{[\theta, \gamma]} \quad \text{and} \quad [i_\theta, i_\gamma] = i_{[\theta, \gamma]}$$

These operators allow us to define a linear map

$$Der_B \mathbb{C}Q \xrightarrow{\tau} \mathbf{dR}_{rel}^1 \mathbb{C}Q \quad \text{by} \quad \tau(\theta) = i_\theta(\omega)$$

We claim that this is an isomorphism. Indeed, every  $B$ -derivation  $\theta$  on  $\mathbb{C}Q^d$  is fully determined by its image on the arrows in  $Q^d$  which satisfy if  $a = \textcircled{j} \xleftarrow{a} \textcircled{i}$

$$\theta(a) = \theta(v_j a v_i) = v_j \theta(a) v_i \in v_j \mathbb{C}Q^d v_i$$

so determines an element  $\theta(a)da^* \in \mathbf{dR}_{rel}^1 \mathbb{C}Q^d$ . Further, we compute

$$\begin{aligned} i_\theta(\omega) &= \sum_{a \in Q_a} i_\theta(da)da^* - i_\theta(da^*)da \\ &= \sum_{a \in Q_a} \theta(a)da^* - \theta(a^*)da \end{aligned}$$

which lies in  $\mathbf{dR}_{rel}^1 \mathbb{C}Q^d$ . As both  $B$ -derivations and 1-forms are determined by their coefficients,  $\tau$  is indeed bijective.

**EXAMPLE 7.50.** For the path algebra of the double quiver  $\mathbb{M}$ , the analogon of the classical isomorphism  $T M \simeq T^* M$  is the isomorphism

$$Der_{\mathbb{C} \times \mathbb{C}} \mathbb{M} \xrightarrow{i_\omega} \mathbf{dR}_{rel}^1 \mathbb{M}$$

as for any  $\mathbb{C} \times \mathbb{C}$ -derivation  $\theta$  we have

$$\begin{aligned} i_\theta \omega &= i_\theta(dx)dy - dx i_\theta(dy) + i_\theta(du)dv - du i_\theta(dv) \\ &= \theta(x)dy - dx\theta(y) + \theta(u)dv - du\theta(v) \\ &\equiv \theta(x)dy - \theta(y)dx + \theta(u)dv - \theta(v)du \end{aligned}$$

and using the relations in  $\mathbb{M}$  we can easily prove that any  $\mathbb{C} \times \mathbb{C}$  derivation on  $\mathbb{M}$  must satisfy

$$\theta(x) \in e\mathbb{M}e \quad \theta(y) \in e\mathbb{M}e \quad \theta(u) \in e\mathbb{M}f \quad \theta(v) \in f\mathbb{M}e$$

so the above expression belongs to  $\mathbf{dR}_{rel}^1 \mathbb{M}$ . Conversely, any  $\theta$  defined by its images on the generators  $x, y, u$  and  $v$  by

$$-\theta(y)dx + \theta(x)dy - \theta(v)du + \theta(u)dv \in \mathbf{dR}_{rel}^1 \mathbb{M}$$

induces a derivation on  $\mathbb{M}$ .

In analogy with the commutative case we define a derivation  $\theta \in Der_B \mathbb{C}Q^d$  to be *symplectic* if and only if  $L_\theta \omega = 0 \in \mathbf{dR}_{rel}^2 \mathbb{C}Q^d$ . We will denote the subspace of symplectic derivations by  $Der_\omega \mathbb{C}Q$ . It follows from the noncommutative analogon of the Cartan homotopy equality

$$L_\theta = i_\theta \circ d + d \circ i_\theta$$

and the fact that  $\omega$  is a closed form, that  $\theta \in Der_\omega \mathbb{C}Q^d$  implies

$$L_\theta \omega = di_\theta \omega = \tau(\theta) = 0$$

That is,  $\tau(\theta)$  is a closed form which by the acyclicity of the Karoubi complex shows that it must be an exact form. That is we have an isomorphism of exact sequences of  $\mathbb{C}$ -vectorspaces

$$\begin{array}{ccccccc} 0 & \longrightarrow & B & \longrightarrow & \mathbf{dR}_{rel}^0 \mathbb{C}Q^d & \xrightarrow{d} & (\mathbf{dR}_{rel}^1 \mathbb{C}Q)_{exact} \longrightarrow 0 \\ & & \downarrow = & & \downarrow \simeq & & \downarrow \tau^{-1} \\ 0 & \longrightarrow & B & \longrightarrow & \frac{\mathbb{C}Q^d}{[\mathbb{C}Q^d, \mathbb{C}Q^d]} & \longrightarrow & Der_\omega \mathbb{C}Q^d \longrightarrow 0 \end{array}$$

in the next section we will show that this is in fact an exact sequence of Lie algebras.




FIGURE 4. Kontsevich bracket  $\{w_1, w_2\}_K$ .

**7.8. Necklace Lie algebras.**


Let  $Q$  be a quiver on  $k$  vertices,  $Q^d$  its double and  $\omega = \sum_{a \in Q_a} dada^*$  the canonical symplectic form on  $\mathbb{C}Q^d$ . Recall from last section the definition of the partial differential operators  $\frac{\partial}{\partial a}$  for an arrow  $a$  in  $Q^d$ .


DEFINITION 7.51. The *Kontsevich bracket* on the necklace words in  $Q^d$ ,  $dR_{rel}^0 \mathbb{C}Q^d$  is defined to be

$$\{w_1, w_2\}_K = \sum_{a \in Q_a} \left( \frac{\partial w_1}{\partial a} \frac{\partial w_2}{\partial a^*} - \frac{\partial w_1}{\partial a^*} \frac{\partial w_2}{\partial a} \right) \text{ mod } [\mathbb{C}Q^d, \mathbb{C}Q^d]$$

That is, to compute  $\{w_1, w_2\}_K$  we consider for every arrow  $a \in Q_a$  all occurrences of  $a$  in  $w_1$  and  $a^*$  in  $w_2$ . We then open up the necklaces removing these factors and gluing the open ends together to form a new necklace word. We then replace the roles of  $a^*$  and  $a$  and redo this operation (with a minus sign), see figure 4. Finally, we add all the obtained necklace words.

EXAMPLE 7.52. For the path algebra  $M$  the canonical symplectic form is  $\omega = dx dy + du dv$ . Using the above graphical description we have that the Kontsevich bracket  $\{w_1, w_2\}_K$  is equal to





Using this graphical description of the Kontsevich bracket, it is an enjoyable exercise to verify that the bracket turns  $dR_{rel}^0 \mathbb{C}Q^d$  into a Lie algebra. That is, for all necklace words  $w_i$ , the bracket satisfies the *Jacobi identity*

$$\{\{w_1, w_2\}_K, w_3\}_K + \{\{w_2, w_3\}_K, w_1\}_K + \{\{w_3, w_1\}_K, w_2\}_K = 0$$

see figure 5.

Recall the exact commutative diagram from last section

$$\begin{array}{ccccccc} 0 & \longrightarrow & B & \longrightarrow & dR_{rel}^0 \mathbb{C}Q^d & \xrightarrow{d} & (dR_{rel}^1 \mathbb{C}Q)_{exact} \longrightarrow 0 \\ & & \downarrow = & & \downarrow \simeq & & \downarrow \tau^{-1} \\ 0 & \longrightarrow & B & \longrightarrow & \frac{\mathbb{C}Q^d}{[\mathbb{C}Q^d, \mathbb{C}Q^d]} & \longrightarrow & Der_w \mathbb{C}Q^d \longrightarrow 0 \end{array}$$

Clearly, the symplectic derivations  $Der_w \mathbb{C}Q^d$  are equipped with a Lie algebra structure via  $[\theta_1, \theta_2] = \theta_1 \circ \theta_2 - \theta_2 \circ \theta_1$ .

For every necklace word  $w$  we have a derivation  $\theta_w = \tau^{-1}dw$  which is defined by

$$\begin{cases} \theta_w(a) &= \frac{\partial w}{\partial a^*} \\ \theta_w(a^*) &= -\frac{\partial w}{\partial a} \end{cases}$$

With this notation we get the following interpretations of the Kontsevich bracket

$$\{w_1, w_2\}_K = i_{\theta_{w_1}}(i_{\theta_{w_2}}\omega) = L_{\theta_{w_1}}(w_2) = -L_{\theta_{w_2}}(w_1)$$

where the next to last equality follows because  $i_{\theta_{w_2}}\omega = dw_2$  and the fact that  $i_{\theta_{w_1}}(dw) = L_{\theta_{w_1}}(w)$  for any  $w$ . More generally, for any  $B$ -derivation  $\theta$  and any necklace word  $w$  we have the equation

$$i_{\theta}(i_{\theta_w}\omega) = L_{\theta}(w)$$

By the commutation relations for the operators  $L_{\theta}$  and  $i_{\theta}$  we have for all  $B$ -derivations  $\theta_i$  the equalities

$$\begin{aligned} L_{\theta_1}i_{\theta_2}i_{\theta_3}\omega - i_{\theta_2}i_{\theta_3}L_{\theta_1}\omega &= [L_{\theta_1}, i_{\theta_2}]i_{\theta_3}\omega + i_{\theta_2}L_{\theta_1}i_{\theta_3}\omega \\ &\quad - i_{\theta_2}L_{\theta_1}i_{\theta_3}\omega + i_{\theta_2}[L_{\theta_1}, i_{\theta_3}]\omega \\ &= i_{[\theta_1, \theta_2]}i_{\theta_3}\omega + i_{\theta_2}i_{[\theta_1, \theta_3]}\omega \end{aligned}$$




FIGURE 5. Jacobi identity for the Kontsevich bracket. Term 1a vanishes against 2c, term 1b against 3d, 1c against 3a, 1d against 2b, 2a against 3c and 2d against 3b.

By the homotopy formula we have  $L_{\theta_w}\omega = 0$  for every necklace word  $w$ , whence we get

$$L_{\theta_{w_1}}i_{\theta_2}i_{\theta_3}\omega = i_{[\theta_{w_1},\theta_2]}i_{\theta_3}\omega + i_{\theta_2}i_{[\theta_{w_1},\theta_3]}\omega$$

Take  $\theta_2 = \theta_{w_2}$ , then the left hand side is equal to

$$\begin{aligned} L_{\theta_{w_1}}i_{\theta_{w_2}}i_{\theta_3}\omega &= -L_{\theta_{w_1}}i_{\theta_3}i_{\theta_{w_2}}\omega \\ &= -L_{\theta_{w_1}}L_{\theta_3}w_2 \end{aligned}$$

whereas the last term on the right equals

$$\begin{aligned} i_{\theta_{w_2}}i_{[\theta_{w_1},\theta_3]}\omega &= -i_{[\theta_{w_1},\theta_3]}i_{\theta_{w_2}}\omega \\ &= -L_{[\theta_{w_1},\theta_3]}w_2 = -L_{\theta_{w_1}}L_{\theta_3}w_2 + L_{\theta_3}L_{\theta_{w_1}}w_2 \end{aligned}$$

and substituting this we obtain that

$$\begin{aligned} i_{[\theta_{w_1},\theta_{w_2}]}i_{\theta_3}\omega &= -L_{\theta_{w_1}}L_{\theta_3}w_2 + L_{\theta_{w_1}}L_{\theta_3}w_2 - L_{\theta_3}L_{\theta_{w_1}}w_2 \\ &= -L_{\theta_3}L_{\theta_{w_1}}w_2 = -L_{\theta_3}\{w_1, w_2\}_K \\ &= -i_{\theta_3}i_{\theta_{\{w_1, w_2\}_K}}\omega = i_{\theta_{\{w_1, w_2\}_K}}i_{\theta_3}\omega \end{aligned}$$

Finally, if we take  $\theta = [\theta_{w_1}, \theta_{w_2}] - \theta_{\{w_1, w_2\}_K}$  we have that  $i_\theta\omega$  is a closed 1-form and that  $i_\theta i_{\theta_3}\omega = -i_{\theta_3}i_\theta\omega = 0$  for all  $\theta_3$ . But then by the homotopy formula  $L_{\theta_3}i_\theta\omega = 0$  whence  $i_\theta\omega = 0$ , which finally implies that  $\theta = 0$ . This concludes the proof of :

**THEOREM 7.53.** *With notations as before, the necklace words  $\mathfrak{dR}_{rel}^0 \mathbb{C}Q^d$  is a Lie algebra for the Kontsevich bracket, and the sequence*

$$0 \longrightarrow B \longrightarrow \mathfrak{dR}_{rel}^0 \mathbb{C}Q^d \xrightarrow{\tau^{-1}d} Der_w \mathbb{C}Q^d \longrightarrow 0$$

*is an exact sequence (hence a central extension) of Lie algebras.*

This result will be crucial in the study of coadjoint orbits in the final chapter.

### References.

The formal structure on smooth affine varieties is due to M. Kapranov [34] and its extension to module varieties is due to L. Le Bruyn [54]. The semi-invariants of quivers have been obtained independently by various people, among which A. Schofield, M. Van den Bergh [72], H. Derksen, J. Weyman [20] and M. Domokos , A. Zubkov [21]. We follow here the approach of [72]. The results of section 7.3 are due to A. Schofield [69] or based on discussions with him. The results of section 7.4 are due to L. Le Bruyn [53] and is inspired by prior work of B. Westbury [80]. The results of section 7.5 are due to J. Cuntz and D. Quillen [18]. The acyclicity results of section 7.6 for the free algebra is due to M. Kontsevich [41] and in the quiver case to R. Bockland , L. Le Bruyn [8] and V. Ginzburg [25] independently as is the description of the necklace Lie algebra.



## Moduli Spaces

### 8.1. Moment maps.

In section 2.8 we have studied in some detail the *real moment map* of  $m$ -tuples of  $n \times n$  matrices. In this section we will first describe the obvious extension to representation spaces of quivers and then to prove the properties of the *real moment map* for moduli spaces of  $\theta$ -semistable representations.

We fix a quiver  $Q$  on  $k$  vertices  $\{v_1, \dots, v_k\}$  and a dimension vector  $\alpha = (a_1, \dots, a_k) \in \mathbb{N}^k$ . We take the standard Hermitian inproduct on each of the vertex spaces  $\mathbb{C}^{\oplus a_i}$  and this induces the standard *operator inner product* on every arrow-component of  $\text{rep}_\alpha Q$ . That is, for every arrow

$$\textcircled{j} \xleftarrow{a} \textcircled{i} \text{ we define } (V_a, W_a) = \text{tr}(V_a W_a^*)$$

on the component  $\text{Hom}_{\mathbb{C}}(\mathbb{C}^{\oplus a_i}, \mathbb{C}^{\oplus a_j})$  for all  $V, W \in \text{rep}_\alpha$  and where  $W_a^*$  is the *adjoint matrix*  $(\overline{w_{ji}})_{i,j}$  of  $W_a = (w_{ij})_{i,j}$ . The *Hermitian inproduct* on  $\text{rep}_\alpha Q$  is defined to be

$$(V, W) = \sum_{a \in Q_a} \text{tr}(V_a W_a^*)$$

The *maximal compact subgroup* of the basechange group  $GL(\alpha) = \prod_{i=1}^k GL_{a_i}$  is the *multiple unitary group*

$$U(\alpha) = \prod_{i=1}^k U_{a_i}$$

which preserves the Hermitian inproduct under the basechange action as subgroup of  $GL(\alpha)$ . The Lie algebra  $\text{Lie } U(\alpha)$  is the algebra of multiple skew-Hermitian matrices

$$\text{Lie } U(\alpha) = \bigoplus_{j=1}^k i\text{Herm}_{a_j} = \{ h = (h_1, \dots, h_k) \mid h_j = -h_j^* \}$$

and the induced action of  $\text{Lie } U(\alpha)$  on  $\text{rep}_\alpha Q$  is given by the rule

$$(h.V)_a = h_j V_a - V_a h_i \text{ for } \textcircled{j} \xleftarrow{a} \textcircled{i}$$

for all  $V \in \text{rep}_\alpha Q$ . This action allows us to define the *real moment map*  $\mu$  for the action of  $U(\alpha)$  on the representation space  $\text{rep}_\alpha Q$  by the assignment

$$\text{rep}_\alpha Q \xrightarrow{\mu} (i\text{Lie } U(\alpha))^* \quad V \longrightarrow (h \mapsto i(h.V, V))$$

That is, the moment map is determined by

$$\begin{aligned} (h.V, V) &= \sum_{\textcircled{j} \xleftarrow{a} \textcircled{i}} \text{tr}(h_j V_a V_a^* - V_a h_i V_a^*) \\ &= \sum_{v_i \in Q_v} \text{tr}(h_i ( \sum_{\textcircled{i} \xleftarrow{a} \textcircled{i}} V_a V_a^* - \sum_{\textcircled{i} \xrightarrow{a} \textcircled{i}} V_a^* V_a )) \end{aligned}$$

Using the nondegeneracy of the Killing form on  $Lie U(\alpha)$  we have the identification

$$\mu^{-1}(\underline{0}) = \{V \in rep_\alpha Q \mid \sum_{\textcircled{i} \xleftarrow{a} \textcircled{i}} V_a V_a^* = \sum_{\textcircled{i} \xrightarrow{a} \textcircled{i}} V_a^* V_a \ \forall v_i \in Q_v\}$$

The *real moment map*  $\mu_{\mathbb{R}}$  is then defined to be

$$rep_\alpha Q \xrightarrow{\mu_{\mathbb{R}}} Lie U(\alpha) \quad V \mapsto i[V, V^*] = i( \sum_{\textcircled{i} \xleftarrow{a} \textcircled{i}} V_a V_a^* - \sum_{\textcircled{i} \xrightarrow{a} \textcircled{i}} V_a^* V_a )_j$$

Reasoning as in section 2.8 we can prove the following moment map description of the isomorphism classes of semi-simple  $\alpha$ -dimensional representations of  $Q$ .

**THEOREM 8.1.** *There are natural one-to-one correspondences between*

- (1) *points of  $iss_\alpha Q$ , and*
- (2)  *$U(\alpha)$ -orbits in  $\mu_{\mathbb{R}}^{-1}(\underline{0})$ .*

Next, we will prove a similar result to describe the points of  $M_\alpha^{ss}(Q, \theta)$ , the moduli space of  $\theta$ -semistable  $\alpha$ -dimensional representations of  $Q$ , introduced and studied in section 6.3. Fix, an integral  $k$ -tuple  $\theta = (t_1, \dots, t_k) \in \mathbb{Z}^k$  with associated *character*

$$GL(\alpha) \xrightarrow{\chi_\theta} \mathbb{C}^* \quad g = (g_1, \dots, g_k) \mapsto \prod_{i=1}^k \det(g_i)^{t_i}$$

We have seen in section 6.3 that in order to describe  $M_\alpha^{ss}(Q, \theta)$  we consider the extended representation space  $rep_\alpha Q \oplus \mathbb{C}$ . We introduce a function  $N$  on this extended space replacing the norm in the above discussion.

$$rep_\alpha Q \oplus \mathbb{C} \xrightarrow{N} \mathbb{R}_+ \quad (V, z) \mapsto |z| e^{\frac{1}{2} \|V\|^2}$$

where  $\|V\|$  is the norm coming from the Hermitian inproduct on  $rep_\alpha Q$ . Sometimes, the function  $N$  is called the *Kähler potential* for the inproduct on  $rep_\alpha Q$ . We will investigate the properties of  $N$ .

**LEMMA 8.2.** *Let  $X$  be a closed subvariety of  $rep_\alpha Q \oplus \mathbb{C}$  disjoint from  $rep'_\alpha Q = \{(V, 0) \mid V \in rep_\alpha Q\} \hookrightarrow rep_\alpha Q \oplus \mathbb{C}$ . Then, the restriction of  $N$  to  $X$  is proper and therefore achieves its minimum.*

**PROOF.** Because  $X$  and  $rep'_\alpha Q$  are disjoint closed subvarieties of  $rep_\alpha Q \oplus \mathbb{C}$ , there is a polynomial  $f \in \mathbb{C}[rep_\alpha Q \oplus \mathbb{C}] = \mathbb{C}[rep_\alpha Q][z]$  such that  $f \mid X = 1$  and  $f \mid rep'_\alpha Q = 0$ . That is,  $X$  is contained in the hypersurface

$$\mathbb{V}(f - 1) = \mathbb{V}(zP_1(V) + \dots + z^n P_n(V) - 1) \hookrightarrow rep_\alpha Q \oplus \mathbb{C}$$

where the  $P_i \in \mathbb{C}[rep_\alpha Q]$ .

Now,  $N$  is *proper* if the inverse images  $N^{-1}([0, r])$  are compact for all  $r \in \mathbb{R}_+$ , that is, there exist constants  $r_1$  and  $r_2$  depending on  $X$  and  $r$  such that

$$N(z, V) \leq r \text{ implies } |z| \leq r_1 \text{ and } \|V\| \leq r_2.$$

We can always take  $r_1 = r$  so we only need to bound  $\|V\|$ . If  $|z| \leq r e^{-\frac{1}{2}\|V\|^2}$ , then we have that

$$|zP_1(V) + \dots + z^n P_n(V)| \leq r|P_1(V)|e^{-\frac{1}{2}\|V\|^2} + \dots + r^n|P_n(V)|e^{-\frac{n}{2}\|V\|^2}$$

Choose  $r_2$ , depending on  $r$  and  $P_i$  such that the condition

$$\|V\| > r_2 \text{ implies that } |P_i(V)| < \frac{1}{n} r^{-i} e^{\frac{i}{2}\|V\|^2} \quad \forall 1 \leq i \leq n$$

But then if  $\|V\| > r_2$ , we have  $|zP_1(V) + \dots + z^n P_n(V)| < 1$  and so  $(V, z)$  does not belong to  $X$ .  $\square$

Recall that  $GL(\alpha)$  acts on the extended representation space  $rep_\alpha Q \oplus \mathbb{C}$  via

$$g.(V, z) = (g.V, \chi_\theta^{-1}(g)z)$$

LEMMA 8.3. *Let  $\mathcal{O}$  be a  $GL(\alpha)$ -orbit in the extended representation space  $rep_\alpha Q \oplus \mathbb{C}$  which is disjoint from  $rep'_\alpha Q$ . Then, if the restriction of  $N$  to  $\mathcal{O}$  achieves its minimum, then  $\mathcal{O}$  is a closed orbit.*

PROOF. Assume that  $N$  achieves its minimum in the point  $V_z = (V, z) \in \mathcal{O}$ . If  $\mathcal{O}$  is not a closed orbit we can by the Hilbert criterium find a one-parameter subgroup  $\lambda$  of  $GL(\alpha)$  such that

$$\lim_{t \rightarrow 0} \lambda(t).V_z \notin \mathcal{O}$$

and the limit exists in  $rep_\alpha Q \oplus \mathbb{C}$ . Decompose the representation  $V = \sum_{n \in \mathbb{Z}} V_n$  into eigenspaces with respect to the one-parameter subgroup  $\lambda$ , that is,

$$\lambda(t).V = \sum_{n \in \mathbb{Z}} t^n V_n$$

Because the limit exists, we have that  $V_n = 0$  whenever  $n < 0$  and  $\theta(\lambda) \leq 0$ . Because the limit is not contained in  $\mathcal{O}$  we have that  $V_n \neq 0$  for some  $n > 0$ . Further, by conjugating  $\lambda$  if necessary we may assume that the weight space decomposition  $V = \sum_n V_n$  is orthogonal with respect to the inproduct in  $rep_\alpha Q$ .

Using these properties we then have that

$$N(\lambda(t).(V, z)) = |z| e^{\frac{1}{2}\|V_0\|^2} |t|^{-\theta(\lambda)} e^{\frac{1}{2} \sum_{n>0} |t|^n \|V_n\|^2}$$

This expression will decrease when  $t$  approaches zero, contradicting the assumption that the minimum of  $N \mid \mathcal{O}$  was achieved in  $(V, z)$ . This contradiction implies that  $\mathcal{O}$  must be a closed orbit.  $\square$

Recall from section 6.3 that an orbit  $\mathcal{O}(V, z)$  is closed and disjoint from  $rep'_\alpha Q$  for some  $z \in \mathbb{C}^*$  if and only if  $V$  is the direct sum of  $\theta$ -stable representations of  $Q$ . Recall the real moment map

$$rep_\alpha Q \xrightarrow{\mu} (iLie U(\alpha))^*$$

And consider the special real valued function  $d\chi_\theta$  on  $Lie U(\alpha)$  which is the restriction to  $Lie U(\alpha)$  of the differential of  $GL(\alpha) \xrightarrow{\chi_\theta} \mathbb{C}^*$  at the identity element (which

takes real values). In fact, for any  $m = (m_1, \dots, m_k) \in \text{Lie } GL(\alpha) = M_\alpha(\mathbb{C})$  we have that

$$d\chi_\theta(m) = \sum_{v_j \in Q_v} t_j \text{tr}(m_j) = \sum_{v_j \in Q_v} \text{tr}(m_j t_j \mathbb{1}_{a_j})$$

With these notations we have the promised extension to moduli spaces of  $\theta$ -semistable representations.

**THEOREM 8.4.** *There are natural one-to-one correspondences between*

- (1) *points of  $M_\alpha^{ss}(Q, \theta)$ , and*
- (2)  *$U(\alpha)$ -orbits in  $\mu^{-1}(d\chi_\theta)$ .*

**PROOF.** Let  $V_z = (V, z) \in \text{rep}_\alpha Q \oplus \mathbb{C}$  with  $z \neq 0$ . For any  $h = (h_1, \dots, h_k)$  in  $i\text{Lie } U(\alpha)$  we define the functions

$$\begin{aligned} m_V(h) &= \frac{d}{dt} \Big|_{t=0} \log N(e^{th} \cdot V_z) \\ &= (h \cdot V, V) - d\chi_\theta(h) \end{aligned}$$

$$\begin{aligned} m_V^{(2)}(h) &= \frac{d}{dt} \Big|_{t=0} \log N(e^{th} \cdot V_z) \\ &= 2\|h \cdot V\|^2 \end{aligned}$$

The function  $m_V$  is the zero map if and only if the restriction of  $N$  to the orbit  $\mathcal{O}(V_z)$  has a critical point at  $V_z$ . As the basechange action of  $U(\alpha)$  on the extended representation space  $\text{rep}_\alpha Q \oplus \mathbb{C}$  preserves the Kähler potential  $N$ ,  $N$  induces a function on the quotient  $\mathcal{O}(V_z)/U(\alpha)$ . The formula for  $m_V^{(2)}$  shows that this function is strictly convex (except in directions along the fibers  $\{(V, c) \mid c \in \mathbb{C}\}$  where it is linear). Hence, a critical point is a minimum and there can be at most one such critical point. From the lemmas above we have that  $N$  has a minimum on  $\mathcal{O}(V_z)$  if and only if  $\mathcal{O}(V_z)$  is a closed orbit, which in its turn is equivalent to  $V$  being the direct sum of  $\theta$ -stable representations, whence determining a point of  $M_\alpha^{ss}(Q, \theta)$ .  $\square$

Finally, for any  $h \in i\text{Lie } U(\alpha)$  we have the formulas

$$\begin{aligned} \mu(V)(h) &= i \sum_{v_i \in Q_v} \text{tr}(h_i ( \sum_{\textcircled{i} \leftarrow \textcircled{a}} V_a V_a^* - \sum_{\textcircled{a} \rightarrow \textcircled{i}} V_a^* V_a )) \\ d\chi_\theta(h) &= \sum_{v_i \in Q_v} \text{tr}(h_i t_i \mathbb{1}_{a_i}) \end{aligned}$$

whence by nondegeneracy of the Killing form, the equality  $\mu(V) = d\chi_\theta$  is equivalent to the conditions

$$\sum_{\textcircled{i} \leftarrow \textcircled{a}} V_a V_a^* - \sum_{\textcircled{a} \rightarrow \textcircled{i}} V_a^* V_a = it_j \mathbb{1}_{a_j} \quad \forall v_j \in Q_v$$

We can assign to  $\theta = (t_1, \dots, t_k) \in \mathbb{Z}^k$  the element  $i\theta \mathbb{1}_\alpha = (it_1 \mathbb{1}_{a_1}, \dots, it_k \mathbb{1}_{a_k}) \in \text{Lie } U(\alpha)$ . We then can rephrase the results of this section as

**THEOREM 8.5.** *There are natural identification between the spaces*


$$\text{iss}_\alpha Q \longleftrightarrow \mu_{\mathbb{R}}^{-1}(\mathbb{0})/U(\alpha)$$

and between the spaces

$$M_\alpha^{ss}(Q, \theta) \longleftrightarrow \mu_{\mathbb{R}}^{-1}(i\theta\mathbb{1}_\alpha)/U(\alpha)$$

### 8.2. Dynamical systems.

In this chapter we will illustrate what we have learned on the simplest *wild quiver*  $Q$  which is neither Dynkin nor extended Dynkin




In this section we will show that the representation theory of this quiver is of importance in system theory.

A *linear time invariant dynamical system*  $\Sigma$  is determined by the following system of differential equations

$$(8.1) \quad \begin{cases} \frac{dx}{dt} = Bx + Au \\ y = Cx. \end{cases}$$

Here,  $u(t) \in \mathbb{C}^m$  is the *input* or *control* of the system at time  $t$ ,  $x(t) \in \mathbb{C}^n$  the *state* of the system and  $y(t) \in \mathbb{C}^p$  the *output* of the system  $\Sigma$ . *Time invariance* of  $\Sigma$  means that the matrices  $A \in M_{n \times m}(\mathbb{C})$ ,  $B \in M_n(\mathbb{C})$  and  $C \in M_{p \times n}(\mathbb{C})$  are constant, that is  $\Sigma = (A, B, C)$  is a representation of the quiver  $\tilde{Q}$



of dimension vector  $\alpha = (m, n, p)$ . The system  $\Sigma$  can be represented as a *black box*



which is in a certain state  $x(t)$  that we can try to change by using the input controls  $u(t)$ . By reading the output signals  $y(t)$  we can try to determine the state of the system.

Recall that the *matrix exponential*  $e^B$  of any  $n \times n$  matrix  $B$  is defined by the infinite series

$$e^B = \mathbb{1}_n + B + \frac{B^2}{2!} + \dots + \frac{B^m}{m!} + \dots$$

The importance of this construction is clear from the fact that  $e^{Bt}$  is the *fundamental matrix* for the homogeneous differential equation  $\frac{dx}{dt} = Bx$ . That is, the columns of  $e^{Bt}$  are a basis for the  $n$ -dimensional space of solutions of the equation  $\frac{dx}{dt} = Bx$ .

Motivated by this, we look for a solution to equation (8.1) as the form  $x(t) = e^{Bt}g(t)$  for some function  $g(t)$ . Substitution gives the condition

$$\frac{dg}{dt} = e^{-Bt}Au \quad \text{whence} \quad g(\tau) = g(\tau_0) + \int_{\tau_0}^{\tau} e^{-Bt}Au(t)dt.$$

Observe that  $x(\tau_0) = e^{B\tau_0}g(\tau_0)$  and we obtain the solution of the linear dynamical system  $\Sigma = (A, B, C)$  :

$$\begin{cases} x(\tau) &= e^{(\tau-\tau_0)B}x(\tau_0) + \int_{\tau_0}^{\tau} e^{(\tau-t)B} Au(t)dt \\ y(\tau) &= Ce^{B(\tau-\tau_0)}x(\tau_0) + \int_{\tau_0}^{\tau} Ce^{(\tau-t)B} Au(t)dt. \end{cases}$$

Differentiating we see that this is indeed a solution and it is the unique one having a prescribed starting state  $x(\tau_0)$ . Indeed, given another solution  $x_1(\tau)$  we have that  $x_1(\tau) - x(\tau)$  is a solution to the homogeneous system  $\frac{dx}{dt} = Bx$ , but then

$$x_1(\tau) = x(\tau) + e^{\tau B}e^{-\tau_0 B}(x_1(\tau_0) - x(\tau_0)).$$

We call the system  $\Sigma$  *completely controllable* if we can steer any starting state  $x(\tau_0)$  to the zero state by some control function  $u(t)$  in a finite time span  $[\tau_0, \tau]$ . That is, the equation

$$0 = x(\tau_0) + \int_{\tau_0}^{\tau} e^{(\tau_0-t)B} Au(t)dt$$

has a solution in  $\tau$  and  $u(t)$ . As the system is time-invariant we may always assume that  $\tau_0 = 0$  and have to satisfy the equation

$$(8.2) \quad 0 = x_0 + \int_0^{\tau} e^{tB} Au(t)dt \quad \text{for every } x_0 \in \mathbb{C}^n$$

Consider the *control matrix*  $c(\Sigma)$  which is the  $n \times mn$  matrix

$$c(\Sigma) = \left[ \begin{array}{|c|} \hline A \\ \hline \end{array} \right] \left[ \begin{array}{|c|} \hline BA \\ \hline \end{array} \right] \left[ \begin{array}{|c|} \hline B^2A \\ \hline \end{array} \right] \cdots \left[ \begin{array}{|c|} \hline B^{n-1}A \\ \hline \end{array} \right]$$

Assume that  $rk\ c(\Sigma) < n$  then there is a non-zero state  $s \in \mathbb{C}^n$  such that  $s^{tr}c(\Sigma) = 0$ , where  $s^{tr}$  denotes the transpose (row column) of  $s$ . Because  $B$  satisfies the characteristic polynomial  $\chi_B(t)$ ,  $B^n$  and all higher powers  $B^m$  are linear combinations of  $\{1_n, B, B^2, \dots, B^{n-1}\}$ . Hence,  $s^{tr}B^m A = 0$  for all  $m$ . Writing out the power series expansion of  $e^{tB}$  in equation (8.2) this leads to the contradiction that  $0 = s^{tr}x_0$  for all  $x_0 \in \mathbb{C}^n$ . Hence, if  $rk\ c(\Sigma) < n$ , then  $\Sigma$  is not completely controllable.

Conversely, let  $rk\ c(\Sigma) = n$  and assume that  $\Sigma$  is not completely controllable. That is, the space of all states

$$s(\tau, u) = \int_0^{\tau} e^{-tB} Au(t)dt$$

is a proper subspace of  $\mathbb{C}^n$ . But then, there is a non-zero state  $s \in \mathbb{C}^n$  such that  $s^{tr}s(\tau, u) = 0$  for all  $\tau$  and all functions  $u(t)$ . Differentiating this with respect to  $\tau$  we obtain

$$(8.3) \quad s^{tr}e^{-\tau B} Au(\tau) = 0 \quad \text{whence} \quad s^{tr}e^{-\tau B} A = 0$$

for any  $\tau$  as  $u(\tau)$  can take on any vector. For  $\tau = 0$  this gives  $s^{tr}A = 0$ . If we differentiate (8.3) with respect to  $\tau$  we get  $s^{tr}Be^{-\tau B}A = 0$  for all  $\tau$  and for  $\tau = 0$  this gives  $s^{tr}BA = 0$ . Iterating this process we show that  $s^{tr}B^m A = 0$  for any  $m$ , whence

$$s^{tr} [A \quad BA \quad B^2A \quad \dots \quad B^{n-1}A] = 0$$

contradicting the assumption that  $rk\ c(\Sigma) = n$ . That is, we have proved :

PROPOSITION 8.6. *A linear time-invariant dynamical system  $\Sigma$  determined by the matrices  $(A, B, C)$  is completely controllable if and only if  $\text{rk } c(\Sigma)$  is maximal.*

We say that a state  $x(\tau)$  at time  $\tau$  is *unobservable* if  $Ce^{(\tau-t)B}x(\tau) = 0$  for all  $t$ . Intuitively this means that the state  $x(\tau)$  cannot be detected uniquely from the output of the system  $\Sigma$ . Again, if we differentiate this condition a number of times and evaluate at  $t = \tau$  we obtain the conditions

$$Cx(\tau) = CBx(\tau) = \dots = CB^{n-1}x(\tau) = 0.$$

We say that  $\Sigma$  is *completely observable* if the zero state is the only unobservable state at any time  $\tau$ . Consider the *observation matrix*  $o(\Sigma)$  of the system  $\Sigma$  which is the  $pn \times n$  matrix

$$o(\Sigma) = [C^{tr} \quad (CB)^{tr} \quad \dots \quad (CB^{n-1})^{tr}]^{tr}$$

An analogous argument as in the proof of proposition 8.6 gives us that a linear time-invariant dynamical system  $\Sigma$  determined by the matrices  $(A, B, C)$  is completely observable if and only if  $\text{rk } o(\Sigma)$  is maximal.

Assume we have two systems  $\Sigma$  and  $\Sigma'$ , determined by matrix triples from  $\text{rep}_\alpha Q = M_{n \times m}(\mathbb{C}) \times M_n(\mathbb{C}) \times M_{p \times n}(\mathbb{C})$  producing the same output  $y(t)$  when given the same input  $u(t)$ , for all possible input functions  $u(t)$ . We recall that the output function  $y$  for a system  $\Sigma = (A, B, C)$  is determined by

$$y(\tau) = Ce^{B(\tau-\tau_0)}x(\tau_0) + \int_{\tau_0}^{\tau} Ce^{(\tau-t)B}Au(t)dt.$$

Differentiating this a number of times and evaluating at  $\tau = \tau_0$  as in the proof of proposition 8.6 equality of input/output for  $\Sigma$  and  $\Sigma'$  gives the conditions

$$CB^i A = C' B'^i A' \quad \text{for all } i.$$

But then, we have for any  $v \in \mathbb{C}^{mn}$  that  $c(\Sigma)(v) = 0 \Leftrightarrow c(\Sigma')(v) = 0$  and we can decompose  $\mathbb{C}^{pn} = V \oplus W$  such that the restriction of  $c(\Sigma)$  and  $c(\Sigma')$  to  $V$  are the zero map and the restrictions to  $W$  give isomorphisms with  $\mathbb{C}^n$ . Hence, there is an invertible matrix  $g \in GL_n$  such that  $c(\Sigma') = gc(\Sigma)$  and from the commutative diagram

$$\begin{array}{ccccc} \mathbb{C}^{mn} & \xrightarrow{c(\Sigma)} & \mathbb{C}^n & \xrightarrow{o(\Sigma)} & \mathbb{C}^{pn} \\ & & \downarrow g & & \\ \mathbb{C}^{mn} & \xrightarrow{c(\Sigma')} & \mathbb{C}^n & \xrightarrow{o(\Sigma')} & \mathbb{C}^{pn} \end{array}$$

we obtain that also  $o(\Sigma') = o(\Sigma)g^{-1}$ .

Consider the system  $\Sigma_1 = (A_1, B_1, C_1)$  *equivalent* with  $\Sigma$  under the base-change matrix  $g$ . That is,  $\Sigma_1 = g.\Sigma = (gA, gBg^{-1}, Cg^{-1})$ . Then,

$$[A_1, B_1 A_1, \dots, B_1^{n-1} A_1] = gc(\Sigma) = c(\Sigma') = [A', B' A', \dots, B'^{n-1} A']$$

and so  $A_1 = A'$ . Further, as  $B_1^{i+1} A_1 = B'^{i+1} A'$  we have by induction on  $i$  that the restriction of  $B_1$  on the subspace of  $B'^i \text{Im}(A')$  is equal to the restriction of  $B'$  on this space. Moreover, as  $\sum_{i=0}^{n-1} B'^i \text{Im}(A') = \mathbb{C}^n$  it follows that  $B_1 = B'$ . Because  $o(\Sigma') = o(\Sigma)g^{-1}$  we also have  $C_1 = C'$ . This finishes the proof of :

PROPOSITION 8.7. Let  $\Sigma = (A, B, C)$  and  $\Sigma' = (A', B', C')$  be two completely controllable and completely observable dynamical systems. The following are equivalent

- (1) The input/output behavior of  $\Sigma$  and  $\Sigma'$  are equal.
- (2) The systems  $\Sigma$  and  $\Sigma'$  are equivalent, that is, there exists an invertible matrix  $g \in GL_n$  such that


$$A' = gA, \quad B' = gBg^{-1} \quad \text{and} \quad C' = Cg^{-1}.$$

Hence, in *system identification* it is important to classify completely controllable and observable systems  $\Sigma \in \text{rep}_\alpha \tilde{Q}$  under this restricted basechange action. We will concentrate on the input part and consider *completely controllable minisystems*, that is, representations  $\Sigma = (A, B) \in \text{rep}_\alpha Q$  where  $\alpha = (m, n)$  such that  $c(\Sigma)$  is of maximal rank. First, we relate the system theoretic notion to that of  $\theta$ -semistability for  $\theta = (-n, m)$  (observe that  $\theta(\alpha) = 0$ ).

LEMMA 8.8. If  $\Sigma = (A, B) \in \text{rep}_\alpha Q$  is  $\theta$ -semistable, then  $\Sigma$  is completely controllable and  $m \leq n$ .

PROOF. If  $m > n$  then  $(\text{Ker } A, 0)$  is a proper subrepresentation of  $\Sigma$  of dimension vector  $\beta = (\dim \text{Im } A - m, 0)$  with  $\theta(\beta) < 0$  so  $\Sigma$  cannot be  $\theta$ -semistable. If  $\Sigma$  is not completely controllable then the subspace  $W$  of  $\mathbb{C}^{\oplus n}$  spanned by the images of  $A, BA, \dots, B^{n-1}A$  has dimension  $k < n$ . But then,  $\Sigma$  has a proper subrepresentation of dimension vector  $\beta = (m, k)$  with  $\theta(\beta) < 0$ , contradicting the  $\theta$ -semistability assumption.  $\square$

We introduce a combinatorial gadget : the *Kalman code* . It is an array consisting of  $(n + 1) \times m$  boxes each having a position label  $(i, j)$  where  $0 \leq i \leq n$  and  $1 \leq j \leq m$ . These boxes are ordered *lexicographically* that is  $(i', j') < (i, j)$  if and only if either  $i' < i$  or  $i' = i$  and  $j' < j$ . Exactly  $n$  of these boxes are painted black subject to the rule that if box  $(i, j)$  is black, then so is box  $(i', j)$  for all  $i' < i$ . That is, a Kalman code looks like



We assign to a completely controllable couple  $\Sigma = (A, B)$  its Kalman code  $K(\Sigma)$  as follows : let  $A = [A_1 \ A_2 \ \dots \ A_m]$ , that is  $A_i$  is the  $i$ -th column of  $A$ . Paint the box  $(i, j)$  black if and only if the column vector  $B^i A_j$  is linearly independent of the column vectors  $B^k A_l$  for all  $(k, l) < (i, j)$ .






FIGURE 1. Kalman code and barcode.

The painted array  $K(\Sigma)$  is indeed a Kalman code. Assume that box  $(i, j)$  is black but box  $(i', j)$  white for  $i' < i$ , then

$$B^{i'} A_j = \sum_{(k,l) < (i',j)} \alpha_{kl} B^k A_l \quad \text{but then,} \quad B^i A_j = \sum_{(k,l) < (i',j)} \alpha_{kl} B^{k+i-i'} A_l$$

and all  $(k + i - i', l) < (i, l)$ , a contradiction. Moreover,  $K(\Sigma)$  has exactly  $n$  black boxes as there are  $n$  linearly independent columns of the control matrix  $c(\Sigma)$  when  $\Sigma$  is completely controllable.

The Kalman code is a discrete invariant of the orbit  $\mathcal{O}(\Sigma)$  under the restricted basechange action by  $GL_n$ . This follows from the fact that  $B^i A_j$  is linearly independent of the  $B^k A_l$  for all  $(k, l) < (i, j)$  if and only if  $gB^i A_j$  is linearly independent of the  $gB^k A_l$  for any  $g \in GL_n$  and the observation that  $gB^k A_l = (gBg^{-1})^k(gA)_l$ .

With  $rep_\alpha^c Q$  we will denote the open subset of  $rep_\alpha Q$  of all completely controllable couples  $(A, B)$ . We consider the map

$$rep_\alpha Q \xrightarrow{\psi} M_{n \times (n+1)m}(\mathbb{C})$$

$$(A, B) \mapsto [A \quad BA \quad B^2A \quad \dots \quad B^{n-1}A \quad B^nA]$$

The matrix  $\psi(A, B)$  determines a linear map  $\psi_{(A,B)} : \mathbb{C}^{(n+1)m} \rightarrow \mathbb{C}^n$  and  $(A, B)$  is a completely controllable couple if and only if the corresponding linear map  $\psi_{(A,B)}$  is surjective. Moreover, there is a linear action of  $GL_n$  on  $M_{n \times (n+1)m}(\mathbb{C})$  by left multiplication and the map  $\psi$  is  $GL_n$ -equivariant.

The Kalman code induces a *barcode* on  $\psi(A, B)$ , that is the  $n \times n$  minor of  $\psi(A, B)$  determined by the columns corresponding to black boxes in the Kalman code, see figure 1. By construction this minor is an invertible matrix  $g^{-1} \in GL_n$ . We can choose a canonical point in the orbit  $\mathcal{O}(\Sigma) : g.(A, B)$ . It does have the characteristic property that the  $n \times n$  minor of its image under  $\psi$ , determined by the Kalman code is the identity matrix  $\mathbb{1}_n$ . The matrix  $\psi(g.(A, B))$  will be denoted by  $b(A, B)$  and is called barcode of the completely controllable pair  $\Sigma = (A, B)$ . We claim that the barcode determines the orbit uniquely.

The map  $\psi$  is injective on the open set  $rep_\alpha^c Q$ . Indeed, if

$$[A \quad BA \quad \dots \quad B^nA] = [A' \quad B'A' \quad \dots \quad B'^nA']$$

then  $A = A'$ ,  $B \mid Im(A) = B' \mid Im(A)$  and hence by induction also

$$B \mid B^i Im(A) = B' \mid B'^i Im(A') \quad \text{for all } i \leq n - 1.$$

But then,  $B = B'$  as both couples  $(A, B)$  and  $(A', B')$  are completely controllable. Hence, the barcode  $b(A, B)$  determines the orbit  $\mathcal{O}(\Sigma)$  and is a point in the

Grassmannian  $Grass_n(m(n+1))$ . We have

$$\begin{array}{ccc} V_c & \xrightarrow{\psi} & M_{n \times m(n+1)}^{max}(\mathbb{C}) \\ & \searrow b(\cdot) & \downarrow \chi \\ & & Grass_n(m(n+1)) \end{array}$$

where  $\psi$  is a  $GL_n$ -equivariant embedding and  $\chi$  the orbit map. Observe that the barcode matrix  $b(A, B)$  shows that the stabilizer of  $(A, B)$  is trivial. Indeed, the minor of  $g.b(A, B)$  determined by the Kalman code is equal to  $g$ . Moreover, continuity of  $b$  implies that the orbit  $\mathcal{O}(\Sigma)$  is closed in  $rep_\alpha^c Q$ .

Consider the differential of  $\psi$ . For all  $(A, B) \in rep_\alpha^c Q$  and  $(X, Y) \in T_{(A,B)} rep_\alpha^c Q \simeq rep_\alpha^c Q$  we have

$$(B + \epsilon Y)^j (A + \epsilon X) = B^n A + \epsilon (B^n X + \sum_{i=0}^{j-1} B^i Y B^{n-1-i} A).$$

Therefore the differential of  $\psi$  in  $(A, B) \in rep_\alpha^c Q$ ,  $d\psi_{(A,B)}(X, Y)$  is equal to

$$\begin{bmatrix} X & BX + YA & B^2X + BYA + YBA & \dots & B^n X + \sum_{i=0}^{n-1} B^i Y B^{n-1-i} A \end{bmatrix}.$$

Assume  $d\psi_{(A,B)}(X, Y)$  is the zero matrix, then  $X = 0$  and substituting in the next term also  $YA = 0$ . Substituting in the third gives  $YBA = 0$ , then in the fourth  $YB^2A = 0$  and so on until  $YB^{n-1}A = 0$ . But then,

$$Y \begin{bmatrix} A & BA & B^2A & \dots & B^{n-1}A \end{bmatrix} = 0.$$

If  $(A, B)$  is a completely controllable pair, this implies that  $Y = 0$  and hence shows that  $d\psi_{(A,B)}$  is injective for all  $(A, B) \in rep_\alpha^c Q$ . By the implicit function theorem,  $\psi$  induces a  $GL_n$ -equivariant diffeomorphism between  $rep_\alpha^c Q$  and a locally closed submanifold of  $M_{n \times (n+1)m}^{max}(\mathbb{C})$ . The image of this submanifold under the orbit map  $\chi$  is again a manifold as all fibers are equal to  $GL_n$ . This concludes the difficult part of the *Kalman theorem* :

**THEOREM 8.9.** *The orbit space  $O_c = rep_\alpha^c Q/GL_n$  of equivalence classes of completely controllable couples is a locally closed submanifold of dimension  $m.n$  of the Grassmannian  $Grass_n(m(n+1))$ . In fact  $rep_\alpha^c Q \xrightarrow{b} O_c$  is a principal  $GL_n$ -bundle.*

To prove the dimension statement, consider  $rep_\alpha^c(K)$  the set of completely controllable pairs  $(A, B)$  having Kalman code  $K$  and let  $O_c(K)$  be the image under the orbit map. After identifying  $rep_\alpha^c(K)$  with its image under  $\psi$ , the barcode matrix  $b(A, B)$  gives a section  $O_c(K) \xrightarrow{s} rep_\alpha^c(K)$ . In fact,

$$GL_n \times O_c(K) \longrightarrow V_c(K) \quad (g, x) \mapsto g.s(x)$$

is a  $GL_n$ -equivariant diffeomorphism because the  $n \times n$  minor determined by  $K$  of  $g.b(A, B)$  is  $g$ . Consider the *generic* Kalman code  $K^g$  of figure 2 obtained by painting the top boxes black from left to right until one has  $n$  black boxes. Clearly  $rep_\alpha^c(K^g)$  is open in  $rep_\alpha^c$  and one deduces

$$\dim O_c = \dim O_c(K^g) = \dim V_c(K^g) - \dim GL_n = mn + n^2 - n^2 = mn.$$




FIGURE 2. Generic Kalman code.

The Kalman orbit space also naturally defines an order over the moduli space  $M_\alpha^{ss}(Q, \theta)$ . First, observe that whenever  $m \leq n$  we have  $\theta$ -stable representations of dimension vector  $\alpha = (m, n)$  for  $\theta = (-n, m)$ . Then,

$$\dim M_\alpha^{ss}(Q, \theta) = \dim \text{rep}_\alpha Q - \dim GL(\alpha) + 1 = n^2 + mn - n^2 - m^2 + 1 = m(n - m) + 1$$

By the lemma we have that  $\text{rep}_\alpha^{ss} Q$  is an open subset of  $\text{rep}_\alpha^c Q$  and let  $O_{ss}$  be the open subset of  $O_c$  it determines. Then, the natural quotient map

$$O_{ss} \longrightarrow M_\alpha^{ss}(Q, \theta)$$

is generically a principal  $PGL_m$ -fibration, so determines a central simple algebra over the function field of  $M_\alpha^{ss}(Q, \theta)$ .

In particular, if  $m = 1$  then  $O_{ss} \simeq M_\alpha^{ss}(Q, \theta)$  and both are isomorphic to  $\mathbb{A}^n$  and the orbits are parametrized by an old acquaintance, the *companion matrix* and its canonical *cyclic vector*

$$A = \begin{bmatrix} 1 \\ 0 \\ \dots \\ 0 \\ 0 \end{bmatrix} \quad B = \begin{bmatrix} 0 & & & x_n \\ -1 & 0 & & x_{n-1} \\ & \ddots & \ddots & \vdots \\ & & -1 & 0 & x_2 \\ & & & -1 & x_1 \end{bmatrix}$$

Trivial as this case seems, we will see that it soon gets interesting if we consider its extension to the double quiver  $Q^d$  and to deformed preprojective algebras.

### 8.3. Deformed preprojective algebras.

Recall the construction of *deformed preprojective algebras* given in section 5.6. Let  $Q$  be a quiver on  $k$  vertices and  $Q^d$  its *double quiver*, that is to each arrow  $a \in Q_a$  we add an arrow  $a^*$  with the reverse orientation in  $Q_a^d$  and define the *commutator element*  $c = \sum_{a \in Q_a} [a, a^*]$  in the path algebra  $\mathbb{C}Q^d$ . For a weight  $\lambda = (\lambda_1, \dots, \lambda_k) \in \mathbb{C}^k$  we define the *deformed preprojective algebra*

$$\Pi_\lambda = \frac{\mathbb{C}Q^d}{c - \lambda}$$

In this section we will give an outline of the determination of the dimension vectors of simple  $\Pi_\lambda$ -representations due to W. Crawley-Boevey [14].

We know already that a dimension vector  $\alpha = (a_1, \dots, a_k)$  can be the dimension vector of a  $\Pi_\lambda$ -representation only if  $\lambda \cdot \alpha = 0$ , so we will denote this subset of  $\mathbb{N}^k$  by  $\mathbb{N}_\lambda^k$ . With  $\Delta_\lambda^+$  we will denote the subset of *positive roots*  $\alpha$  of  $Q$  lying in  $\mathbb{N}_\lambda^k$  and with  $\mathbb{N}\Delta_\lambda^+$  the additive semigroup they generate.

If  $v_i$  is a loop-free vertex of  $Q$  we have defined the *reflection*  $\mathbb{Z}^k \xrightarrow{r_i} \mathbb{Z}^k$  by

$$r_i(\alpha) = \alpha - T_Q(\alpha, \epsilon_i)$$

and we define its *dual reflection*  $\mathbb{C}^k \xrightarrow{s_i} \mathbb{C}^k$  by the formula

$$s_i(\lambda)_j = \lambda_j - T_Q(\epsilon_i, \epsilon_j)\lambda_i$$

Clearly, we have  $s_i(\lambda) \cdot \alpha = \lambda \cdot r_i(\alpha)$ . We say that a loop-free vertex  $v_i$  in  $Q$  is *admissible* for  $(\lambda, \alpha)$  (or for  $\lambda$ ) if  $\lambda_i \neq 0$ . We define an equivalence relation  $\sim$  on pairs  $(\lambda, \alpha) \in \mathbb{C}^k \times \mathbb{Z}^k$  induced by  $(\lambda, \alpha) \sim (s_i(\lambda), r_i(\alpha))$  whenever  $v_i$  is an admissible vertex for  $(\lambda, \alpha)$ . We want to relate the representation theory of  $\Pi_\lambda$  to that of  $\Pi_{s_i(\lambda)}$ .

**THEOREM 8.10.** *If  $v_i$  is an admissible vertex for  $\lambda$ , then there is an equivalence of categories*

$$\Pi_\lambda - \text{rep} \xrightarrow{E_i} \Pi_{s_i(\lambda)} - \text{rep}$$

*that acts as the reflection  $r_i$  on the dimension vectors.*

**PROOF.** Because the definition of  $\Pi_\lambda$  does not depend on the orientation of the quiver  $Q$  we may assume that there are no arrows in  $Q$  having starting vertex  $v_i$ . Let  $V \in \text{rep}_\alpha \Pi_\lambda$  and consider  $V$  as a representation of the double quiver  $Q^d$ . Consider the vector space

$$V_\oplus = \bigoplus_{\textcircled{i} \xleftarrow{a} \textcircled{j}} V_j$$

where the sum is taken over all arrows  $a \in Q_a$  terminating in  $v_i$ . Let  $\mu_a$  and  $\pi_a$  be the inclusion and projection between  $V_j$  and  $V_\oplus$  and define maps  $V_i \xrightarrow{\mu} V_\oplus$  and  $V_\oplus \xrightarrow{\pi} V_i$  by the formulas

$$\pi = \frac{1}{\lambda_i} \sum_{\textcircled{i} \xleftarrow{a} \textcircled{j}} V_a \circ \pi_a \quad \text{and} \quad \mu = \sum_{\textcircled{i} \xleftarrow{a} \textcircled{j}} \mu_a \circ V_a$$

then  $\pi \circ \mu = \mathbb{1}_{V_i}$  whence  $\mu \circ \pi$  is an idempotent endomorphism on  $V_\oplus$ .

We define the representation  $V'$  of  $Q^d$  by the following data :  $V'_j = V_j$  for  $j \neq i$ ,  $V'_a = V_a$  and  $V'_{a^*} = V_{a^*}$  whenever the terminating vertex of  $a$  is not  $v_i$ . Further,

$$V'_i = \text{Im } \mathbb{1} - \mu \circ \pi = \text{Ker } \pi$$

and for an arrow  $\textcircled{i} \xleftarrow{a} \textcircled{j}$  in  $Q$  we define

$$\begin{cases} V'_a &= -\lambda_i(\mathbb{1} - \mu \circ \pi) \circ \mu_a : V'_j \longrightarrow V'_i \\ V'_{a^*} &= \pi_a \mid V'_i : V'_i \longrightarrow V'_j \end{cases}$$

We claim that  $V'$  is a representation of  $\Pi_{s_i(\lambda)}$ . Indeed, for a vertex  $v_i$  we have

$$\sum_{\textcircled{i} \xleftarrow{a} \textcircled{j}} V'_a V'_{a^*} = \sum_{\textcircled{i} \xleftarrow{a} \textcircled{j}} -\lambda_i(\mathbb{1} - \mu \circ \pi) \circ \mu_a \circ \pi_a \mid V'_i = -\lambda_i(\mathbb{1} - \mu \circ \pi) \mid V'_i = -\lambda_i \mathbb{1}_{V'_i}$$

and  $(s_i\lambda)_i = -\lambda_i$ . Further, for an arrow  $\textcircled{i} \xleftarrow{a} \textcircled{j}$  in  $Q$  then

$$V'_{a^*} V'_a = \pi_a \circ (-\lambda_i(\mathbb{1} - \mu \circ \pi) \circ \mu_a) = -\lambda_i \pi_a \circ \mu_a + \lambda_i \pi_a \circ \mu \circ \pi \circ \mu_a = -\lambda_i \mathbb{1}_{V_j} + V_{a^*} V_a$$

but then, whenever  $j \neq i$  we have the equality

$$\sum_{\textcircled{i} \xleftarrow{a} \textcircled{\ominus}} V'_a V'_{a^*} - \sum_{\textcircled{\ominus} \xleftarrow{a} \textcircled{j}} V'_a V'_{a^*} = \sum_{\textcircled{j} \xleftarrow{a} \textcircled{\ominus}} V_a V_{a^*} - \sum_{\textcircled{\ominus} \xleftarrow{a} \textcircled{j}} V_{a^*} V_a - T_Q(\epsilon_j, \epsilon_i) \lambda_i \mathbb{1}_{V_j}$$

because there are  $-T_Q(\epsilon_j, \epsilon_i)$  arrows from  $v_j$  to  $v_i$ . Then, this reduces to

$$\lambda_j \mathbb{1}_{V_j} - T_Q(\epsilon_j, \epsilon_i) \lambda_i \mathbb{1}_{V_i} = (s_i \lambda)_j \mathbb{1}_{V_j}$$

The assignment  $V \mapsto V'$  extends to a functor  $E_i$  and the exact sequence

$$0 \longrightarrow V'_i \longrightarrow V_{\oplus} \xrightarrow{\pi} V_i \longrightarrow 0$$

shows that it acts as  $r_i$  on the dimension vectors. Finally, the reflection also defines a functor  $E'_i : \Pi_{s_i(\lambda)} - \mathbf{rep} \longrightarrow \Pi_{\lambda} - \mathbf{rep}$  and one shows that there is a natural equivalence  $V \longrightarrow E'_i(E_i(V))$  finishing the proof.  $\square$

Recall from section 5.6 that for a fixed dimension vector  $\alpha$  we have the complex moment map

$$\mathit{rep}_{\alpha} Q^d \xrightarrow{\mu_{\alpha}} M_{\alpha} \quad \mu_{\alpha}(V)_i = \sum_{\textcircled{\ominus} \xleftarrow{a} \textcircled{i}} V_a V_{a^*} - \sum_{\textcircled{i} \xleftarrow{a} \textcircled{\ominus}} V_{a^*} V_a$$

and that we have the identification  $\mathit{rep}_{\alpha} \Pi_{\lambda} = \mu_{\alpha}^{-1}(\lambda)$ . A geometric interpretation of the proof of the foregoing theorem tells us that the schemes  $\mu_{\alpha}^{-1}(\lambda)$  and  $\mu_{r_i(\alpha)}^{-1}(s_i(\lambda))$  have the same number of irreducible components and that

$$\dim \mu_{\alpha}^{-1}(\lambda) - \alpha \cdot \alpha = \dim \mu_{r_i(\alpha)}^{-1}(s_i(\lambda)) - r_i(\alpha) \cdot r_i(\alpha)$$

see [14, lemma 1.2] for full details. The set of  $\lambda$ -Schur roots  $S_{\lambda}$  was defined to be the set of  $\alpha \in \mathbb{N}^k$  such that

$$p_Q(\alpha) \geq p_Q(\beta_1) + \dots + p_Q(\beta_r)$$

for all decompositions  $\alpha = \beta_1 + \dots + \beta_r$  with the  $\beta_i \in \Delta_{\lambda}^+$ . If we demand a proper inequality  $>$  for all decompositions we get a subset  $\Sigma_{\lambda}$  and call it the *set of  $\lambda$ -simple roots*. Recall that  $S_{\lambda}$  and hence  $\Sigma_{\lambda}$  consists of Schur roots of  $Q$ .

As in the case of Kac's theorem where one obtains the set of all roots from the subsets  $\Pi = \{\epsilon_i \mid v_i \text{ has no loops}\}$  and the *fundamental set of roots*  $F_Q = \{\alpha \in \mathbb{N}^k - \underline{0} \mid T_Q(\alpha, \epsilon_i) \leq 0 \text{ and } \mathit{supp}(\alpha) \text{ is connected}\}$ , we can use the above reflection functors  $E_i$  to reduce pairs  $(\lambda, \alpha)$  under the equivalence relation  $\sim$  to a particularly nice form, see [14, Thm. 4.8].

**THEOREM 8.11.** *If  $\alpha \in \Sigma_{\lambda}$ , then  $(\lambda, \alpha) \sim (\lambda', \alpha')$  with*

$$\begin{cases} \alpha' \in \Pi & \text{if } \alpha \text{ is a real root,} \\ \alpha' \in F_Q & \text{if } \alpha \text{ is an imaginary root.} \end{cases}$$

**PROPOSITION 8.12.** *If  $(\lambda, \alpha)$  is such that  $\alpha \in \Sigma_{\lambda}$ , then  $\mathit{rep}_{\alpha} \Pi_{\lambda} = \mu_{\alpha}^{-1}(\lambda)$  is irreducible and*

$$\dim \mu_{\alpha}^{-1}(\lambda) = \alpha \cdot \alpha - 1 + 2p_Q(\alpha)$$

*In particular,  $\mu_{\alpha}^{-1}(\lambda)$  is a complete intersection.*

PROOF. If  $\alpha \in \Sigma_\lambda$ , then we know by theorem 5.42 that

$$\dim \mu_\alpha^{-1}(\lambda) = \alpha \cdot \alpha - \chi_Q(\alpha, \alpha) + p_Q(\alpha) = \alpha \cdot \alpha - 1 + 2p_Q(\alpha)$$

as  $p_Q(\alpha) = 1 - \chi_Q(\alpha, \alpha)$ . Moreover, this number is also the relative dimension of the complex moment map  $\mu_\alpha$ . Therefore,  $\mu_\alpha^{-1}(\lambda)$  is equidimensional and we only have to prove that it is irreducible.

By theorem 8.11 and the geometric interpretation of the reflexion functor equivalence we may reduce to the case where  $\alpha$  is either a coordinate vector or lies in the fundamental region. The former case being trivial, we assume  $\alpha \in F_Q$ . Consider the projection map

$$\mu_\alpha^{-1}(\lambda) \xrightarrow{\pi} \text{rep}_\alpha Q$$

then the image of  $\pi$  is described in theorem 5.40 and any non-empty fiber  $\pi^{-1}(V) \simeq (\text{Ext}_{\mathbb{C}Q}^1(V, V))^*$  is irreducible. As in the proof of theorem 5.42 we can decompose  $\text{rep}_\alpha Q$  according to representation types in  $\text{rep}_\alpha(\tau)$ . Because  $\alpha \in \Sigma_\lambda$  we have that  $\dim \pi^{-1}(\text{rep}_\alpha(\tau)) < d = \alpha \cdot \alpha - 1 + 2p_Q(\alpha)$ . for all  $\tau \neq (1, \alpha)$ .

Because  $\alpha$  is a Schur root,  $\text{rep}_\alpha(1, \alpha)$  is an open set and  $\pi^{-1}(\text{rep}_\alpha Q - \text{rep}_\alpha(1, \alpha))$  has dimension less than  $d$ , whence it is sufficient to prove that  $\pi^{-1}(\text{rep}_\alpha(1, \alpha))$  is irreducible. Because it is an open subset of  $\mu_\alpha^{-1}(\lambda)$  it is equidimensional of dimension  $d$  and every fiber is irreducible. But, if  $X \longrightarrow Y$  is a dominant map with  $Y$  irreducible and all fibers irreducible of the same dimension, then  $X$  is irreducible, finishing the proof.  $\square$

The term  $\lambda$ -simple roots for  $\Sigma_\lambda$  is justified by the following result.

**THEOREM 8.13.** *Let  $(\lambda, \alpha)$  be such that  $\alpha \in \Sigma_\lambda$ . Then,  $\underline{\text{rep}}_\alpha \Pi_\lambda = \mu_\alpha^{-1}(\lambda)$  is a reduced and irreducible complete intersection of dimension  $d = \alpha \cdot \alpha - 1 + 2p_Q(\alpha)$  and the general element of  $\mu_\alpha^{-1}(\lambda)$  is a simple representation of  $\Pi_\lambda$ .*

*In particular,  $\text{iss}_\alpha \Pi_\lambda$  is an irreducible variety of dimension  $2p_Q(\alpha)$ .*

PROOF. We know that  $\mu_{|\alpha}^{-1}(\lambda)$  is irreducible of dimension  $d$ . By the type stratification, it is enough to prove the existence of one simple representation of dimension vector  $\alpha$ . The reflection functors being equivalences of categories, we may assume that  $\alpha$  is either in  $\Pi$  or in  $F_Q$ . Clearly, for  $\alpha$  a dimension vector, there is a simple representation, whence assume  $\alpha \in F_Q$ .

Assume there is no simple  $\alpha$ -dimensional representation of  $\Pi_\lambda$ . Because  $\underline{\text{rep}}_\alpha \Pi_\lambda$  is irreducible, there is a dimension vector  $\beta < \alpha$  and an open subset of representations containing a subrepresentation of dimension vector  $\beta$ . As the latter condition is closed, every  $\alpha$ -dimensional representation of  $\Pi_\lambda$  contains a  $\beta$ -dimensional subrepresentation.

Because  $\alpha$  is a Schur root for  $Q$ , the general  $\alpha$ -dimensional representation of  $Q$  extends to  $\Pi_\lambda$  and hence contains a subrepresentation of dimension vector  $\beta$ , that is  $\beta \xrightarrow{Q} \alpha$ . Applying the same argument to the quiver  $Q^\circ$  we also have  $\beta \xrightarrow{Q^\circ} \alpha$ .

If we now consider duals, this implies that the general  $\alpha$ -dimensional representation of  $Q$  has a subrepresentation of dimension vector  $\alpha - \beta$ . But then, by the results of section 6.2 we have  $\text{ext}(\beta, \alpha - \beta) = 0 = \text{ext}(\alpha - \beta, \beta)$  whence a general  $\alpha$ -dimensional representation of  $Q$  decomposes as a direct sum of representations of dimension  $\beta$  and  $\alpha - \beta$ , contradicting the fact that  $\alpha$  is a Schur root. Hence, there are  $\alpha$ -dimensional simple representations of  $\Pi_\lambda$ .

Let  $V$  be a simple representation in  $\mu_\alpha^{-1}(\lambda)$ , then computing differentials it follows that  $\mu_\alpha$  is smooth at  $V$ , whence  $\mu_\alpha^{-1}(\lambda)$  is generically reduced. But then, being a complete intersection, it is Cohen-Macaulay and therefore reduced.  $\square$

This finishes the proof of the *easy* part of the characterization of simple roots for  $\Pi_\lambda$  due to W. Crawley-Boevey, [14].

THEOREM 8.14. *The following are equivalent*


- (1)  $\Pi_\lambda$  has  $\alpha$ -dimensional simple representations.
- (2)  $\alpha \in \Sigma_\lambda$ .

The proof of [14] involves a lengthy case-by-case study and awaits a more transparent argument, perhaps along the lines of hyper-Kähler reduction as in section 8.5.


If  $\alpha \in \Sigma_\lambda$ , then  $\Pi_\lambda(\alpha)$  is an order in a central simple algebra over the function-field of  $\text{iss}_\alpha \Pi_\lambda$ .

### 8.4. Hilbert schemes.

In this section we will illustrate some of the foregoing results in the special case of the quiver  $Q$  coming from the study of linear dynamical systems, and its double quiver  $Q^d$



In order to avoid heavy use of stars, we denote as in the previous chapters,  $a = u$ ,  $a^* = v$ ,  $b = x$  and  $b^* = y$ , so the path algebra of the double  $Q^d$



is the algebra  $\mathbb{M}$  considered before. We fix the dimension vector  $\alpha = (1, n)$  and the character  $\theta = (-n, 1)$  and recall from section 8.2 that the moduli space  $M_\alpha^{ss}(Q, \theta) \simeq \mathbb{C}^n$ .

We say that  $u$  is a *cyclic vector* for the matrix-couple  $(X, Y) \in M_n(\mathbb{C}) \oplus M_n(\mathbb{C})$  if there is no proper subspace of  $\mathbb{C}^n$  containing  $u$  which is stable under left multiplication by  $X$  and  $Y$ .

LEMMA 8.15. *A representation  $V = (X, Y, u, v) \in \text{rep}_\alpha \mathbb{M}$  is  $\theta$ -semistable if and only if  $u$  is a cyclic vector for  $(X, Y)$ . Moreover, in this case  $V$  is even  $\theta$ -stable.*

PROOF. If there is a proper subspace of  $\mathbb{C}^n$  of dimension  $k$  containing  $u$  and stable under the multiplication with  $X$  and  $Y$  then  $V$  contains a subrepresentation of dimension  $\beta = (1, k)$  and  $\theta(\beta) < 0$ . If  $u$  is cyclic for  $(X, Y)$  then the only proper subrepresentations of  $V$  are of dimension  $(0, k)$  for some  $k$ , but for those  $\theta(\beta) > 0$  whence  $V$  is  $\theta$ -stable.  $\square$

The complex moment map  $\mu = \mu_\alpha$  for this situation is

$$\text{rep}_\alpha Q^d = M_n(\mathbb{C}) \oplus M_n(\mathbb{C}) \oplus \mathbb{C}^n \oplus \mathbb{C}^{n*} \xrightarrow{\mu} \mathbb{C} \oplus M_n(\mathbb{C})$$

$$(X, Y, u, v) \mapsto (-v.u, [Y, X] + u.v)$$

Observe that the image is contained in  $M_\alpha^0(\mathbb{C}) = \{(c, M) \mid c + \text{tr}(M) = 0\}$ . The differential  $d\mu$  in the point  $(X, Y, u, v)$  is equal to

$$d\mu_{(X, Y, u, v)}(A, B, c, d) = (-v.c - d.u, [B, X] + [Y, A] + u.d + c.v).$$

LEMMA 8.16. *The second component of the differential  $d\mu$  is surjective in  $(X, Y, u, v)$  if  $u$  is a cyclic vector for  $(X, Y)$ .*

PROOF. Consider the *nondegenerate* symmetric bilinear form  $\text{tr}(MN)$  on  $M_n(\mathbb{C})$ . With respect to this inproduct on  $M_n(\mathbb{C})$  the space orthogonal to the image of (the second component of)  $d\mu_{(X, Y, u, v)}$  is equal to

$$\{M \in M_n(\mathbb{C}) \mid \text{tr}([B, X]M + [Y, A]M + u.dM + c.vM) = 0, \forall (A, B, c, d)\}$$

Because the trace does not change under cyclic permutations and is nondegenerate we see that this space is equal to

$$\{M \in M_n(\mathbb{C}) \mid [M, X] = 0 \quad [Y, M] = 0 \quad Mu = 0 \quad \text{and} \quad vM = 0\}$$

But then, the kernel  $\ker M$  is a subspace of  $\mathbb{C}^n$  containing  $u$  and stable under left multiplication by  $X$  and  $Y$ . By the cyclicity assumption this implies that  $\ker M = \mathbb{C}^n$  or equivalently that  $M = 0$ . As  $d\mu_{(X, Y, u, v)}^\perp = 0$  and  $\text{tr}$  is nondegenerate, this implies that the differential is surjective.  $\square$

Let  $\text{rep}_\alpha^{ss} Q^d = \text{rep}_\alpha^s Q^d = \text{rep}_\alpha^s \mathbb{M}$  be the open variety of  $\theta$ -(semi)stable representations.

PROPOSITION 8.17. *For every matrix  $(c, M) \in M_\alpha^0(\mathbb{C})$  in the image of the map*

$$\text{rep}_\alpha^s \mathbb{M} \xrightarrow{\mu} M_\alpha^0(\mathbb{C})$$

*the inverse image  $\mu^{-1}(M)$  is a submanifold of  $\text{rep}_\alpha \mathbb{M}$  of dimension  $n^2 + 2n$ .*

This is a special case of theorem 5.44. Observe that for the quiver  $Q$  we have  $p_Q(m, n) = mn + 1 - m^2$ . As any decomposition of  $\alpha = (1, n)$  is of the form

$$(1, n) = (1, a_1) + (0, a_2) + \dots + (0, a_k) \quad \text{with} \quad \sum_i a_i = n$$

we have that  $p_Q(\alpha) = n \geq \sum_i p_Q(\beta_i) = a_1 + 1 + \dots + 1$  and equality only occurs for  $(1, 1) + (0, 1) + \dots + (0, 1)$ . Therefore  $\alpha \in S_{\mathbf{0}}$ .


We now turn to the description of the moduli space  $M_\alpha^{ss}(Q^d, \theta)$ . In this particular case we clearly have.

LEMMA 8.18. *For  $\alpha = (1, n)$  and  $\theta = (-n, 1)$  there is a natural one-to-one correspondence between*

- (1)  $GL(\alpha)$ -orbits in  $\text{rep}_\alpha^s \mathbb{M}$ , and
- (2)  $GL_n$ -orbits in  $\text{rep}_\alpha^s \mathbb{M}$  under the induced action.



For the investigation of the  $GL_n(\mathbb{C})$ -orbits on  $rep_\alpha^s \mathbb{M}$  we introduce a combinatorial gadget : the *Hilbert  $n$ -stair*. This is the lower triangular part of a square  $n \times n$  array of boxes



filled with *go-stones* according to the following two rules :

- each row contains exactly one stone, and
- each column contains at most one stone of each color.


For example, the set of all possible Hilbert 3-stairs is given below.




To every Hilbert stair  $\sigma$  we will associate a sequence of monomials  $W(\sigma)$  in the free noncommutative algebra  $\mathbb{C}\langle x, y \rangle$ , that is  $W(\sigma)$  is a sequence of words in  $x$  and  $y$ .

At the top of the stairs we place the identity element 1. Then, we descend the stairs according to the following rule.

- Every go-stone has a *top word*  $T$  which we may assume we have constructed before and a *side word*  $S$  and they are related as indicated below



For example, for the Hilbert 3-stairs we have the following sequences of non-commutative words



We will evaluate a Hilbert  $n$ -stair  $\sigma$  with associated sequence of non-commutative words  $W(\sigma) = \{1, w_2(x, y), \dots, w_n(x, y)\}$  on

$$rep_\alpha \mathbb{M} = M_n(\mathbb{C}) \oplus M_n(\mathbb{C}) \oplus \mathbb{C}^n \oplus \mathbb{C}^{n*}$$

For a quadruple  $(X, Y, u, v)$  we replace every occurrence of  $x$  in the word  $w_i(x, y)$  by  $X$  and every occurrence of  $y$  by  $Y$  to obtain an  $n \times n$  matrix  $w_i = w_i(X, Y) \in M_n(\mathbb{C})$

and by left multiplication on  $u$  a column vector  $w_i.v$ . The *evaluation of  $\sigma$  on  $(X, Y, u, v)$*  is the determinant of the  $n \times n$  matrix

$$\sigma(X, Y, u, v) = \det \begin{bmatrix} u & w_2.u & w_3.u & \cdots & w_n.u \end{bmatrix}$$


For a fixed Hilbert  $n$ -stair  $\sigma$  we denote with  $rep(\sigma)$  the subset of quadruples  $(X, Y, u, v)$  in  $rep_\alpha \mathbb{M}$  such that the evaluation  $\sigma(v, X, Y) \neq 0$ .

**THEOREM 8.19.** *For every Hilbert  $n$ -stair,  $rep(\sigma) \neq \emptyset$*

**PROOF.** Let  $u$  be the basic column vector

$$e_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

Let every black stone in the Hilbert stair  $\sigma$  fix a column of  $X$  by the rule



That is, one replaces every black stone in  $\sigma$  by 1 at the same spot in  $X$  and fills the remaining spots in the same column by zeroes. The same rule applies to  $Y$  for white stones. We say that such a quadruple  $(X, Y, u, v)$  is in  $\sigma$ -*standard form*.

With these conventions one easily verifies by induction that

$$w_i(X, Y)e_1 = e_i \quad \text{for all } 2 \leq i \leq n.$$

Hence, filling up the remaining spots in  $X$  and  $Y$  arbitrarily one has that  $\sigma(X, Y, u, v) \neq 0$  proving the claim.  $\square$

Hence,  $rep(\sigma)$  is an open subset of  $rep_\alpha \mathbb{M}$  (and even of  $rep_\alpha^s \mathbb{M}$ ) for every Hilbert  $n$ -stair  $\sigma$ . Further, for every word (monomial)  $w(x, y)$  and every  $g \in GL_n(\mathbb{C})$  we have that

$$w(gXg^{-1}, gYg^{-1})gv = gw(X, Y)v$$

and therefore the open sets  $rep(\sigma)$  are stable under the  $GL_n(\mathbb{C})$ -action on  $rep_\alpha \mathbb{M}$ . We will give representatives of the orbits in  $rep(\sigma)$ .

Let  $W_n = \{1, x, \dots, x^n, xy, \dots, y^n\}$  be the set of all words in the non-commuting variables  $x$  and  $y$  of length  $\leq n$ , ordered lexicographically.

For every quadruple  $(X, Y, u, v) \in rep_\alpha \mathbb{M}$  consider the  $n \times m$  matrix

$$\psi(X, Y, u, v) = [u \quad Xu \quad X^2u \quad \dots \quad Y^nu]$$

where  $m = 2^{n+1} - 1$  and the  $j$ -th column is the column vector  $w(X, Y)v$  with  $w(x, y)$  the  $j$ -th word in  $W_n$ .

Hence,  $(X, Y, u, v) \in \text{rep}(\sigma)$  if and only if the  $n \times n$  minor of  $\psi(X, Y, u, v)$  determined by the word-sequence  $\{1, w_2, \dots, w_n\}$  of  $\sigma$  is invertible. Moreover, as

$$\psi(gXg^{-1}, gYg^{-1}, gu, vg^{-1}) = g\psi(v, X, Y)$$

we deduce that the  $GL_n(\mathbb{C})$ -orbit of  $(X, Y, u, v) \in \text{rep}_\alpha \mathbb{M}$  contains a *unique* quadruple  $(X_1, Y_1, u_1, v_1)$  such that the corresponding minor of  $\psi(X_1, Y_1, u_1, v_1) = \mathbb{1}_n$ .

Hence, each  $GL_n(\mathbb{C})$ -orbit in  $\text{rep}(\sigma)$  contains a unique representant in  $\sigma$ -standard form. Therefore,

PROPOSITION 8.20. *The action of  $GL_n(\mathbb{C})$  on  $\text{rep}(\sigma)$  is free and the orbit space*

$$\text{rep}(\sigma)/GL_n(\mathbb{C})$$

*is an affine space of dimension  $n^2 + 2n$ .*

PROOF. The dimension is equal to the number of non-forced entries in  $X, Y$  and  $v$ . As we fixed  $n - 1$  columns in  $X$  or  $Y$  this dimension is equal to

$$k = 2n^2 - (n - 1)n + n = n^2 + 2n.$$

The argument above shows that every  $GL_n(\mathbb{C})$ -orbit contains a unique quadruple in  $\sigma$ -standard form so the orbit space is an affine space.  $\square$

THEOREM 8.21. *For  $\alpha = (1, n)$  and  $\theta = (-n, 1)$ , the moduli space*

$$M_\alpha^{ss}(Q^d, \theta) = M_\alpha^{ss}(\mathbb{M}, \theta)$$

*is a complex manifold of dimension  $n^2 + 2n$  and is covered by the affine spaces  $\text{rep}(\sigma)$ .*

PROOF. Recall that  $\text{rep}_\alpha^s \mathbb{M}$  is the open submanifold consisting of quadruples  $(x, Y, u, v)$  such that  $u$  is a cyclic vector of  $(X, Y)$  or equivalently such that

$$\mathbb{C}\langle X, Y \rangle u = \mathbb{C}^n$$

where  $\mathbb{C}\langle X, Y \rangle$  is the not necessarily commutative subalgebra of  $M_n(\mathbb{C})$  generated by the matrices  $X$  and  $Y$ .

Hence, clearly  $\text{rep}(\sigma) \subset \text{rep}_n \mathbb{M}$  for any Hilbert  $n$ -stair  $\sigma$ . Conversely, we claim that a quadruple  $(X, Y, u, v) \in \text{rep}_\alpha^s \mathbb{M}$  belongs to at least one of the open subsets  $\text{rep}(\sigma)$ .

Indeed, either  $Xu \notin \mathbb{C}u$  or  $Yu \notin \mathbb{C}u$  as otherwise the subspace  $W = \mathbb{C}u$  would contradict the cyclicity assumption. Fill the top box of the stairs with the corresponding stone and define the 2-dimensional subspace  $V_2 = \mathbb{C}u_1 + \mathbb{C}u_2$  where  $u_1 = u$  and  $u_2 = w_2(X, Y)u$  with  $w_2$  the corresponding word (either  $x$  or  $y$ ).

Assume by induction we have been able to fill the first  $i$  rows of the stairs with stones leading to the sequence of words  $\{1, w_2(x, y), \dots, w_i(x, y)\}$  such that the subspace  $V_i = \mathbb{C}u_1 + \dots + \mathbb{C}u_i$  with  $u_i = w_i(X, Y)v$  has dimension  $i$ .

Then, either  $Xu_j \notin V_i$  for some  $j$  or  $Yu_j \notin V_i$  (if not,  $V_i$  would contradict cyclicity). Then, fill the  $j$ -th box in the  $i + 1$ -th row of the stairs with the corresponding stone. Then, the top  $i + 1$  rows of the stairs form a Hilbert  $i + 1$ -stair as there can be no stone of the same color lying in the same column. Define  $w_{i+1}(x, y) = xw_i(x, y)$  (or  $yw_i(x, y)$ ) and  $u_{i+1} = w_{i+1}(X, Y)u$ . Then,  $V_{i+1} = \mathbb{C}u_1 + \dots + \mathbb{C}u_{i+1}$  has dimension  $i + 1$ .

Continuing we end up with a Hilbert  $n$ -stair  $\sigma$  such that  $(X, Y, u, v) \in \text{rep}(\sigma)$ . This concludes the proof.  $\square$

EXAMPLE 8.22. The moduli space  $M_\alpha^{ss}(Q^d, \theta)$  when  $n = 3$ .

Representatives for the  $GL_3(\mathbb{C})$ -orbits in  $\text{rep}(\sigma)$  are given by the following quadruples for  $\sigma$  a Hilbert 3-stair :

$X$	$\begin{bmatrix} 0 & a & b \\ 1 & c & d \\ 0 & e & f \end{bmatrix}$	$\begin{bmatrix} 0 & 0 & a \\ 1 & 0 & b \\ 0 & 1 & c \end{bmatrix}$	$\begin{bmatrix} 0 & a & b \\ 1 & c & d \\ 0 & e & f \end{bmatrix}$	$\begin{bmatrix} 0 & a & b \\ 0 & c & d \\ 1 & e & f \end{bmatrix}$	$\begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$	$\begin{bmatrix} a & 0 & b \\ c & 0 & d \\ e & 1 & f \end{bmatrix}$
$Y$	$\begin{bmatrix} 0 & g & h \\ 0 & i & j \\ 1 & k & l \end{bmatrix}$	$\begin{bmatrix} d & e & f \\ g & h & i \\ j & k & l \end{bmatrix}$	$\begin{bmatrix} g & 0 & h \\ i & 0 & j \\ k & 1 & l \end{bmatrix}$	$\begin{bmatrix} 0 & g & h \\ 1 & i & j \\ 0 & k & l \end{bmatrix}$	$\begin{bmatrix} 0 & 0 & j \\ 1 & 0 & k \\ 0 & 1 & l \end{bmatrix}$	$\begin{bmatrix} 0 & g & h \\ 1 & i & j \\ 0 & k & l \end{bmatrix}$
$u$	$\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$	$\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$	$\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$	$\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$	$\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$	$\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$
$v$	$[m \ n \ o]$	$[m \ n \ o]$	$[m \ n \ o]$	$[m \ n \ o]$	$[m \ n \ o]$	$[m \ n \ o]$

We now turn to the deformed preprojective algebras. Let  $\lambda = (-n\lambda, \lambda \uparrow_n) \in M_\alpha^0(\mathbb{C})$  for  $\lambda \in \mathbb{C}$ . Then,

$$\Pi_\lambda = \frac{\mathbb{M}}{(v.u + \lambda v_1, [Y, X] + u.v - \lambda v_2)}$$

then if we denote by  $M_\alpha^{ss}(\Pi_\lambda, \theta)$  the moduli space of  $\theta$ -semistable representations of  $\Pi_\lambda$ , then we have the following situation

$$\begin{array}{ccc} \mu^{-1}(\lambda) \cap \text{rep}_\alpha^s \mathbb{M} & \hookrightarrow & \text{rep}_\alpha^s \mathbb{M} \\ \downarrow & & \downarrow \\ M_\alpha^{ss}(\Pi_\lambda, \theta) & \hookrightarrow & M_\alpha^{ss}(\mathbb{M}, \theta) \end{array}$$

and from the theorem above we obtain :

THEOREM 8.23. For a  $\lambda \in M_\alpha^0(\mathbb{C})$ , the orbit space of  $\theta$ -semistable representations of the deformed preprojective algebra

$$M_\alpha^{ss}(\Pi_\lambda, \theta)$$

is a submanifold of  $M_\alpha^{ss}(\mathbb{M}, \theta)$  of dimension  $2n$ .

We will identify the special case of the preprojective algebra (that is  $\lambda = \underline{0}$  with the Hilbert scheme of  $n$  points in the plane .

Consider a codimension  $n$  ideal  $\mathfrak{i} \triangleleft \mathbb{C}[x, y]$  and fix a basis  $\{v_1, \dots, v_n\}$  of the quotient space

$$V_{\mathfrak{i}} = \frac{\mathbb{C}[x, y]}{\mathfrak{i}} = \mathbb{C}v_1 + \dots + \mathbb{C}v_n.$$

Multiplication by  $x$  on  $\mathbb{C}[x, y]$  induces a linear operator on the quotient  $V_{\mathfrak{i}}$  and hence determines a matrix  $X_{\mathfrak{i}} \in M_n(\mathbb{C})$  with respect to the chosen basis  $\{v_1, \dots, v_n\}$ . Similarly, multiplication by  $y$  determines a matrix  $Y_{\mathfrak{i}} \in M_n(\mathbb{C})$ .

Moreover, the image of the unit element  $1 \in \mathbb{C}[x, y]$  in  $V_i$  determines with respect to the basis  $\{v_1, \dots, v_n\}$  a column vector  $u \in \mathbb{C}^n = V_i$ . Clearly, this vector and matrices satisfy :

$$[X_i, Y_i] = 0 \quad \text{and} \quad \mathbb{C}[X_i, Y_i]u = \mathbb{C}^n.$$

Here,  $\mathbb{C}[X_i, Y_i]$  is the  $n$ -dimensional subalgebra of  $M_n(\mathbb{C})$  generated by the two matrices  $X_i$  and  $Y_i$ . In particular,  $u$  is a cyclic vector for the matrix-couple  $(X, Y)$ .

Conversely, if  $(X, Y, u) \in M_n(\mathbb{C}) \oplus M_n(\mathbb{C}) \oplus \mathbb{C}^n$  is a cyclic triple such that  $[X, Y] = 0$ , then  $\mathbb{C}\langle X, Y \rangle = \mathbb{C}[X, Y]$  is an  $n$ -dimensional commutative subalgebra of  $M_n(\mathbb{C})$ , then the kernel of the natural epimorphism

$$\mathbb{C}[x, y] \longrightarrow \mathbb{C}[X, Y] \quad x \mapsto X \quad y \mapsto Y$$

is a codimension  $n$  ideal  $\mathfrak{i}$  of  $\mathbb{C}[x, y]$ .

However, there is some redundancy in the assignment  $\mathfrak{i} \longrightarrow (X_i, Y_i, u_i)$  as it depends on the choice of basis of  $V_i$ . If we choose a different basis  $\{v'_1, \dots, v'_n\}$  with basechange matrix  $g \in GL_n(\mathbb{C})$ , then the corresponding triple is

$$(X'_i, Y'_i, u'_i) = (g.X_i.g^{-1}, g.Y_i.g^{-1}, gu_i)$$

The above discussion shows that there is a one-to-one correspondence between

- codimension  $n$  ideals  $\mathfrak{i}$  of  $\mathbb{C}[x, y]$ , and
- $GL_n(\mathbb{C})$ -orbits of cyclic triples  $(X, Y, u)$  in  $M_n(\mathbb{C}) \oplus M_n(\mathbb{C}) \oplus \mathbb{C}^n$  such that  $[X, Y] = 0$ .

EXAMPLE 8.24. The Hilbert scheme  $Hilb_2$ .

Consider a triple  $(X, Y, u) \in M_2(\mathbb{C}) \oplus M_2(\mathbb{C}) \oplus \mathbb{C}^2$  and assume that either  $X$  or  $Y$  has distinct eigenvalues (type a). As

$$\left[ \begin{bmatrix} \nu_1 & 0 \\ 0 & \nu_2 \end{bmatrix}, \begin{bmatrix} a & b \\ c & d \end{bmatrix} \right] = \begin{bmatrix} 0 & (\nu_1 - \nu_2)b \\ (\nu_2 - \nu_1)c & 0 \end{bmatrix}$$

we have a representant in the orbit of the form

$$\left( \begin{bmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{bmatrix}, \begin{bmatrix} \mu_1 & 0 \\ 0 & \mu_2 \end{bmatrix}, \begin{bmatrix} u_1 \\ u_2 \end{bmatrix} \right)$$

where cyclicity of the column vector implies that  $u_1 u_2 \neq 0$ .

The stabilizer subgroup of the matrix-pair is the group of diagonal matrices  $\mathbb{C}^* \times \mathbb{C}^* \hookrightarrow GL_2(\mathbb{C})$ , hence the orbit has a unique representant of the form

$$\left( \begin{bmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{bmatrix}, \begin{bmatrix} \mu_1 & 0 \\ 0 & \mu_2 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right)$$

The corresponding ideal  $\mathfrak{i} \triangleleft \mathbb{C}[x, y]$  is then

$$\mathfrak{i} = \{f(x, y) \in \mathbb{C}[x, y] \mid f(\lambda_1, \mu_1) = 0 = f(\lambda_2, \mu_2)\}$$

hence these orbits correspond to sets of two *distinct* points in  $\mathbb{C}^2$ .

The situation is slightly more complicated when  $X$  and  $Y$  have only one eigenvalue (type b). If  $(X, Y, u)$  is a cyclic commuting triple, then either  $X$  or  $Y$  is not diagonalizable. But then, as

$$\left[ \begin{bmatrix} \nu & 1 \\ 0 & \nu \end{bmatrix}, \begin{bmatrix} a & b \\ c & d \end{bmatrix} \right] = \begin{bmatrix} c & d - a \\ 0 & c \end{bmatrix}$$

we have a representant in the orbit of the form

$$\left( \begin{bmatrix} \lambda & \alpha \\ 0 & \lambda \end{bmatrix}, \begin{bmatrix} \mu & \beta \\ 0 & \mu \end{bmatrix}, \begin{bmatrix} u_1 \\ u_2 \end{bmatrix} \right)$$

with  $[\alpha : \beta] \in \mathbb{P}^1$  and  $u_2 \neq 0$ . The stabilizer of the matrixpair is the subgroup

$$\left\{ \begin{bmatrix} c & d \\ 0 & c \end{bmatrix} \mid c \neq 0 \right\} \hookrightarrow GL_2(\mathbb{C})$$

and hence we have a unique representant of the form

$$\left( \begin{bmatrix} \lambda & \alpha \\ 0 & \lambda \end{bmatrix}, \begin{bmatrix} \mu & \beta \\ 0 & \mu \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right)$$

The corresponding ideal  $i \triangleleft \mathbb{C}[x, y]$  is


$$i = \{f(x, y) \in \mathbb{C}[x, y] \mid f(\lambda, \mu) = 0 \text{ and } \alpha \frac{\partial f}{\partial x}(\lambda, \mu) + \beta \frac{\partial f}{\partial y}(\lambda, \mu) = 0\}$$

as one proves by verification on monomials because

$$\begin{bmatrix} \lambda & \alpha \\ 0 & \lambda \end{bmatrix}^k \begin{bmatrix} \mu & \beta \\ 0 & \mu \end{bmatrix}^l \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} k\alpha\lambda^{k-1}\mu^l + l\beta\lambda^k\mu^{l-1} \\ \lambda^k\mu^l \end{bmatrix}$$

Therefore,  $i$  corresponds to the set of two points at  $(\lambda, \mu) \in \mathbb{C}^2$  infinitesimally attached to each other in the direction  $\alpha \frac{\partial}{\partial x} + \beta \frac{\partial}{\partial y}$ . For each point in  $\mathbb{C}^2$  there is a  $\mathbb{P}^1$  family of such *fat points*.

Thus, points of  $Hilb_2$  correspond to either of the following two situations :



The Hilbert-Chow map  $Hilb_2 \xrightarrow{\pi} S^2 \mathbb{C}^2$  (where  $S^2 \mathbb{C}^2$  is the symmetric power of  $\mathbb{C}^2$ , that is  $S_2 = \mathbb{Z}/2\mathbb{Z}$  orbits of couples of points from  $\mathbb{C}^2$ ) sends a point of type a to the formal sum  $[p] + [p']$  and a point of type b to  $2[p]$ . Over the complement of (the image of) the diagonal, this map is a one-to-one correspondence.

However, over points on the diagonal the fibers are  $\mathbb{P}^1$  corresponding to the directions in which two points can approach each other in  $\mathbb{C}^2$ . As a matter of fact, the symmetric power  $S^2 \mathbb{C}^2$  has singularities and the Hilbert-Chow map  $Hilb_2 \xrightarrow{\pi} S^2 \mathbb{C}^2$  is a *resolution of singularities*.

**THEOREM 8.25.** *Let  $rep_\alpha \mathbb{M} \xrightarrow{\mu} M_\alpha^0(\mathbb{C})$  be the complex moment map, then*

$$Hilb_n \simeq M_\alpha^{ss}(\Pi_0, \theta)$$

*and is therefore a complex manifold of dimension  $2n$ .*

**PROOF.** We identify the triples  $(X, Y, u) \in M_n(\mathbb{C}) \oplus M_n(\mathbb{C}) \oplus \mathbb{C}^n$  such that  $u$  is a cyclic vector of  $(X, Y)$  and  $[X, Y] = 0$  with the subspace

$$\{(X, Y, u, \underline{0}) \mid [X, Y] = 0 \text{ and } u \text{ is cyclic}\} \hookrightarrow rep_\alpha^s \mathbb{M}$$

which is clearly contained in  $\mu^{-1}(0)$ . To prove the converse inclusion assume that  $(X, Y, u, v)$  is a cyclic quadruple such that

$$[X, Y] + uv = 0.$$

Let  $m(x, y)$  be any word in the noncommuting variables  $x$  and  $y$ . We claim that

$$v.m(X, Y).u = 0.$$

We will prove this by induction on the length  $l(m)$  of the word  $m(x, y)$ . When  $l(m) = 0$  then  $l(x, y) = 1$  and we have

$$v.l(X, Y).u = v.u = tr(u.v) = tr([X, Y]) = 0.$$

Assume we proved the claim for all words of length  $< l$  and take a word of the form  $m(x, y) = m_1(x, y)yxm_2(x, y)$  with  $l(m_1) + l(m_2) + 2 = l$ . Then, we have

$$\begin{aligned} wm(X, Y) &= wm_1(X, Y)YXm_2(X, Y) \\ &= wm_1(X, Y)([Y, X] + XY)m_2(X, Y) \\ &= (wm_1(X, Y)v).wm_2(X, Y) + wm_1(X, Y)XYm_2(X, Y) \\ &= wm_1(X, Y)XYm_2(X, Y) \end{aligned}$$

where we used the induction hypotheses in the last equality (the bracketed term vanishes).

Hence we can reorder the terms in  $m(x, y)$  if necessary and have that  $wm(X, Y) = wX^{l_1}Y^{l_2}$  with  $l_1 + l_2 = l$  and  $l_1$  the number of occurrences of  $x$  in  $m(x, y)$ . Hence, we have to prove the claim for  $X^{l_1}Y^{l_2}$ .

$$\begin{aligned} wX^{l_1}Y^{l_2}v &= tr(X^{l_1}Y^{l_2}vw) \\ &= -tr(X^{l_1}Y^{l_2}[X, Y]) \\ &= -tr([X^{l_1}Y^{l_2}, X]Y) \\ &= -tr(X^{l_1}[Y^{l_2}, X]Y) \\ &= -\sum_{i=0}^{l_2-1} tr(X^{l_1}Y^i[Y, X]Y^{l_2-i}) \\ &= -\sum_{i=0}^{l_2-1} tr(Y^{l_2-i}X^{l_1}Y^i[Y, X]) \\ &= -\sum_{i=0}^{l_2-1} tr(Y^{l_2-i}X^{l_1}Y^i v.w) \\ &= -\sum_{i=0}^{l_2-1} wY^{l_2-i}X^{l_1}Y^i v \end{aligned}$$

But we have seen that  $wY^{l_2-i}X^{l_1}Y^i = wX^{l_1}Y^{l_2}$  hence the above implies that  $wX^{l_1}Y^{l_2}v = -l_2wX^{l_1}Y^{l_2}v$ . But then  $wX^{l_1}Y^{l_2}v = 0$ , proving the claim.

Consequently,  $w.C(X, Y).v = 0$  and by the cyclicity condition we have  $w.C^n = 0$  hence  $w = 0$ . Finally, as  $v.w + [X, Y] = 0$  this implies that  $[X, Y] = 0$  and we can identify the fiber  $\mu^{-1}(0)$  with the indicated subspace. From this the result follows.  $\square$

We can use the affine covering of  $M_{\alpha}^{ss}(\mathbb{M}, \theta)$  by Hilbert stairs, to cover the Hilbert scheme  $Hilb_n$  by the intersections  $Hilb(\sigma) = rep(\sigma) \cap Hilb_n$ .

EXAMPLE 8.26. The Hilbert scheme  $Hilb_2$ .

Consider  $Hilb_2(\blacksquare)$ . Because

$$\begin{bmatrix} 0 & a \\ 1 & b \end{bmatrix}, \begin{bmatrix} c & d \\ e & f \end{bmatrix} = \begin{bmatrix} ae - d & af - ac - bd \\ c + be - f & d - ae \end{bmatrix}$$


this subset can be identified with  $\mathbb{C}^4$  using the equalities

$$d = ar \quad \text{and} \quad f = c + be.$$

Similarly,  $Hilb_2(\square) \simeq \mathbb{C}^4$ .

EXAMPLE 8.27. The Hilbert scheme  $Hilb_3$ .

Up to change of colors there are three 3-stairs to consider



We claim that

$$Hilb_3(\begin{smallmatrix} \bullet \\ \square \\ \square \end{smallmatrix}) \simeq \mathbb{C}^6.$$

For consider the commutator

$$\left[ \begin{bmatrix} 0 & a & b \\ 1 & c & d \\ 0 & e & f \end{bmatrix}, \begin{bmatrix} 0 & g & h \\ 0 & i & j \\ 1 & k & k \end{bmatrix} \right] = \begin{bmatrix} b-g & ai+bk-cg-eh & aj+bl-dg-fh \\ d-i & g+dk-ej & h+cj+dl-di-fj \\ f-k & -a-ck-el+ei+fk & -b-dk+ej \end{bmatrix}$$

Taking the Groebner basis of these relations one finds the following relations

$$\begin{cases} f = k \\ g = ej - ik \\ d = i \\ h = i^2 - cj + jk - il \\ b = g \\ a = ei - ck + k^2 - el \end{cases}$$

from which the claim follows. In a similar manner one proves that

$$\text{Hilb}_3 \left( \begin{array}{c|c} \bullet & \\ \hline & \bullet \end{array} \right) \simeq \mathbb{C}^6.$$

However, the situation for

$$\text{Hilb}_3 \left( \begin{array}{c|c} \bullet & \\ \hline & \circ \end{array} \right)$$

is more complicated.

**THEOREM 8.28.** *The Hilbert scheme  $\text{Hilb}_n$  of  $n$  points in  $\mathbb{C}^2$  is a complex connected manifold of dimension  $2n$ .*

**PROOF.** The symmetric power  $S^n \mathbb{C}^1$  parametrizes *sets* of  $n$ -points on the line  $\mathbb{C}^1$  and can be identified with  $\mathbb{C}^n$ . Consider the map

$$\text{Hilb}_n \xrightarrow{\pi} S^n \mathbb{C}^1$$

defined by mapping a cyclic triple  $(X, Y, u)$  with  $[X, Y] = 0$  in the orbit corresponding to the point of  $\text{Hilb}_n$  to the set  $\{\lambda_1, \dots, \lambda_n\}$  of eigenvalues of  $X$ . Observe that this map does not depend on the point chosen in the orbit.

Let  $\Delta$  be the *big diagonal* in  $S^n \mathbb{C}^1$ , that is,  $S^n \mathbb{C}^1 - \Delta$  is the space of all sets of  $n$  distinct points from  $\mathbb{C}^1$ . Clearly,  $S^n \mathbb{C}^1 - \Delta$  is a connected  $n$ -dimensional manifold. We claim that

$$\pi^{-1}(S^n \mathbb{C}^1 - \Delta) \simeq (S^n \mathbb{C}^1 - \Delta) \times \mathbb{C}^n$$

and hence is connected.

Indeed, take a matrix  $X$  with  $n$  distinct eigenvalues  $\{\lambda_1, \dots, \lambda_n\}$ . We may diagonalize  $X$ . But then, as

$$\left[ \begin{array}{ccc} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{array} \right], \left[ \begin{array}{ccc} y_{11} & \cdots & y_{1n} \\ \vdots & & \vdots \\ y_{n1} & \cdots & y_{nn} \end{array} \right] = \left[ \begin{array}{ccc} (\lambda_1 - \lambda_1)y_{11} & \cdots & (\lambda_1 - \lambda_n)y_{1n} \\ \vdots & & \vdots \\ (\lambda_n - \lambda_1)y_{n1} & \cdots & (\lambda_n - \lambda_n)y_{nn} \end{array} \right]$$

we see that also  $Y$  must be a diagonal matrix with entries  $(\mu_1, \dots, \mu_n) \in \mathbb{C}^n$  where  $\mu_i = y_{ii}$ . But then the cyclicity condition implies that all coordinates of  $v$  must be non-zero.

Now, the stabilizer subgroup of the commuting (diagonal) matrix-pair  $(X, Y)$  is the *maximal torus*  $T_n = \mathbb{C}^* \times \dots \times \mathbb{C}^*$  of diagonal invertible  $n \times n$  matrices. Using its action we may assume that all coordinates of  $v$  are equal to 1. That is,



the points in  $\pi^{-1}(\{\lambda_1, \dots, \lambda_n\})$  with  $\lambda_i \neq \lambda_j$  have unique (up to permutation as before) representatives of the form

$$\left( \begin{bmatrix} \lambda_1 & & & \\ & \lambda_2 & & \\ & & \ddots & \\ & & & \lambda_n \end{bmatrix}, \begin{bmatrix} \mu_1 & & & \\ & \mu_2 & & \\ & & \ddots & \\ & & & \mu_n \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{bmatrix} \right)$$

that is  $\pi^{-1}(\{\lambda_1, \dots, \lambda_n\})$  can be identified with  $\mathbb{C}^n$ , proving the claim.

Next, we claim that *all* the fibers of  $\pi$  have dimension at most  $n$ . Let  $\{\lambda_1, \dots, \lambda_n\} \in S^n \mathbb{C}^1$  then there are only finitely many  $X$  in Jordan normalform with eigenvalues  $\{\lambda_1, \dots, \lambda_n\}$ . Fix such an  $X$ , then the subset  $T(X)$  of cyclic triples  $(X, Y, u)$  with  $[X, Y] = 0$  has dimension at most  $n + \dim C(X)$  where  $C(X)$  is the centralizer of  $X$  in  $M_n(\mathbb{C})$ , that is,

$$C(X) = \{Y \in M_n(\mathbb{C}) \mid XY = YX\}.$$

The stabilizer subgroup  $Stab(X) = \{g \in GL_n(\mathbb{C}) \mid gXg^{-1} = X\}$  is an open subset of the vectorspace  $C(X)$  and acts freely on the subset  $T(X)$  because the action of  $GL_n(\mathbb{C})$  on  $\mu^{-1}(0) \cap rep_\alpha^s \mathbb{M}$  has trivial stabilizers.

But then, the orbit space for the  $Stab(X)$ -action on  $T(X)$  has dimension at most

$$n + \dim C(X) - \dim Stab(X) = n.$$

As we only have to consider finitely many  $X$  this proves the claim. The diagonal  $\Delta$  has dimension  $n-1$  in  $S^n \mathbb{C}^1$  and hence by the foregoing we know that the dimension of  $\pi^{-1}(\Delta)$  is at most  $2n-1$ . Let  $H$  be the connected component of  $Hilb_n$  containing the connected subset  $\pi^{-1}(S^n \mathbb{C}^1 - \Delta)$ . If  $\pi^{-1}(\Delta)$  were not entirely contained in  $H$ , then  $Hilb_n$  would have a component of dimension less than  $2n$ , which we proved not to be the case. This finishes the proof.  $\square$

We can give a representation theoretic interpretation of the *resolution of singularities Hilbert-Chow morphism*

$$Hilb_n \xrightarrow{\pi} S^n \mathbb{C}^2$$

$\Sigma_0 = \{(1, 0), (0, 1)\}$ , that is the only simple  $\Pi_0$ -representations are one-dimensional. Any semi-simple representation of  $\Pi_0$  of dimension vector  $\alpha = (1, n)$  therefore decomposes as  $T_0 \oplus S_1^{\oplus e_1} \oplus \dots \oplus S_r^{\oplus e_r}$  with  $T_0$  the unique simple  $(1, 0)$ -dimensional representation and the  $S_i$  in the two-dimensional family of  $(0, 1)$ -simple representations of  $\Pi_0$  (corresponding to couples  $(\lambda_i, \mu_i) \in \mathbb{C}^2$ ). Therefore we have the projective bundle morphism

$$Hilb_n = M_\alpha^{ss}(\Pi_0, \theta) \xrightarrow{\pi'} iss_\alpha \Pi_0 = S^n \mathbb{C}^2$$

where the mapping sends a point of  $Hilb_n$  determined by a cyclic triple  $(X, Y, u)$  to the  $n$ -tuple of eigenvalues  $(\lambda_i, \mu_i)$  of  $X$  and  $Y$ .

### 8.5. Hyper Kähler structure.

Again,  $Q$  is a quiver on  $k$  vertices and  $Q^d$  its double. We fix a dimension vector  $\alpha = (a_1, \dots, a_k) \in \mathbb{N}^k$  and a character  $\theta = (t_1, \dots, t_k) \in \mathbb{Z}^k$  and a weight

$\lambda = (\lambda_1, \dots, \lambda_k) \in \mathbb{C}^k$  such that the numerical conditions

$$\theta(\alpha) = \sum_{i=1}^k t_i a_i = 0 \quad \text{and} \quad \lambda(\alpha) = \sum_{i=1}^n \lambda_i a_i = 0$$

are satisfied. The first is required to have  $\theta$ -semistable representations, the second for  $\lambda$  to lie in the image of the complex moment map

$$\text{rep}_\alpha Q^d \xrightarrow{\mu_{\mathbb{C}}} M_\alpha^0(\mathbb{C}) \quad V \mapsto \sum_{a \in Q_a} [V_a, V_{a^*}]$$

where  $a^*$  is the arrow in  $Q_a^d$  corresponding to  $a \in Q_a$  (that is with the opposite direction).

Recall that the *quaternion algebra*  $\mathbb{H}$  is the 4-dimensional division algebra over  $\mathbb{R}$  defined by

$$\mathbb{H} = \mathbb{R}.1 \oplus \mathbb{R}.i \oplus \mathbb{R}.j \oplus \mathbb{R}.k \quad i^2 = j^2 = k^2 = -1 \quad k = ij = -ji$$

DEFINITION 8.29. A  $C^\infty$  (real) manifold  $M$  is said to be a hyper-Kähler manifold if  $\mathbb{H}$  acts on  $H$  by diffeomorphisms.

LEMMA 8.30. *For any quiver  $Q$ , the representation space  $\text{rep}_\alpha Q^d$  is a hyper-Kähler manifold.*

PROOF. We have to specify the actions. They are defined as follows, for  $V \in \text{rep}_\alpha Q^d$  for all arrows  $b \in Q_a^d$  and all arrows  $a \in Q_a$

$$\begin{aligned} (i.V)_b &= iV_b \\ (j.V)_a &= -V_{a^*}^\dagger \quad (j.V)_{a^*} = V_a^\dagger \\ (k.V)_a &= -iV_{a^*}^\dagger \quad (k.V)_{a^*} = iV_a^\dagger \end{aligned}$$

where this time we denote the *Hermitian adjoint* of a matrix  $M$  by  $M^\dagger$  to distinguish it from the star-operation on the arrows of  $Q^d$ . A calculation shows that these operations satisfy the required relations.  $\square$

In section 8.1 we introduced the *real moment map* for quiver representations. If we apply this to the double quiver  $Q^d$  we can take

$$\text{rep}_\alpha Q^d \xrightarrow{\mu_{\mathbb{R}}} \text{Lie } U(\alpha) \quad V \mapsto \sum_{b \in Q_a^d} \frac{i}{2} [V_b, V_b^\dagger]$$

We will use the action by non-zero elements of  $\mathbb{H}$  to obtain  $C^\infty$ -diffeomorphisms between certain subsets of  $\text{rep}_\alpha Q^d$ . Let  $h = \frac{i-k}{\sqrt{2}}$  then we have

$$\begin{aligned} \mu_{\mathbb{C}}(h.V) &= \frac{1}{2} \sum_{a \in Q_a} [iV_a + iV_{a^*}^\dagger, iV_{a^*} - iV_a^\dagger] \\ &= \frac{1}{2} \sum_{a \in Q_a} ( -[V_a, V_{a^*}] + [V_a, V_a^\dagger] - [V_{a^*}^\dagger, V_{a^*}] + [V_{a^*}^\dagger, V_a^\dagger] ) \\ &= \frac{1}{2} \sum_{a \in Q_a} ( [V_a, V_{a^*}]^\dagger - [V_a, V_{a^*}] ) + \frac{1}{2} \sum_{a \in Q_a} ( [V_a, V_a^\dagger] + [V_{a^*}^\dagger, V_{a^*}] ) \\ &= \frac{1}{2} (\mu_{\mathbb{C}}(V)^\dagger - \mu_{\mathbb{C}}(V)) - i\mu_{\mathbb{R}}(V) \end{aligned}$$

and

$$\begin{aligned} \mu_{\mathbb{R}}(h.V) &= \frac{i}{4} ( \sum_{a \in Q_a} [iV_a + iV_{a^*}^\dagger, -iV_a^\dagger - iV_{a^*}] + \sum_{a \in Q_a} [iV_{a^*} - iV_a^\dagger, -iV_{a^*}^\dagger + iV_a] ) \\ &= \frac{i}{4} \sum_{a \in Q_a} ( [V_a, V_a^\dagger] + [V_a, V_{a^*}] + [V_{a^*}^\dagger, V_a^\dagger] + [V_{a^*}^\dagger, V_{a^*}] \\ &\quad + [V_{a^*}^\dagger, V_{a^*}] - [V_{a^*}^\dagger, V_a] - [V_a^\dagger, V_{a^*}^\dagger] + [V_a^\dagger, V_a] ) \\ &= \frac{i}{4} (2\mu_{\mathbb{C}}(V) + 2\mu_{\mathbb{C}}(V)^\dagger) \end{aligned}$$

In particular we have

PROPOSITION 8.31. *If  $\lambda \in \mathbb{R}^k$ , then we have a homeomorphism between the real varieties*

$$\mu_{\mathbb{C}}^{-1}(\lambda \mathbb{1}_\alpha) \cap \mu_{\mathbb{R}}^{-1}(\underline{0}) \xrightarrow{h.} \mu_{\mathbb{C}}^{-1}(\underline{0}) \cap \mu_{\mathbb{R}}^{-1}(i\lambda \mathbb{1}_\alpha)$$

Moreover, the hyper-Kähler structure commutes with the base-change action of  $U(\alpha)$ , whence we have a natural one-to-one correspondence between the quotient spaces

$$(\mu_{\mathbb{C}}^{-1}(\lambda \mathbb{1}_\alpha) \cap \mu_{\mathbb{R}}^{-1}(\underline{0}))/U(\alpha) \xrightarrow{h.} (\mu_{\mathbb{C}}^{-1}(\underline{0}) \cap \mu_{\mathbb{R}}^{-1}(i\lambda \mathbb{1}_\alpha))/U(\alpha)$$

By the results of section 8.1 we can identify both sides. To begin, by definition of the complex moment map  $\mu_{\mathbb{C}}$  we have that

$$\mu_{\mathbb{C}}^{-1}(\underline{0}) = \underline{\text{rep}}_\alpha \Pi_0 \quad \text{and} \quad \mu_{\mathbb{C}}^{-1}(\lambda \mathbb{1}_\alpha) = \underline{\text{rep}}_\alpha \Pi_\lambda$$

Moreover, applying theorem 8.5 to the double quiver  $Q^d$  we have

$$\text{iss}_\alpha Q^d \simeq \mu_{\mathbb{R}}^{-1}(\underline{0})/U(\alpha) \quad \text{and} \quad M_\alpha^{ss}(Q^d, \lambda) \simeq \mu_{\mathbb{R}}^{-1}(i\lambda \mathbb{1}_\alpha)/U(\alpha)$$

when  $\lambda \in \mathbb{Z}^k$ . This concludes the proof of

THEOREM 8.32. *For a character  $\theta = (t_1, \dots, t_k) \in \mathbb{Z}^k$  such that  $\theta(\alpha) = 0$ , there is a natural one-to-one correspondence between*

$$\text{iss}_\alpha \Pi_\theta \xrightarrow{h.} M_\alpha^{ss}(\Pi_0, \theta)$$

which is an homeomorphism in the (induced) real topology.

Note however that this bijection does *not* respect the complex structures of these varieties. This is already clear from the fact that  $iss_\alpha \Pi_\theta$  is an affine complex variety and  $M_\alpha^{ss}(\Pi_0, \theta)$  is a projective bundle over  $iss_\alpha \Pi_0$ .

If  $V \in rep_\alpha Q^d$  belongs to  $\mu_{\mathbb{R}}^{-1}(0)$  we know that  $V$  is a semisimple representation, that is,

$$V = S_1^{\oplus e_1} \oplus \dots \oplus S_r^{\oplus e_r}$$

with the  $S_i$  simple representations of dimension vector  $\beta_i$ . Further, if  $W \in rep_\alpha^{-1}(i\theta|_\alpha)$ , then  $W$  is a direct sum of  $\theta$ -stable representations, that is,

$$W = T_1^{\oplus f_1} \oplus \dots \oplus T_s^{\oplus f_s}$$

with the  $T_i$   $\theta$ -stable representations of dimension vector  $\gamma_i$ . By the explicit form of the map, we have that if  $W = h.V$  that  $r = s$ ,  $e_i = f_i$  and  $\beta_i = \gamma_i$ . That is,

PROPOSITION 8.33. *Let  $\theta$  be a character such that  $\theta(\alpha) = 0$ , then the deformed preprojective algebra  $\Pi_\theta$  has semi-simple representations of dimension vector  $\alpha$  of representation type  $\tau = (e_1, \beta_1; \dots; e_r, \beta_r)$  if and only if the preprojective algebra  $\Pi_0$  has  $\theta$ -stable representations of dimension vectors  $\beta_i$  for all  $1 \leq i \leq r$ .*

*In particular,  $\Pi_\theta$  has a simple representation of dimension vector  $\alpha$  if and only if  $\Pi_0$  has a  $\theta$ -stable representation of dimension vector  $\alpha$ .*

The variety  $M_\alpha^{ss}(\Pi_0, \theta)$  is locally controlled by noncommutative algebras. Indeed, as in the case of moduli spaces of  $\theta$ -semistable quiver-representations, it is locally isomorphic to  $iss_\alpha(\Pi_0)_\Sigma$  for some universal localization of  $\Pi_0$ . We can determine the  $\alpha$ -smooth locus of the corresponding sheaf of Cayley-Hamilton algebras.

PROPOSITION 8.34. *Let  $\alpha \in \Sigma_\theta$ , then the  $\alpha$ -smooth locus of  $M_\alpha^{ss}(\Pi_0, \theta)$  is the open subvariety  $M_\alpha^s(\Pi_0, \theta)$  of  $\theta$ -stable representations of  $\Pi_0$ .*

*In particular, if the sheaf of Cayley-Hamilton algebras over  $M_\alpha^{ss}(\Pi_0, \theta)$  is a sheaf of  $\alpha$ -smooth algebras if and only if  $\alpha$  is a minimal dimension vector in  $\Sigma_\theta$ .*

PROOF. As  $\alpha \in \Sigma_\theta$  we know that  $iss_\alpha \Pi_\theta$  has dimension  $2p_Q(\alpha) = 2 - T_Q(\alpha, \alpha)$ . By the hyper-Kähler correspondence so is the dimension of  $M_\alpha^{ss}(\Pi_0, \theta)$ , whence the open subset of  $\mu_{\mathbb{C}}^{-1}(0)$  consisting of  $\theta$ -semistable representations has dimension

$$\alpha \cdot \alpha - 1 + 2p_Q(\alpha)$$

as there are  $\theta$ -stable representations in it. Take a  $GL(\alpha)$ -closed orbit  $\mathcal{O}(V)$  in this open set. That is,  $V$  is the direct sum of  $\theta$ -stable subrepresentations

$$V = S_1^{\oplus e_1} \oplus \dots \oplus S_r^{\oplus e_r}$$

with  $S_i$  a  $\theta$ -stable representation of  $\Pi_0$  of dimension vector  $\beta_i$  occurring in  $V$  with multiplicity  $e_i$  whence  $\alpha = \sum_i e_i \beta_i$ .

As all  $S_i$  are  $\Pi_0$ -representations we can determine the local quiver  $Q_V$  by the knowledge of all  $Ext_{\Pi_0}^1(S_i, S_j)$  from proposition 5.45

$$Ext_{\Pi_0}^1(S_i, S_j) = 2\delta_{ij} - T_Q(\beta_i, \beta_j)$$

But then the dimension of the normal space to the orbit is

$$dim Ext_{\Pi_0}^1(V, V) = 2 \sum_{i=1}^r e_i - T_Q(\alpha, \alpha)$$

whence the étale local structure in an  $n$ -smooth point is of the form  $GL(\alpha) \times^G L(\tau)Ext^1(V, V)$  where  $\tau = (e_1, \dots, e_r)$  and is therefore of dimension

$$\alpha \cdot \alpha + \sum_{i=1}^2 e_i^2 - T_Q(\alpha, \alpha)$$

This number is equal to the dimension of the subvariety of  $\theta$ -semistable representations of  $\Pi_0$  which has dimension  $\alpha \cdot \alpha - 1 + 2 - T_Q(\alpha, \alpha)$  if and only if  $r = 1$  and  $e_1 = 1$ , that is if  $V$  is  $\theta$ -stable.  $\square$


Even in points of  $M_\alpha^{ss}(\Pi_0, \theta)$  which are not in the  $\alpha$ -smooth locus we can use the local quiver to deduce combinatorial properties of the set of dimension vectors  $\Sigma_\theta$  of simple representations of  $\Pi_\theta$ .

PROPOSITION 8.35. *Let  $\alpha, \beta \in \Sigma_\theta$ , then*

- (1) *If  $T_Q(\alpha, \beta) \leq -2$  then  $\alpha + \beta \in \Sigma_\theta$ ,*
- (2) *If  $T_Q(\alpha, \beta) \geq -1$  then  $\alpha + \beta \notin \Sigma_\theta$ .*

PROOF. The property that  $\alpha$  and  $\beta$  are Schur roots of  $Q$  such that  $T_Q(\alpha, \beta) \leq -2$  ensures that  $\gamma = \alpha + \beta$  is a Schur root of  $Q$  and hence that  $\mu_{\mathbb{C}}^{-1}(\theta \mathbb{1}_\gamma)$  has dimension  $\gamma \cdot \gamma - 1 + 2p_Q(\gamma)$ , whence so is the subvariety of  $\theta$ -semistable  $\gamma$ -dimensional representations of  $\Pi_0$ . We have to prove that  $\Pi_0$  has a  $\theta$ -stable  $\gamma$ -dimensional representation.

Let  $V = S \oplus T$  with  $S$  resp.  $T$  a  $\theta$ -stable representation of  $\Pi_0$  of dimension vector  $\alpha$  resp.  $\beta$  (they exist by the hyper-Kähler correspondence). But then the local quiver  $Q_V$  has the following form



and by a calculation similar to the one in the foregoing proof we see that the image of the slice morphism in the space  $GL(\gamma) \times^{\mathbb{C}^* \times \mathbb{C}^*} rep_{(1,1)} Q_V$  has codimension 1. However, as  $T_Q(\alpha, \beta) \leq -2$  there are at least 3 algebraically independent new invariants coming from the non-loop cycles in  $Q_V$ , so they cannot all vanish on the image. This means that  $(1, \alpha; 1, \beta)$  cannot be the generic type for  $\theta$ -semistables of dimension  $\gamma$ , so by the stratification result, there must exist  $\theta$ -stables of dimension  $\gamma$ .

For the second assertion, assume that  $\gamma = \alpha + \beta$  is the dimension vector of a simple representation of  $\Pi_\theta$ , then  $iss_\gamma \Pi_\theta$  has dimension  $2p_Q(\gamma) = 2 - T_Q(\alpha, \beta, \alpha + \beta) = 2p_Q(\alpha) + 2p_Q(\beta)$  whence so is the dimension of  $M_\gamma^{ss}(\Pi_0, \theta)$ . By assumption  $(1, \alpha; 1, \beta)$  cannot be the generic type for  $\theta$ -semistable representations, but the stratum consisting of direct sums  $S \oplus T$  with  $S \in M_\alpha^s(\Pi_0, \theta)$  and  $T \in M_\beta^s(Q, \theta)$  has the same dimension as the total space, a contradiction.  $\square$

The first part of the foregoing proof can also be used to show that usually the moduli spaces  $M_\alpha^{ss}(\Pi, \theta)$  and the quotient varieties  $iss_\alpha \Pi_\theta$  have lots of singularities.

PROPOSITION 8.36. *Let  $\alpha \in |\sigma_{\theta}|$  such that  $\alpha = \beta + \gamma$  with  $\beta, \gamma \in \Sigma_\theta$ . Then,*

$$M_\alpha^{ss}(\Pi_0, \theta) \quad \text{and} \quad iss_\alpha \Pi_\theta$$

*is singular along the stratum of points of type  $(1, \beta; 1, \gamma)$ .*




that two such  $2n$  tuples are determined only up to permuting the couples  $(x_i, y_i)$ . The  $n$ -functions  $F_j$  give a completely integrable system on the phase space via *Liouville's theorem*, see for example [1].

In the classical case, all points are assumed to lie on the real axis and the potential is repulsive so that collisions do not appear. G. Wilson [82] considered an alternative where the points are assumed to lie in the complex numbers and such that the potential is attractive (to allow for collisions), that is, the Hamiltonian is of the form

$$H = \frac{1}{2} \sum_i y_i^2 - \sum_{i < j} \frac{1}{(x_i - x_j)^2}$$

giving again rise to a dynamical system via the equations of motion. One recovers the classical situation back if the particles are assumed only to move on the imaginary axis.



In general, we want to extend the phase space of  $n$  distinct points analytically to allow for collisions.

When all the points are distinct, that is, if all eigenvalues of  $X$  are distinct we will see in a moment that there is a unique  $GL_n(\mathbb{C})$ -orbit of couples of  $n \times n$  matrices (up to permuting the  $n$  couples  $(x_i, y_i)$ ).


$$X = \begin{bmatrix} x_1 & & & & \\ & \ddots & & & \\ & & x_n & & \end{bmatrix} \quad \text{and} \quad Y = \begin{bmatrix} y_1 & \frac{1}{x_1-x_2} & \cdots & \cdots & \frac{1}{x_1-x_n} \\ \frac{1}{x_2-x_1} & y_2 & \ddots & & \vdots \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ \vdots & & \ddots & \ddots & \frac{1}{x_{n-1}-x_n} \\ \frac{1}{x_n-x_1} & \cdots & \cdots & \frac{1}{x_n-x_{n-1}} & y_n \end{bmatrix}$$

For matrix couples in this standard form one verifies that

$$[Y, X] + \begin{bmatrix} 1 & \cdots & 1 \\ \vdots & \ddots & \vdots \\ 1 & \cdots & 1 \end{bmatrix} = \mathbb{1}_n$$

This equality suggests an approach to extend the phase space of  $n$  distinct complex Calogero particles to allow for collisions.

Assign the representation  $(X, Y, u, v) \in \text{rep}_\alpha \mathbb{M}$  where  $\alpha = (1, n)$  and  $\mathbb{M}$  is the path algebra of the quiver  $Q^d$  is



where  $X$  and  $Y$  are the matrices above and where

$$u = \begin{bmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{bmatrix} \quad v = [1 \quad 1 \quad \dots \quad 1]$$

Recall that the complex moment map for this quiver-setting is defined to be

$$\text{rep}_\alpha Q^d = M_n(\mathbb{C}) \oplus M_n(\mathbb{C}) \oplus \mathbb{C}^n \oplus \mathbb{C}^{n*} \xrightarrow{\mu} \mathbb{C} \oplus M_n(\mathbb{C})$$

$$(X, Y, u, v) \mapsto (-v.u, [Y, X] + u.v)$$

Therefore, the above equation entails that  $(X, Y, u, v) \in \mu_{\mathbb{C}}^{-1}(\theta\mathbb{1}_\alpha)$  where  $\theta = (-n, 1)$ , that is  $(X, Y, u, v) \in \text{rep}_\alpha \Pi_\theta$ . Observe that  $\alpha = (1, n) \in \Sigma_\theta$  (in fact,  $\alpha$  is a minimal element in  $\Sigma_\theta$ ), whence theorem 8.13,  $\text{rep}_\alpha \Pi_\theta$  is an irreducible complete intersection of dimension  $d = n^2 + 2n$  and there are  $\alpha$ -dimensional simple representations of  $\Pi_\theta$ . In particular,  $\text{iss}_\alpha \Pi_\theta$  is an irreducible variety of dimension  $2n$ .

We can define the *phase space for Calogero collisions* of  $n$  particles to be the quotient space

$$\text{Calo}_n = \text{iss}_\alpha \Pi_\theta$$

In a moment we will show that this is actually an orbit-space and :

**THEOREM 8.37.** *The phase space  $\text{Calo}_n$  of Calogero collisions of  $n$ -particles is a connected complex manifold of dimension  $2n$ .*

**THEOREM 8.38.** *Let  $\text{rep}_\alpha \mathbb{M} \xrightarrow{\mu_{\mathbb{C}}} M_\alpha^0(\mathbb{C})$  be the complex moment map, then any  $V = (X, Y, u, v) \in \text{rep}_\alpha \Pi_\theta$  is a  $\theta$ -stable representation. Therefore,*

$\text{Calo}_n = \mu_{\mathbb{C}}^{-1}(\theta\mathbb{1}_\alpha)/GL(\alpha) = \text{iss}_\alpha \Pi_\theta \simeq (\mu_{\mathbb{C}}^{-1}(\theta\mathbb{1}_\alpha) \cap \text{rep}_\alpha^s \mathbb{M})/GL(\alpha) = M_\alpha^s(\Pi_\theta, \theta)$  and is therefore a complex manifold of dimension  $2n$ , which is connected by theorem 8.13.

**PROOF.** The result will follow if we can prove that any Calogero quadruple  $(X, Y, u, v)$  has the property that  $u$  is a cyclic vector, that is, lies in  $\text{rep}_\alpha^s \mathbb{M}$ .

Assume that  $U$  is a subspace of  $\mathbb{C}^n$  stable under  $X$  and  $Y$  and containing  $u$ .  $U$  is then also stable under left multiplication with the matrix

$$A = [X, Y] + \mathbb{1}_n$$

and we have that  $\text{tr}(A | U) = \text{tr}(\mathbb{1}_n | U) = \dim U$ . On the other hand,  $A = u.v$  and therefore

$$A \cdot \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix} = \begin{bmatrix} u_1 \\ \vdots \\ u_n \end{bmatrix} \cdot [v_1 \quad \dots \quad v_n] \cdot \begin{bmatrix} c_1 \\ \vdots \\ c_n \end{bmatrix} = \left( \sum_{i=1}^n v_i c_i \right) \begin{bmatrix} u_1 \\ \vdots \\ u_n \end{bmatrix}$$



Hence, if we take a basis for  $U$  containing  $u$ , then we have that

$$\text{tr}(A | U) = a$$

where  $A.u = au$ , that is  $a = \sum u_i v_i$ .

But then,  $\text{tr}(A | U) = \dim U$  is independent of the choice of  $U$ . Now,  $\mathbb{C}^n$  is clearly a subspace stable under  $X$  and  $Y$  and containing  $u$ , so we must have that  $a = n$  and so the only subspace  $U$  possible is  $\mathbb{C}^n$  proving cyclicity of  $u$  with respect to the matrix-couple  $(X, Y)$ .  $\square$

Again, it follows that we can cover the phase space  $\text{Calo}_n$  by open subsets

$$\text{Calo}_n(\sigma) = \{(X, Y, u, v) \text{ in } \sigma\text{-standard form such that } [Y, X] + u.v = \mathbb{1}_n \}$$

where  $\sigma$  runs over the Hilbert  $n$ -stairs.

EXAMPLE 8.39. The phase space  $\text{Calo}_2$ .

Consider  $\text{Calo}_2(\begin{smallmatrix} \square \\ \square \end{smallmatrix})$ . Because

$$\left[ \begin{bmatrix} 0 & a \\ 1 & b \end{bmatrix}, \begin{bmatrix} c & d \\ e & f \end{bmatrix} \right] + \begin{bmatrix} 1 \\ 0 \end{bmatrix} \cdot [g \quad h] - \mathbb{1}_2 = \begin{bmatrix} g - d + ae - 1 & h + af - ac - bd \\ c - f + be & d - ae - 1 \end{bmatrix}$$

We obtain after taking Groebner bases that the defining equations are

$$\begin{cases} g &= 2 \\ h &= b \\ f &= c + eh \\ d &= 1 + ae \end{cases}$$

In particular we find

$$\text{Calo}_2(\begin{smallmatrix} \square \\ \square \end{smallmatrix}) = \left\{ \left( \begin{bmatrix} 0 & a \\ 1 & b \end{bmatrix}, \begin{bmatrix} c & 1+ae \\ e & c+be \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \end{bmatrix}, [2 \quad b] \right) \mid a, b, c, e \in \mathbb{C} \right\} \simeq \mathbb{C}^4$$

and a similar description holds for  $\text{Calo}_2(\begin{smallmatrix} \blacksquare \\ \blacksquare \end{smallmatrix})$ .

EXAMPLE 8.40. The phase space  $\text{Calo}_3$ .

We claim that

$$\text{Calo}_3(\begin{smallmatrix} \square & \square \\ \blacksquare & \square \end{smallmatrix}) \simeq \mathbb{C}^6$$

For, if we compute the  $3 \times 3$  matrix

$$\left[ \begin{bmatrix} 0 & a & b \\ 1 & c & d \\ 0 & e & f \end{bmatrix}, \begin{bmatrix} 0 & g & h \\ 0 & i & j \\ 1 & k & l \end{bmatrix} \right] + \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \cdot [m \quad n \quad o] - \mathbb{1}_3$$

then the Groebner basis for its entries gives the following defining equations

$$\begin{cases} m &= 3 \\ n &= c + k \\ o &= i + l \\ f &= k \\ d &= o - l \\ g &= 2 + b \\ l &= g - ej - kl + ko \\ h &= 2jk + 2l^2 - jn - 3lo + o^2 \\ a &= 2k^2 - 2el - kn + eo \end{cases}$$

In a similar manner one can show that

$$\text{Calo}_3(\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}) \simeq \mathbb{C}^6 \quad \text{but} \quad \text{Calo}_3(\begin{smallmatrix} \square & \square \\ \square & \blacksquare \end{smallmatrix})$$

is again more difficult to describe.

We can identify the classical Calogero situations as an open subset of  $Calo_n$ .

PROPOSITION 8.41. *Let  $(X, Y, u, v) \in rep_\alpha \Pi_\theta$  and suppose that  $X$  is diagonalizable. Then*

- (1) *all eigenvalues of  $X$  are distinct, and*
- (2) *the  $GL(\alpha)$ -orbit contains a representative such that*

$$X = \begin{bmatrix} \lambda_1 & & & & \\ & \ddots & & & \\ & & \ddots & & \\ & & & \ddots & \\ & & & & \lambda_n \end{bmatrix} \quad Y = \begin{bmatrix} \alpha_1 & \frac{1}{\lambda_1 - \lambda_2} & \cdots & \cdots & \frac{1}{\lambda_1 - \lambda_n} \\ \frac{1}{\lambda_2 - \lambda_1} & \alpha_2 & \ddots & & \vdots \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ \vdots & & \ddots & \ddots & \frac{1}{\lambda_{n-1} - \lambda_n} \\ \frac{1}{\lambda_n - \lambda_1} & \cdots & \cdots & \frac{1}{\lambda_n - \lambda_{n-1}} & \alpha_n \end{bmatrix}$$

$$u = \begin{bmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{bmatrix} \quad v = [1 \quad 1 \quad \cdots \quad 1]$$

and this representative is unique up to permutation of the  $n$  couples  $(\lambda_i, \alpha_i)$ .

PROOF. Choose a representative with  $X$  a diagonal matrix as indicated. Equating the diagonal entries in  $[Y, X] + u.v = \mathbb{1}_n$  we obtain that for all  $1 \leq i \leq n$  we have  $u_i v_i = 1$ . Hence, none of the entries of

$$[X, Y] + \mathbb{1}_n = u.v$$

is zero. Consequently, by equating the  $(i, j)$ -entry it follows that  $\lambda_i \neq \lambda_j$  for  $i \neq j$ .

The representative with  $X$  a diagonal matrix is therefore unique up to the action of a diagonal matrix  $D$  and of a permutation. The freedom in  $D$  allows us to normalize  $u$  and  $v$  as indicated, the effect of the permutation is described in the last sentence.

Finally, the precise form of  $Y$  can be calculated from the normalized forms of  $X$ ,  $u$  and  $v$  and the equation  $[Y, X] + u.v = \mathbb{1}_n$ .  $\square$

Invoking the hyper-Kähler structure on  $rep_\alpha \mathbb{M}$  we have by theorem 8.32an homeomorphism, in fact in this case a  $C^\infty$ -diffeomorphism between the Calogero phase-space and the Hilbert scheme

$$Calo_n = iss_\alpha \Pi_\theta \xrightarrow{h.} M_\alpha^{ss}(\Pi_0, \theta) = Hilb_n$$

### 8.7. Coadjoint orbits.

In this section we will give an important application of  $\mathfrak{g}\mathfrak{e}\mathfrak{o} @_n$  developed in the foregoing chapter. If  $\alpha$  is a minimal dimension vector in  $\Sigma_\theta$  we will prove that the quotient variety  $iss_\alpha \Pi_\theta$  is smooth and a coadjoint orbit for the dual of the necklace algebra. In particular, the phase space of Calogero particles is a coadjoint orbit.

We fix a quiver  $Q$  on  $k$  vertices, a dimension vector  $\alpha \in \mathbb{N}^k$  and a character  $\theta \in \mathbb{Z}^k$  such that  $\theta(\alpha) = 0$  with corresponding weight  $\theta \mathbb{1}_\alpha$ . Recall that  $\Sigma_\theta$  is the subset of dimension vectors  $\alpha$  such that

$$p_Q(\alpha) > p_Q(\beta_1) + \cdots + p_Q(\beta_r)$$

for all decompositions  $\alpha = \beta_1 + \dots + \beta_r$  with the  $\beta_i \in \Delta^+\theta$ , that is,  $\beta_i$  is a positive root for the quiver  $Q$  and  $\theta(\beta_i) = 0$ . With  $\Sigma_\theta^{min}$  we will denote the subset of *minimal dimension vectors* in  $\Sigma_\theta$ , that is, such that for all  $\beta < \alpha$  we have  $\beta \notin \Sigma_\theta$ .

PROPOSITION 8.42. *If  $\alpha \in \Sigma_\theta^{min}$ , then the deformed preprojective algebra  $\Pi_\theta$  is  $\alpha$ -smooth, that is  $rep_\alpha \Pi_\theta$  is a smooth  $GL(\alpha)$ -variety of dimension  $d = \alpha \cdot \alpha - 1 + 2p_Q(\alpha)$ .*

Moreover, the quotient variety  $iss_\alpha \Pi_\theta$  is a smooth variety of dimension  $2p_Q(\alpha)$ , and the quotient map

$$rep_\alpha \Pi_\theta \longrightarrow iss_\alpha \Pi_\theta$$

is a principal  $PGL(\alpha)$ -fibration, so determines a central simple algebra.

PROOF. Let  $V \in rep_\alpha \Pi_\theta$  and let  $V^{ss}$  be its semisimplification. As  $\Sigma_\theta$  is the set of simple dimension vectors of  $\Pi_\theta$  by theorem 8.14 and  $\alpha$  is a minimal dimension vector in this set,  $V^{ss}$  must be simple. As  $V^{ss}$  is the direct sum of the Jordan-Hölder components of  $V$ , it follows that  $V \simeq V^{ss}$  is simple and hence its orbit  $\mathcal{O}(V)$  is closed. As the stabilizer subgroup of  $V$  is  $\mathbb{C}^*\mathbb{1}_\alpha$  computing the differential of the complex moment map shows that  $V$  is a smooth point of  $\mu_{\mathbb{C}}^{-1}(\theta\mathbb{1}_\alpha = rep_\alpha \Pi_\theta)$ .

Therefore,  $rep_\alpha \Pi_\theta$  is a smooth  $GL(\alpha)$ -variety whence  $\Pi_\theta$  is  $\alpha$ -smooth. Because each  $\alpha$ -dimensional representation is simple, the quotient map

$$rep_\alpha \Pi_\theta \xrightarrow{\pi} iss_\alpha \Pi_\theta$$

is a principal  $PGL(\alpha)$ -fibration in the étale topology. The total space being smooth, so is the basespace  $iss_\alpha \Pi_\theta$ .  $\square$

The trace pairing identifies  $rep_\alpha Q^d$  with the *cotangent bundle*  $T^* rep_\alpha Q$  and as such it comes equipped with a *canonical symplectic structure*. More explicit, for every arrow  $\textcircled{j} \xleftarrow{a} \textcircled{i}$  in  $Q$  we have an  $a_j \times a_i$  matrix of coordinate functions  $A_{uv}$  with  $1 \leq u \leq a_j$  and  $1 \leq v \leq a_i$  and for the adjoined arrow  $\textcircled{i} \xrightarrow{a^*} \textcircled{j}$  in  $Q^d$  an  $a_i \times a_j$  matrix of coordinate functions  $A_{vu}^*$ . The canonical symplectic structure on  $rep_\alpha Q^d$  is then induced by the closed 2-form

$$\omega = \sum_{\substack{1 \leq v \leq a_i \\ 1 \leq u \leq a_j}} dA_{uv} \wedge dA_{vu}^* \\ \textcircled{j} \xleftarrow{a} \textcircled{i}$$

This symplectic structure induces a *Poisson bracket* on the coordinate ring  $\mathbb{C}[rep_\alpha Q^d]$  by the formula

$$\{f, g\} = \sum_{\substack{1 \leq v \leq a_i \\ 1 \leq u \leq a_j}} \left( \frac{\partial f}{\partial A_{uv}} \frac{\partial g}{\partial A_{vu}^*} - \frac{\partial f}{\partial A_{vu}^*} \frac{\partial g}{\partial A_{uv}} \right) \\ \textcircled{j} \xleftarrow{a} \textcircled{i}$$

The basechange action of  $GL(\alpha)$  on the representation space  $rep_\alpha Q^d$  is *symplectic* which means that for all *tangentvectors*  $t, t' \in T rep_\alpha Q^d$  we have for the induced  $GL(\alpha)$  action that

$$\omega(t, t') = \omega(g.t, g.t')$$

for all  $g \in GL(\alpha)$ .

The infinitesimal  $GL(\alpha)$  action gives a Lie algebra homomorphism

$$\text{Lie } PGL(\alpha) \longrightarrow \text{Vect}_\omega \text{ rep}_\alpha Q^d$$

which factorizes through a Lie algebra morphism  $H$  to the coordinate ring making the diagram below commute

$$\begin{array}{ccc} & \text{Lie } PGL(\alpha) & \\ & \swarrow \scriptstyle H = \mu_{\mathbb{C}}^* & \searrow \\ \mathbb{C}[\text{rep}_\alpha Q^d] & \xrightarrow{f \mapsto \xi_f} & \text{Vect}_\omega \text{ rep}_\alpha Q^d \end{array}$$

where  $\mu_{\mathbb{C}}$  is the complex moment map introduced before. We say that the action of  $GL(\alpha)$  on  $\text{rep}_\alpha M$  is *Hamiltonian*.

This makes the ring of polynomial invariants  $\mathbb{C}[\text{rep}_\alpha Q^d]^{GL(\alpha)}$  into a *Poisson algebra* and we will write

$$\mathfrak{lie} = (\mathbb{C}[\text{rep}_\alpha Q^d]^{GL(\alpha)}, \{-, -\})$$

for the corresponding abstract *infinite dimensional Lie algebra*.

The dual space of this Lie algebra  $\mathfrak{lie}^*$  is then a *Poisson manifold* equipped with the *Kirillov-Kostant bracket*.

*Evaluation at a point* in the quotient variety  $\text{iss}_\alpha Q^d$  defines a linear function on  $\mathfrak{lie}$  and therefore evaluation gives an embedding

$$\text{iss}_\alpha Q^d \hookrightarrow \mathfrak{lie}^*$$

as Poisson varieties. That is, the induced map on the polynomial functions is a morphism of Poisson algebras.

Let us return to the setting of deformed preprojective algebras. So let  $\theta$  be a character with  $\theta(\alpha) = 0$  and corresponding weight  $\theta \mathbb{1}_\alpha \in \text{Lie } PGL(\alpha)$ .

**THEOREM 8.43.** *Let  $\alpha \in \Sigma_\theta^{\text{min}}$ , then  $\text{iss}_\alpha \Pi_\theta$  is an affine symplectic manifold and the Poisson embeddings*

$$\text{iss}_\alpha \Pi_\theta \hookrightarrow \text{iss}_\alpha Q^d \hookrightarrow \mathfrak{lie}^*$$

*make  $\text{iss}_\alpha \Pi_\theta$  into a closed coadjoint orbit of the infinite dimensional Lie algebra  $\mathfrak{lie}^*$ .*

**PROOF.** We know from proposition 8.42 that  $\text{iss}_\alpha \Pi_\theta$  is a smooth affine variety and that  $PGL(\alpha)$  acts freely on  $\mu_{\mathbb{C}}^{-1}(\theta \mathbb{1}_\alpha) = \text{rep}_\alpha \Pi_\theta$ . Moreover, the infinitesimal coadjoint action of  $\mathfrak{lie}$  on  $\mathfrak{lie}^*$  preserves  $\text{iss}_\alpha \Pi_\theta$  and therefore  $\mathbb{C}[\text{iss}_\alpha \Pi_\theta]$  is a quotient Lie  $\overline{\mathfrak{lie}}$  algebra (for the induced bracket) of  $\mathfrak{lie}$ .

In general, if  $X$  is a smooth affine variety, then the differentials of polynomial functions on  $X$  span the tangent spaces at all points  $x$  of  $X$ . Therefore, if  $X$  is in addition symplectic, the infinitesimal Hamiltonian action of the Lie algebra  $\mathbb{C}[X]$  (with the natural Poisson bracket) on  $X$  is infinitesimally transitive. But then, the evaluation map makes  $X$  a coadjoint orbit of the dual Lie algebra  $\mathbb{C}[X]^*$ .

Hence, the quotient variety  $\text{iss}_\alpha \Pi_\theta$  is a coadjoint orbit in  $\overline{\mathfrak{lie}}^*$ . Therefore, the infinite dimensional group  $\text{Ham}$  generated by all *Hamiltonian flows* on  $\text{iss}_\alpha \Pi_\theta$  acts with *open* orbits.

By proposition 8.42  $\text{iss}_\alpha \Pi_\theta$  is an irreducible variety, whence is a single *Ham*-orbit, finishing the proof.  $\square$

The Lie algebra  $\mathfrak{lie}$  depends on the dimension vector  $\alpha$ . By the general principle of  $\mathfrak{geo} @_{\mathbf{n}}$  we would like to construct a noncommutative variety from a family of coadjoint quotient spaces of deformed preprojective algebras. For this reason we need a larger Lie algebra, the *necklace Lie algebra*.

Recall that the necklace Lie algebra introduced in section 7.8

$$\mathfrak{neck} = \mathfrak{dR}_{rel}^0 \mathbb{C}Q^d = \frac{\mathbb{C}Q^d}{[\mathbb{C}Q^d, \mathbb{C}Q^d]}$$

is the vectorspace with basis all the necklace words  $w$  in the quiver  $Q^d$ , that is, all equivalence classes of oriented cycles in the quiver  $Q^d$ , equipped with the Kontsevich bracket

$$\{w_1, w_2\}_K = \sum_{a \in Q_a} \left( \frac{\partial w_1}{\partial a} \frac{\partial w_2}{\partial a^*} - \frac{\partial w_1}{\partial a^*} \frac{\partial w_2}{\partial a} \right) \pmod{[\mathbb{C}Q^d, \mathbb{C}Q^d]}$$

We recall that the algebra of polynomial quiver invariants  $\mathbb{C}[iss_{\alpha} Q^d] = \mathbb{C}[rep_{\alpha} Q^d]^{GL(\alpha)}$  is generated by traces of necklace words. That is, we have a map

$$\mathfrak{neck} = \frac{\mathbb{C}Q^d}{[\mathbb{C}Q^d, \mathbb{C}Q^d]} \xrightarrow{tr} \mathfrak{lie} = \mathbb{C}[iss_{\alpha} Q^d]$$

Recalling the definition of the Lie bracket on  $\mathfrak{lie}$  we see that this map is actually a Lie algebra map, that is, for all necklace words  $w_1$  and  $w_2$  in  $Q^d$  we have the identity

$$tr \{w_1, w_2\}_K = \{tr(w_1), tr(w_2)\}$$

Now, the image of  $tr$  contains a set of *algebra* generators of  $\mathbb{C}[iss_{\alpha} Q^d]$ , so the elements  $tr \mathfrak{neck}$  are enough to separate points in  $iss_{\alpha} Q^d$  and in the closed subvariety  $iss_{\alpha} \Pi_{\theta}$ . That is, the composition

$$iss_{\alpha} \Pi_{\theta} \hookrightarrow iss_{\alpha} Q^d \xrightarrow{tr^*} \mathfrak{neck}^*$$

is injective. Again, the differentials of functions on  $iss_{\alpha} \Pi_{\theta}$  obtained by restricting traces of necklace words span the tangent spaces at all points if the affine variety  $iss_{\alpha} \Pi_{\theta}$  is smooth. That is, we have :

**THEOREM 8.44.** *Let  $\alpha \in \Sigma_{\theta}^{min}$ . Then, the quotient variety of the preprojective algebra  $iss_{\alpha} \Pi_{\theta}$  is an affine smooth manifold and the embeddings*

$$iss_{\alpha} \Pi_{\theta} \hookrightarrow iss_{\alpha} Q^d \hookrightarrow \mathfrak{lie}^* \hookrightarrow \mathfrak{neck}^*$$

*make  $iss_{\alpha} \Pi_{\theta}$  into a closed coadjoint orbit in the dual of the necklace Lie algebra  $\mathfrak{neck}$ .*

We have proved in section 7.8 that there is an exact sequence of Lie algebras

$$0 \longrightarrow \underbrace{\mathbb{C} \oplus \dots \oplus \mathbb{C}}_k \longrightarrow \mathfrak{neck} \longrightarrow Der_{\omega} \mathbb{C}Q^d \longrightarrow 0$$

That is, the necklace Lie algebra  $\mathfrak{neck}$  is a central extension of the Lie algebra of symplectic derivations of  $\mathbb{C}Q^d$ . This Lie algebra corresponds to the automorphism group of all  $B = \mathbb{C} \times \dots \times \mathbb{C}$ -automorphisms of the path algebra  $\mathbb{C}Q^d$  preserving the *moment map element*, the commutator

$$c = \sum_{a \in Q_a} [a, a^*]$$

That is, we expect a transitive action of an extension of this automorphism group on the quotient varieties of deformed preprojective algebras  $iss_\alpha \Pi_\theta$  when  $\alpha \in \Sigma_\theta^{min}$ . Further, it should be observed that these coadjoint cases are precisely the situations where the preprojective algebra  $\Pi_\theta$  is  $\alpha$ -smooth. That is, whereas the Lie algebra of vectorfields of the smooth noncommutative variety corresponding to  $\mathbb{C}Q^d$  has rather unpredictable behavior on the *singular* noncommutative closed subvariety corresponding to the quotient algebra  $\Pi_\theta$ , it behaves as expected on those  $\alpha$ -dimensional components where  $\Pi_\theta$  is  $\alpha$ -smooth.

**8.8. Adelic Grassmannian.**

At the moment of writing it is unclear which coadjoint orbits  $iss_\alpha \Pi_\theta$  should be taken together to form an object in  $\mathfrak{geo} \ @_{\mathbf{n}}$ , for a general quiver  $Q$ . In this section we will briefly recall how the phase spaces  $Calo_n$  of Calogero particles can be assembled together to form an infinite dimensional cellular complex, the *adelic Grassmannian*  $Gr^{ad}$ .

Let  $\lambda \in \mathbb{C}$ , a subset  $V \subset \mathbb{C}[x]$  is said to be  $\lambda$ -primary if there is some power  $r \in \mathbb{N}_+$  such that

$$(x - \lambda)^r \mathbb{C}[x] \subset V \subset \mathbb{C}[x]$$

A subset  $V \subset \mathbb{C}[x]$  is said to be *primary decomposable* if it is the finite intersection

$$V = V_{\lambda_1} \cap \dots \cap V_{\lambda_r}$$

with  $\lambda_i \neq \lambda_j$  if  $i \neq j$  and  $V_{\lambda_i}$  is a  $\lambda_i$ -primary subset. Let  $k_{\lambda_i}$  be the codimension of  $V_{\lambda_i}$  in  $\mathbb{C}[x]$  and consider the polynomial

$$p_V(x) = \prod_{i=1}^r (x - \lambda_i)^{k_{\lambda_i}}$$

Finally, take  $W = p_V(x)^{-1}V$ , then  $W$  is a vectorsubspace of the rational function-field  $\mathbb{C}(x)$  in one variable.

DEFINITION 8.45. The *adelic Grassmannian*  $Gr^{ad}$  is the set of subspaces  $W \subset \mathbb{C}(x)$  that arise in this way.

We can decompose  $Gr^{ad}$  in affine cells as follows. For a fixed  $\lambda \in \mathbb{C}$  we define

$$Gr_\lambda = \{W \in Gr^{ad} \mid \exists k, l \in \mathbb{N} : (x - \lambda)^k \mathbb{C}[x] \subset W \subset (x - \lambda)^{-l} \mathbb{C}[x]\}$$

Then, we can write every element  $w \in W$  as a *Laurent series*

$$w = \alpha_s (x - \lambda)^s + \text{higher terms}$$

Consider the increasing set of integers  $S = \{s_0 < s_1 < \dots\}$  consisting of all *degrees*  $s$  of elements  $w \in W$ . Now, define natural numbers

$$v_i = i - s_i \quad \text{then} \quad v_0 \geq v_1 \geq \dots \geq v_z = 0 = v_{z+1} = \dots$$

That is, to  $W \in Gr_\lambda$  we can associate a *partition*

$$p(W) = (v_0, v_1, \dots, v_{z-1})$$

Conversely, if  $p$  is a partition of some  $n$ , then the set of all  $W \in Gr_\lambda$  with associated partition  $p_W = p$  form an affine space  $\mathbb{A}^n$  of dimension  $n$ . Hence,  $Gr_\lambda$  has a cellular structure indexed by the set of all partitions.

As  $Gr^{ad} = \prod_{\lambda \in \mathbb{C}} Gr_\lambda$  because for any  $W \in Gr^{ad}$  there are uniquely determined  $W(\lambda_i) \in Gr_{\lambda_i}$  such that  $W = W(\lambda_1) \cap \dots \cap W(\lambda_r)$ , there is a natural number  $n$

associated to  $W$  where  $n = |p_i|$  where  $p_i = p(W(\lambda_i))$  is the partition determined by  $W(\lambda_i)$ . Again, all  $W \in Gr^{ad}$  with corresponding  $(\lambda_1, p_1; \dots; \lambda_r, p_r)$  for an affine cell  $\mathbb{A}^n$  of dimension  $n$ . In his way, the adelic Grassmannian  $Gr^{ad}$  becomes an infinite cellular space with the cells indexed by  $r$ -tuples of complex numbers and partitions for all  $r \geq 0$ . The adelic Grassmannian is an important object in the theory of dynamical systems as it parametrizes rational solutions of the *KP hierarchy*. A surprising connection between  $Gr^{ad}$  and the Calogero system was discovered by G. Wilson in [82].

**THEOREM 8.46.** *Let  $Gr^{ad}(n)$  be the collection of all cells of dimension  $n$  in  $gr^{ad}$ , then there is a set-theoretic bijection*

$$Gr^{ad}(n) \longleftrightarrow Calo_n$$

*between  $Gr^{ad}(n)$  and the phase space of  $n$  Calogero particles.*

The adelic Grassmannian also appears in the study of right ideals of the first *Weyl algebra*

$$A_1(\mathbb{C}) = \frac{\mathbb{C}\langle x, y \rangle}{(xy - yx - 1)}$$

which is an infinite dimensional simple  $\mathbb{C}$ -algebra, having no finite dimensional representations. Consider right ideals of  $A_1(\mathbb{C})$  under isomorphism, that is

$$\mathfrak{p} \simeq \mathfrak{p}' \quad \text{iff} \quad f \cdot \mathfrak{p} = g \cdot \mathfrak{p}' \quad \text{for some } f, g \in A_1(\mathbb{C}).$$

If we denote with  $D_1(\mathbb{C})$  the *Weyl skewfield*, that is, the field of fractions of  $A_1(\mathbb{C})$ , then the foregoing can also be expressed as

$$\mathfrak{p} \simeq \mathfrak{p}' \quad \text{iff} \quad \mathfrak{p} = h \cdot \mathfrak{p}' \quad \text{for some } h \in D_1(\mathbb{C}).$$

The set of isomorphism classes will be denoted by *Weyl*.

The connection between right ideals of  $A_1(\mathbb{C})$  and  $gr^{ad}$  is contained (in disguise) in the paper of R. Cannings and M. Holland [11].  $A_1(\mathbb{C})$  acts as differential operators on  $\mathbb{C}[x]$  and for every right ideal  $I$  of  $A_1(\mathbb{C})$  they show that  $I \cdot \mathbb{C}[x]$  is primary decomposable. Conversely, if  $V \subset \mathbb{C}[x]$  is primary decomposable, they associate the right ideal

$$I_V = \{ \theta \in A_1(\mathbb{C}) \mid \theta \cdot \mathbb{C}[x] \subset V \}$$

of  $A_1(\mathbb{C})$  to it. Moreover, isomorphism classes of right ideals correspond to studying primary decomposable subspaces under multiplication with polynomials. Hence,

$$Gr^{ad} \simeq Weyl$$

The group  $Aut A_1(\mathbb{C})$  of  $\mathbb{C}$ -algebra automorphisms of  $A_1(\mathbb{C})$  acts on the set of right ideals of  $A_1(\mathbb{C})$  and respects the notion of isomorphism whence acts on *Weyl*. The group  $Aut A_1(\mathbb{C})$  is generated by automorphisms  $\sigma_i^f$  defined by

$$\begin{cases} \sigma_1^f(x) = x + f(y) \\ \sigma_1^f(y) = y \end{cases} \quad \text{with } f \in \mathbb{C}[y], \quad \begin{cases} \sigma_2^f(x) = x \\ \sigma_2^f(y) = y + f(x) \end{cases} \quad \text{with } f \in \mathbb{C}[x]$$

We claim that for any polynomial in one variable  $f(z) \in \mathbb{C}[z]$  we have that

$$f(xy) \cdot x^n = x^n \cdot f(xy - n) \quad \text{and} \quad f(xy) \cdot y^n = y^n \cdot f(xy + n)$$

Indeed, we have  $(xy) \cdot x = x \cdot (yx) = x \cdot (xy - 1)$  and therefore

$$f(xy) \cdot x = x \cdot f(xy - 1)$$

from which the claim follows by recursion. In particular, as  $x^n \cdot y^n = x^{n-1}(xy)y^{n-1} = x^{n-1}y^{n-1}(xy+n-1)$  we get by recurrence that

$$x^n y^n = xy(xy+1)(xy+2)\dots(xy+n-1)$$

In calculations with the Weyl algebra it is often useful to decompose  $A_1(\mathbb{C})$  in weight spaces. For  $t \in \mathbb{Z}$  let us define

$$A_1(\mathbb{C})(t) = \{ f \in A_1(\mathbb{C}) \mid [xy, f] = tf \}$$

then the foregoing asserts that  $A_1(\mathbb{C}) = \bigoplus_{t \in \mathbb{Z}} A_1(\mathbb{C})(t)$  where  $A_1(\mathbb{C})(t)$  is equal to

$$\begin{cases} y^t \mathbb{C}[xy] = \mathbb{C}[xy]y^t & \text{for } t \geq 0 \\ x^{-t} \mathbb{C}[xy] = \mathbb{C}[xy]x^{-t} & \text{for } t < 0. \end{cases}$$

For a natural number  $n \geq 1$  we define the  $n$ -th canonical right ideal of  $A_1(\mathbb{C})$  to be

$$\mathfrak{p}_n = x^{n+1}A_1(\mathbb{C}) + (xy+n)A_1(\mathbb{C}).$$

LEMMA 8.47. *The weight space decomposition of  $\mathfrak{p}_n$  is given for  $t \in \mathbb{Z}$*

$$\mathfrak{p}_n(t) = x^{n+1}A_1(\mathbb{C})(t+n+1) + (xy+n)A_1(\mathbb{C})(t)$$

which is equal to

$$\begin{cases} (xy+n)\mathbb{C}[xy]y^t & \text{for } t \geq 0, \\ (xy+n)\mathbb{C}[xy]x^{-t} & \text{for } -n \leq t < 0, \\ \mathbb{C}[xy]x^{-t} & \text{for } t < -n. \end{cases}$$

PROOF. Let  $t = -1$ , then  $\mathfrak{p}_n(-1)$  is equal to

$$x^{n+1}\mathbb{C}[xy]y^n + (xy+n)\mathbb{C}[xy]x$$

Using  $x^{n+1}y^{n+1} = xy(xy+1)\dots(xy+n)$  this is equal to

$$xy(xy+1)\dots(xy+n)\mathbb{C}[xy]y^{-1} + (xy+n)\mathbb{C}[xy]x$$

The first factor is  $(xy+1)\dots(xy+n)\mathbb{C}[xy]x$  from which the claim follows. For all other  $t$  the calculations are similar.  $\square$

One can show that  $\mathfrak{p}_n \not\cong \mathfrak{p}_m$  whenever  $n \neq m$  so the isomorphism classes  $[\mathfrak{p}_n]$  are distinct points in *Weyl* for all  $n$ . We define

$$\text{Weyl}_n = \text{Aut } A_1(\mathbb{C}) \cdot [\mathfrak{p}_n] = \{ [\sigma(\mathfrak{p}_n)] \mid \forall \sigma \in \text{Aut } A_1(\mathbb{C}) \}$$

the orbit in *Weyl* of the point  $[\mathfrak{p}_n]$  under the action of the automorphism group.

EXAMPLE 8.48. The Weyl right ideals *Weyl*<sub>1</sub>.

For a point  $(a, b) \in \mathbb{C}^2$  we define a right ideal of  $A_1(\mathbb{C})$  by

$$\mathfrak{p}_{a,b} = (x+a)^2 A_1(\mathbb{C}) + ((x+a)(y+b)+1)A_1(\mathbb{C}).$$

Observe that  $\mathfrak{p}_1 = \mathfrak{p}_{0,0}$ . Consider the action of the automorphism  $\sigma_2^f$  on these right ideals. As  $f \in \mathbb{C}[x]$  we can write

$$f = f(-a) + (x+a)f_1 \quad \text{with } f_1 \in \mathbb{C}[x].$$

Then, recalling the definition of  $\sigma_2^f$  we have

$$\begin{aligned} \sigma_2^f(\mathfrak{p}_{a,b}) &= (x+a)^2 A_1(\mathbb{C}) + ((x+a)(y+b+f(-a) + (x+a)f_1) + 1)A_1(\mathbb{C}) \\ &= (x+a)^2 A_1(\mathbb{C}) + ((x+a)(y+b+f(-a)) + 1)A_1(\mathbb{C}) = \mathfrak{p}_{a,b+f(-a)} \end{aligned}$$

Now, consider the action of an automorphism  $\sigma_1^f$ . We claim that

$$\mathfrak{p}_{a,b} = A_1(\mathbb{C}) \cap (y+b)^{-1}(x+a)A_1(\mathbb{C})$$



This is easily verified on the special case  $\mathfrak{p}_1$  using the above lemma, the arbitrary case follows by changing variables. We have

$$\begin{aligned} \mathfrak{p}_{a,b} &= A_1(\mathbb{C}) \cap (y+b)^{-1}(x+a)A_1(\mathbb{C}) \\ &\simeq (x+a)^{-1}(y+b)A_1(\mathbb{C}) \cap A_1(\mathbb{C}) \quad (\text{multiply with } h = (x+a)^{-1}(y+b)) \\ &= (y+b)^2A_1(\mathbb{C}) + ((y+b)(x+a) - 1)A_1(\mathbb{C}) \stackrel{\text{def}}{=} \mathfrak{q}_{b,a} \end{aligned}$$

Writing  $f = f(-b) + (y+b)f_1$  with  $f_1 \in \mathbb{C}[y]$  we then obtain by mimicking the foregoing

$$\begin{aligned} \sigma_1^f(\mathfrak{p}_{a,b}) &\simeq \sigma_1^f(\mathfrak{q}_{b,a}) \\ &= \mathfrak{q}_{b,a+f(-b)} \\ &\simeq \mathfrak{p}_{a+f(-b),b} \end{aligned}$$

and therefore there is an  $h \in D_1(\mathbb{C})$  such that  $\sigma_1^f(\mathfrak{p}_{a,b}) = h\mathfrak{p}_{a+f(-b),b}$ .

As the group  $\text{Aut } A_1(\mathbb{C})$  is generated by the automorphisms  $\sigma_1^f$  and  $\sigma_2^f$  we see that

$$\text{Weyl}_1 = \text{Aut } A_1(\mathbb{C}).[\mathfrak{p}_1] \hookrightarrow \{ [\mathfrak{p}_{a,b} \mid a, b \in \mathbb{C}] \}$$

Moreover, this inclusion is clearly surjective by the above arguments. Finally, we claim that  $\text{Weyl}_1 \simeq \mathbb{C}^2$ . That is we have to prove that if

$$\mathfrak{p}_{a,b} = h.\mathfrak{p}_{a',b'} \quad \Rightarrow \quad (a,b) = (a',b').$$

Observe that  $A_1(\mathbb{C}) \hookrightarrow \mathbb{C}(x)[y, \delta]$  where this algebra is the differential polynomial algebra over the field  $\mathbb{C}(x)$  and is hence a right principal ideal domain. That is, we may assume that the element  $h \in D_1(\mathbb{C})$  actually lies in  $\mathbb{C}(x)[y, \delta]$ . Now, induce the filtration by  $y$ -degree on  $\mathbb{C}(x)[y, \delta]$  to the subalgebra  $A_1(\mathbb{C})$ . This is usually called the *Bernstein filtration*. Because  $A_1(\mathbb{C})$  and  $\mathbb{C}(x)[y, \delta]$  are domains we have for all  $f \in A_1(\mathbb{C})$  that

$$\deg(h.f) = \deg(h) + \deg(f).$$

Now, as both  $\mathfrak{p}_{a,b}$  and  $\mathfrak{p}_{a',b'}$  contain elements of degree zero  $x^2 + a$  resp.  $x^2 + a'$  we must have that  $h \in \mathbb{C}(x)$ .

View  $y$  as the differential operator  $-\frac{\partial}{\partial x}$  on  $\mathbb{C}[x]$  and define for every right ideal  $\mathfrak{p}$  of  $A_1(\mathbb{C})$  its *evaluation* to be the *subspace* of polynomials

$$\text{ev}(\mathfrak{p}) = \{ D.f \mid D \in \mathfrak{p}, f \in \mathbb{C}[x] \}$$

where  $D.f$  is the evaluation of the differential operator on  $f$ . One calculates that

$$\text{ev}(\mathfrak{p}_{a,b}) = \mathbb{C}(1 + b(x+a)) + (x+a)^2\mathbb{C}[x]$$

and as from  $\mathfrak{p}_{a,b} = h.\mathfrak{p}_{a',b'}$  and  $h \in \mathbb{C}(x)$  follows that

$$\text{ev}(\mathfrak{p}_{a,b}) = h\text{ev}(\mathfrak{p}_{a',b'})$$

we deduce that  $h \in \mathbb{C}^*$  and hence that  $\mathfrak{p}_{a,b} = \mathfrak{p}_{a',b'}$  and  $(a,b) = (a',b')$ .


Yu. Berest and G. Wilson proved in [6] that the Cannings-Holland correspondence respects the automorphism orbit decomposition.

**THEOREM 8.49.** *We have  $\text{Weyl} = \bigsqcup_n \text{Weyl}_n$  and there are set-theoretic bijections*

$$\text{Weyl}_n \longleftrightarrow \text{Gr}^{\text{ad}}(n)$$

*whence also with  $\text{Calo}_n$ .*

EXAMPLE 8.50. Consider the special case  $n = 1$ . As  $\square$  is the only partition of 1, for every  $\lambda \in \mathbb{C}$ ,  $Gr_\lambda$  is a one-dimensional cell  $\mathbb{A}^1$ , whence  $Gr^{ad}(1) \simeq \mathbb{A}^2$ . In fact we have



where the origin corresponds to the canonical right ideal  $\mathfrak{p}_1$  and the right ideal corresponding to  $(\lambda, \mu)$  is  $\mathfrak{p}_{\lambda, \mu} = (x - \lambda)^2 A_1(\mathbb{C}) + ((x - \lambda)(y - \mu) + 1) A_1(\mathbb{C})$ .

Finally, let us verify that  $\mathfrak{p}_n$  should correspond to a point in  $Gr^{ad}(n)$ . As  $\mathfrak{p}_n = x^{n+1} A_1(\mathbb{C}) + (xy + n) A_1(\mathbb{C})$  we have that

$$\mathfrak{p}_n \cdot \mathbb{C}[x] = \mathbb{C} + \mathbb{C}x + \dots + \mathbb{C}x^{n-1} + (x^{n+1})\mathbb{C}[x]$$

whence  $(x^{n+1})\mathbb{C}[x] \subset \mathfrak{p}_n \cdot \mathbb{C}[x] \subset \mathbb{C}[x]$  and converting this to  $Gr^{ad}$  the corresponding subspace is

$$(x^n)\mathbb{C}[x] \subset x^{-1}\mathfrak{p}_n \cdot \mathbb{C}[x] \subset x^{-1}\mathbb{C}[x]$$

The associated sequence of degrees is  $(-1, 0, 1, \dots, n - 2, n, \dots)$  giving rise to the partition  $p = (\underbrace{1, 1, \dots, 1}_n)$  proving the claim.

If we trace the action of  $Aut A_1(\mathbb{C})$  on  $Weyl_n$  through all the identifications, we get a transitive action of  $Aut A_1(\mathbb{C})$  on  $Calo_n$ . However, this action is non-differentiable hence highly non-algebraic. Berest and Wilson asked whether it is possible to identify  $Calo_n$  with a coadjoint orbit in some infinite dimensional Lie algebra. We have seen before that this is indeed the case if we consider the necklace Lie algebra.

It is our hope that similar results are true for more general quivers and certain families of coadjoint orbits coming from quotient varieties of deformed preprojective algebras.

**References.**

The results of section 8.1 are due to A. King [38]. Section 8.2 is based on the lecture notes of H-P. Kraft [44] and those of A. Tannenbaum [77]. Theorem 8.10 is due to W. Crawley-Boevey and M. Holland [16], the other results of section 8.3 are due to W. Crawley-Boevey [14]. The description of Hilbert schemes via Hilbert stairs is my interpretation of a result of M. Van den Bergh [78]. Example 8.24 is taken from the lecture notes of H. Nakajima [62]. The hyper-Kähler correspondence is classical, see for example the notes of Nakajima, [62], here we follow W. Crawley-Boevey [15]. The applications to the hyper-Kähler correspondence are due to R. Bockland and L. Le Bruyn [8]. The treatment of Calogero particles is taken from G. Wilson’s paper [82]. The fact that the Calogero phase space is a coadjoint orbit is due to V. Ginzburg [24]. The extension to deformed preprojective algebras is independently due to V. Ginzburg [25] and R. Bockland and L. Le Bruyn [8]. The results on adelic Grassmannians are due to G. Wilson [82] and Y. Berest and G. Wilson [6]. More details on the connection with right ideals of the Weyl algebra can be found in the papers of R. C. Cannings and M. Holland [11] and L. Le Bruyn [49].

## Bibliography

1. V.I. Arnold, *Mathematical methods of classical mechanics*, Graduate Texts in Mathematics, vol. 60, Springer-Verlag New York Heidelberg Berlin, 1978.
2. Michael Artin, *On Azumaya algebras and finite-dimensional representations of rings*, J. Algebra **11** (1969), 523–563.
3. ———, *Local structure of maximal orders over surfaces*, Brauer groups in ring theory and algebraic geometry (Fred Van Oystaeyen and Alain Verschoren, eds.), Lecture Notes Mathematics, vol. 917, Springer-Verlag, Berlin Heidelberg New York, 1982, pp. 146–181.
4. ———, *Maximal orders over surfaces*, Invent. Math. (1987).
5. Michael Artin and David Mumford, *Some elementary examples of unirational varieties which are not rational*, Proc. London Math. Soc. **25** (1972), 75–95.
6. Yuri Berest and George Wilson, *Automorphisms and ideals of the Weyl algebra*, preprint, London, 1999.
7. D. Birkes, *Orbits of linear algebraic groups*, Ann. of Math. **93** (1971), 459–475.
8. Raphael Bockland and Lieven LeBruyn, *Necklace Lie algebras and noncommutative symplectic geometry*, preprint-UIA, 2000.
9. N. Bourbaki, *Groupes et algèbres de Lie*, Hermann, Paris.
10. Ken Brown and Ken Goodearl, *Homological aspects of noetherian pi hopf algebras and irreducible modules of maximal dimension*, J. Algebra **198** (1997), 240–265.
11. Robert Cannings and Martin Holland, *Right ideals of rings of differential operators*, J. Algebra **167** (1994), 116–141.
12. J.L. Colliot-Thelene, R.T. Hoobler, and B. Kahn, *The Bloch-Ogus-Gabber theorem*, 1996, preprint.
13. William Crawley-Boevey, *Geometry of representations of algebras*, Lecture Notes, 1993.
14. ———, *Geometry of the moment map for representations of quivers*, preprint, Leeds, 1999.
15. ———, *On the exceptional fibers of Kleinian singularities*, preprint, Leeds, 1999.
16. William Crawley-Boevey and Martin Holland, *Noncommutative deformations of Kleinian singularities*, Duke Math. J. (1998), 605–635.
17. Neil Criss and Victor Ginzburg, *Representation theory and complex geometry*, Birkhäuser, Boston Basel Berlin, 1997.
18. J. Cuntz and D. Quillen, *Algebra extensions and nonsingularity*, J. Amer. Math. Soc. **8** (1995), 251–289.
19. Corrado DeConcini and Claudio Procesi, *Quantum groups, D-modules, representation theory and quantum groups* (L. Boutet de Monvel et al., ed.), Lecture Notes Mathematics, vol. 1565, Springer-Verlag, Berlin Heidelberg New York, 1993, pp. 31–140.
20. Harm Derksen and Jerzy Weyman, *On the canonical decomposition of quiver representations*, preprint, 2000.
21. M. Domokos and A. Zubkov, *Semi-invariants of quivers as determinants*, Transform. Groups (2000).
22. Edward Formanek, *The polynomial identities and invariants of  $n \times n$  matrices*, Regional conference series in mathematics, vol. 78, American Mathematical Society, 1991.
23. Peter Gabriel and Andrei V. Roiter, *Representations of finite-dimensional algebras*, Springer-Verlag, Berlin Heidelberg New York, 1997.
24. Victor Ginzburg, *Non-commutative symplectic geometry and Calogero-Moser space*, preprint Chicago, preliminary version, 1999.
25. ———, *Non-commutative symplectic geometry, quiver varieties and operads*, preprint Chicago, 2000.

26. G. B. Gurevich, *Foundations of the theory of algebraic invariants*, Noordhoff, Groningen, 1964.
27. Dieter Happel and Claus M. Ringel, *Tilted algebras*, Trans. Amer. Math. Soc. **274** (1982), 399–443.
28. Robin Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics, vol. 52, Springer-Verlag, Berlin Heidelberg New York, 1977.
29. Michiel Hazewinkel, *On representations of the symmetric groups, nilpotent matrices, systems, vector bundles and schubert cells*, Representations of Algebras, Third International Conference on Representations of Algebras Held in Puebla, Mexico, August 1980 (M. Auslander and E. Lluís, eds.), Lecture Notes in Mathematics, vol. 903, Springer-Verlag, Berlin Heidelberg New York, 1981.
30. Wim Hesselink, *Desingularization of varieties of nullforms*, Invent. Math. **55** (1979), 141–163.
31. Klaus Hulek, 1979.
32. Victor G. Kac, *Infinite root systems, representations of graphs and invariant theory*, Invent. Math. **56** (1980), 57–92.
33. ———, *Infinite root systems, representations of graphs and invariant theory, ii*, J. Algebra **78** (1982), 141–162.
34. Mikhail Kapranov, *Noncommutative geometry based on commutator expansion*, (1998).
35. Bernhard Keller, *Introduction to A-infinity algebras and modules*, Lecture Notes, 2000.
36. G. Kempf, F. Knudsen, D. Mumford, and B. Saint-Donat, *Toroidal embeddings*, Lecture Notes in Mathematics, vol. 339, Springer-Verlag, Berlin Heidelberg New York, 1973.
37. G. Kempf and L. Ness, *The length of a vector in a representation space*, Algebraic Geometry, Lecture Notes in Mathematics, vol. 732, Springer-Verlag, Berlin Heidelberg New York, 1979, pp. 233–244.
38. Alastair King, *Moduli of representations of finite dimensional algebras*, Quart. J. Math. Oxford Ser. (2) **45** (1994), 515–530.
39. Frances C. Kirwan, *Cohomology of quotients in symplectic and algebraic geometry*, Princeton University Press, 1984.
40. Friedrich Knop, *Beweis des fundamentallemmas und des scheidensatzes*, Algebraische Transformationsgruppen und Invariantentheorie, Algebraic Transformation Groups and Invariant Theory (Hanspeter Kraft, Peter Slodowy, and Tonny A. Springer, eds.), DMV Seminar, vol. 13, Birkhäuser, Basel Boston Berlin, 1989, pp. 110–113.
41. Maxim Kontsevich, *Formal non-commutative symplectic geometry*, The Gelfand mathematical seminars, 1990-1992, Birkhauser verlag, 1993, pp. 173–187.
42. ———, *Triangulated categories and geometry*, Course at the Ecole Normale Sup., Paris, 1998.
43. ———, *Operads and motives in deformation quantization*, mathQA/9904055, 1999.
44. Hanspeter Kraft, *Geometric methods in representation theory*, Representations of Algebras, Workshop Notes of the Third International Conference on Representations of Algebras Held in Puebla, Mexico, August 4-8, 1980 (M. Auslander and E. Lluís, eds.), Lecture Notes in Mathematics, vol. 944, Springer-Verlag, Berlin Heidelberg New York, 1982.
45. ———, *Geometrische methoden in der invariantentheorie*, Aspekte der Mathematik, vol. D1, Vieweg-Verlag, Braunschweig, 1984.
46. ———, *Klassische invariantentheorie : Eine einföhrung*, Algebraische Transformationsgruppen und Invariantentheorie, Algebraic Transformation Groups and Invariant Theory (Hanspeter Kraft, Peter Slodowy, and Tonny A. Springer, eds.), DMV Seminar, vol. 13, Birkhäuser, Basel Boston Berlin, 1989.
47. Hanspeter Kraft and Christine Riedtmann, *Geometry of representations of quivers*, Representations of Algebras, Lecture Notes, vol. 116, London Mathematical Society, 1986, pp. 109–145.
48. Lieven LeBruyn, *Trace rings of generic  $2 \times 2$  matrices*, Mem. Amer. Math. Soc., vol. 363, American Mathematical Society, 1987.
49. ———, *Moduli spaces for right ideals of the Weyl algebra*, J. Algebra **172** (1995), 32–48.
50. ———, *Etale cohomology in non-commutative geometry*, TMR-Lecture Notes, 1997.
51. ———, *Nilpotent representations*, J. Algebra **197** (1997), 153–177.
52. ———, *Orbits of matrix tuples*, Soc. Math. France, Sémin. et congrès **2** (1997), 245–261.
53. ———, *Noncommutative compact manifolds constructed from quivers*, Algebra Montpellier Ann. (e-journal) **1** (1999), 1–5.
54. ———, *Noncommutative geometry @n*, math.AG/9904171, 1999.

55. ———, *Local structure of Schelter-Procesi smooth orders*, Trans. Amer. Math. Soc. (2000), accepted.
56. ———, *Optimal filtrations on representations of finite dimensional algebras*, Trans. Amer. Math. Soc. (2000), accepted.
57. ———, *The singularities of quantum groups*, Proc. London Math. Soc. **80** (2000), 304–336.
58. Lieven LeBruyn and Claudio Procesi, *Semisimple representations of quivers*, Trans. Amer. Math. Soc. **317** (1990), 585–598.
59. Lieven LeBruyn and Aidan Schofield, *Rational invariants of quivers and the ring of matrix-invariants*, Perspectives in Ring Theory (Fred VanOystaeyen and Lieven LeBruyn, eds.), NATO ASI Series, vol. ASI C-233, Kluwer Academic Publishers, Dordrecht Boston London, 1988, pp. 21–30.
60. Domingo Luna, *Slices etales*, Bull. Soc. Math. France Suppl. Mém. **33** (1973), 81–105.
61. J. S. Milne, *Etale cohomology*, Princeton Mathematical Series, vol. 33, Princeton University Press, Princeton New Jersey, 1980.
62. H. Nakajima, *Lectures on Hilbert schemes of points on surfaces*, American Mathematical Society, 1999.
63. Richard S. Pierce, *Associative algebras*, Graduate Texts in Mathematics, vol. 88, Springer-Verlag, Berlin Heidelberg New York, 1982.
64. Claudio Procesi, *The invariant theory of  $n \times n$  matrices*, Adv. in Math. **19** (1976), 306–381.
65. ———, *A formal inverse to the Cayley-Hamilton theorem*, J. Algebra **107** (1987), 63–74.
66. Y. P. Razmyslov, *Trace identities of full matrix algebras over a field of characteristic zero*, Math. USSR-Izv. **8** (1974), 727–760.
67. Irving Reiner, *Maximal orders*, Academic Press, 1975.
68. William Schelter, *Smooth algebras*, J. Algebra **103** (1986), 677–685.
69. Aidan H. Schofield, *Representations of rings over skew fields*, London Mathematical Society Lecture Notes Series, vol. 92, Cambridge University Press, 1985.
70. ———, *General representations of quivers*, Proc. London Math. Soc. **65** (1992), 46–64.
71. ———, *Birational classification of moduli spaces of representations of quivers*, mathAG/9911014, 1999.
72. Aidan H. Schofield and Michel VandenBergh, *Semi-invariants of quivers for arbitrary dimension vectors*, math.RA/9907174, 1999.
73. G. W. Schwartz, *Lifting smooth homotopies of orbit spaces*, Inst. Hautes Études Sci. Publ. Math. **51** (1980), 37–135.
74. S. Shatz, *Profinite groups, arithmetic and geometry*, Annals of Mathematical Studies, vol. 67, Princeton University Press, Princeton New Jersey, 1972.
75. K. S. Siberskii, *Algebraic invariants for a set of matrices*, Siberian Math. J. **9** (1968), 115–124.
76. Peter Slodowy, *Der scheibensatz fur algebraische transformationsgruppen*, Algebraische Transformationsgruppen und Invariantentheorie, Algebraic Transformation Groups and Invariant Theory (Hanspeter Kraft, Peter Slodowy, and Tonny A. Springer, eds.), DMV Seminar, vol. 13, Birkhäuser, Basel Boston Berlin, 1989, pp. 89–114.
77. Allen Tannenbaum, *Invariance and system theory : Algebraic and geometric aspects*, Lecture Notes Mathematics, vol. 845, Springer-Verlag Berlin Heidelberg New York, 1981.
78. Michel VandenBergh, *The Brauer-Severi scheme of the trace ring of generic matrices*, Perspectives in Ring Theory (Fred VanOystaeyen and Lieven LeBruyn, eds.), NATO ASI Series, vol. C 233, Kluwer Academic Publishers, Dordrecht Boston London, 1988, pp. 333–338.
79. Fred VanOystaeyen, *Algebraic geometry for associative algebras*, Marcel Dekker, Inc., 2000.
80. Bruce W. Westbury, *On the character varieties of the modular group*, preprint, Nottingham, 1995.
81. Herman Weyl, *The classical groups*, Princeton University Press, 1946.
82. George Wilson, *Collisions of Calogero-Moser particles and an adelic Grassmannian*, Invent. Math. **133** (1998), 1–41.



# Index

- $GL_n$ 
  - variety, 48
- $\mathfrak{alg}^t$ , 40
- $\mathfrak{alg} @_n$ , 40
- proj, 241
  
- Abelian category, 94
- abelianization, 280
- action
  - contragradient, 17
  - free, 175
  - locally finite, 64
- acyclic
  - object, 95
  - resolution, 95
- additive category
  - of quiver, 285
- adelic Grassmannian, 374
- adjoint matrix, 82
- admissible vertex, 348
- affine scheme, 91
- algebra
  - quantum enveloping, 222
- algebra
  - $A_\infty$ , 322
  - $n$ -th root, 283
  - algebraic functions, 94
  - associated graded, 141, 280
  - Azumaya, 97, 135
  - center, 40
  - connected graded, 23
  - deformed preprojective, 215
  - differential graded, 309
  - differentially graded, 164
  - dual numbers, 140
  - extension, 164
  - formal necklace, 29
  - formal trace, 29
  - free product, 300, 313
  - free trace-, 40
  - group, 303
  - Henselian, 93
  - Hopf, 222
  - infinite necklace, 29
  - infinite trace, 29
  - infinitesimal extension, 150
  - local, 141, 165
  - marked quiver-necklace, 160
  - necklace, 22
  - path, 101
  - Poisson, 280, 331
  - preprojective, 215
  - pull-back, 151
  - quaternion, 362
  - Quillen smooth, 102
  - quiver trace, 109
  - Rees, 281
  - semisimple, 18
  - strict Henselian, 94
  - super, 313
  - variety, 225
  - Weyl, 280, 375
  - Weyl-, 48
  - with trace, 40
  - witness, 71
- algebraic group, 224
- algebraic quotient, 66
- analytic topology, 91
- Artin, M., 88, 135, 229
- Artin-Mumford exact sequence, 131
- associated graded algebra, 141, 280
- associated quiver setting, 248
- associativity constraint, 280
- Atiyah class, 281
- Azumaya algebra, 97, 135
- Azumaya locus, 161
  
- balanced coweight, 232
- barcode, 345
- based linear map, 311
- Berest, Y., 377, 378
- Bernstein filtration, 377
- Berstein-Gelfand-Ponomarev
  - reflection functor, 172
- bimodule, 325
- bipartite double, 181
- Birkes, D., 88
- Bockland, R., 229, 336, 378

- border, 247, 258
- border of corner, 247
- Borel subgroup, 250
- boundary, 106
- bracket invariant, 287
- Brauer group, 100
  
- Calogero system, 366
- Cannings, R., 375, 378
- canonical
  - $A_\infty$ -structure, 322
- canonical decomposition, 168, 192
- canonical right ideal, 376
- capacity, 134
- Cartan
  - decomposition, 63
  - matrix, 166
- Cartan homotopy formula, 331
- Cartan matrix, 222
- category
  - Abelian, 94
- Cayley-Hamilton
  - algebra, 40
  - formal polynomial, 33, 40
  - theorem, 33
- Cayley-smooth
  - $\alpha$ -, 216
  - locus, 193
  - model, 210, 211
- center of algebra, 40
- character, 241, 296, 338
- characteristic polynomial, 3, 33
- Chern numbers, 301
- Chevalley
  - theorem, 168
  - theorem of, 51
- Chinese remainder theorem, 98
- closed form, 312
- cobracket invariant, 287
- cohomology
  - dimension, 123
  - etale, 95
  - Galois, 99
  - pointed set, 96
- commutator element, 215, 347
- commutator filtration, 280
- compact subgroup, 337
- compartment, 180
- completion
  - lifting algebra, 316
- complex moment map, 215
- composition factors, 60
  - multiplicities, 60
- composition series, 60
- cone
  - tangent, 141
- coniveau spectral sequence, 128
- connecting morphism, 299
- connecting sum map, 301
- connection, 319
- constant sheaf, 93
- constructible set, 52
- continuous module, 99
- contraction operator, 321
- coordinate map, 23
- coregular quiver spaces, 226
- corner, 234, 258
  - border, 247
  - optimal, 247
- corner type, 259
- cotangent bundle, 215
- covector, 17
- covers, 91
- Crawley-Boevey, W., 183, 229, 378
- Cuntz, J., 336
- curvature, 311
- cycles, 106
- cyclic vector, 347
  
- deformed preprojective algebra, 215
- deRham
  - cohomology groups, 309
  - complex, 309
- deRham cohomology
  - big, 321
  - big relative, 326
  - noncommutative, 321, 324
  - noncommutative relative, 327
- derivation
  - point, 142
  - trace preserving, 141
  - with respect to morphism, 140
- derived functor, 94
  - right, 94
- Derksen, H., 183, 336
- differential, 143
- differential form
  - noncommutative, 309
  - relative, 325
- differential graded, 309
- differential operator
  - exterior, 309
- differentially graded algebra, 164
- dimension
  - local, 142
- dimension formula, 144
- dimension vector, 69, 73, 105
  - minimal, 371
  - sincere, 166
  - supprt, 114
- direct image, 121
- direct order, 264
- discriminant, 9
- divisor
  - ramification, 134
- dominance order, 73, 74



- dominant weight, 231
- Domokos, M., 336
- double centralizer theorem, 19
- double quiver, 347
- dual numbers, 140
- dual partition, 26
- Dynkin diagram, 166
- equalizer diagram, 93
- equivalent
  - marked quiver, 202
- equivariant map, 22
- etale
  - cohomology functor, 95
  - locally split, 207
  - morphism, 91
  - neighborhood, 93
  - presheaf, 92
  - sheaf, 93
  - site, 92
  - stalk, 94
  - topology, 91
- Euler derivation, 321
- Euler form, 106
- evaluation map, 17
- extension, 105
- extension algebra, 164
- exterior product, 308
- Fedosov product, 311
- field
  - Tate, 124
  - Tsen, 122
- filtration
  - adic, 141
  - finite, 58
- fixed elements, 174
- flag variety, 250
- formal
  - $A_\infty$ -algebra, 322
  - Cayley-Hamilton polynomial, 33, 40
  - necklace algebra, 29
  - substitution, 36
  - trace algebra, 29
  - trace map, 29
- formal completion, 281, 284, 285
- formal structure, 279–281, 297
- formal structure sheaf, 285
- Formanek, E., 45
- free action, 175
- free product, 300
  - algebra, 303
  - group, 303
- free product algebra, 313
- function
  - algebraic, 94
  - elementary symmetric, 6
  - Morse, 86
  - polynomial, 6
  - special, 85
- functor
  - derived, 94
- fundamental group, 302
- fundamental set of roots, 168
- Gabriel, P., 183
- Galois
  - absolute -group, 98
  - cohomology, 99
- general subrepresentation, 236
- generic representation, 168
- generic semi-simple type, 192
- Gerstenhaber, 77
- Ginzburg, V., 336, 378
- global section, 92
- going-down property, 52
- good vertex, 116
- graded localization, 242
- Grassmann variety, 346
- Grassmannian
  - adelic, 374
- Grothendieck topology, 91, 92
- grothendieck topology
  - noncommutative, 283
- group
  - Brauer, 100
  - cohomological dimension, 123
  - multiplicative, 93
  - reductive, 62
  - unitary, 62, 82
- group algebra, 303
- Gurevich, G.B., 44
- Haar measure, 62
- Hamilton equations, 366
- Hamiltonian action, 372
- Hamiltonian flow, 372
- Happel, D., 183
- Hazewinkel, M., 88
- Henselian
  - strict, 94
- Henselian algebra, 93
- Henselization
  - strict, 93
- Hermitian
  - inproduct, 82
  - skew -matrices, 82
  - transpose, 62
- Hermitian inproduct, 337
- Hesselink stratification, 251
- Hesselink stratum, 252
- Hesselink, W., 77, 251, 275
- Hilbert
  - criterion, 55, 57
  - theorem 90, 100
- Hilbert scheme, 356
- Hilbert-Chow map, 361
- Hochschild-Serre spectral sequence, 124

- Holland, M., 229, 375, 378
- homology groups, 162
- Hopf algebra, 222
- Hulek, K., 301
- ideal
  - maximal, 91
  - of relations, 47
  - prime, 91
  - trace relations, 42
- idempotent, 26
- implicit function, 91
- implicit function theorem, 53
- indecomposable
  - representation, 165
  - roots, 166
- indecomposable projective, 295
- index
  - ramification, 133
- inertia, 134
- inertial degree, 134
- infinitesimal extension
  - of algebra, 150
  - trivial, 150
  - versal, 150
- injective
  - object, 94
  - resolution, 95, 164
- integrable system, 366
- integral extension, 50
- invariant
  - polynomial, 12, 64
- isotypical
  - component, 64
  - decomposition, 64
  - Gcomponent, 146
- Jacobi identity, 334
- Jacobian matrix, 91, 140
- Jordan
  - Hölder theorem, 60
  - Weierstrass theorem, 4
  - moves, 3
  - normal form, 3
- Jordan move
  - permutation, 231
- Kähler differential, 308
- Kähler potential, 338
- Kac, V., 172, 183
- Kalman code, 344
  - generic, 346
- Kalman theorem, 346
- Kapranov, M., 279, 336
- Karoubi complex, 323
  - relative, 327
- Keller, B., 165, 183
- Kempf, G., 88
- Killing form, 83
- King, A., 275, 378
- Kirillov-Kostant bracket, 372
- Kirwan, F., 275
- Knop, F., 183
- Kontsevich bracket, 333
- Kontsevich, M., 183, 336
- KP hierarchy, 375
- Kraft, H-P., 45, 59, 88, 253, 275, 378
- Kraft, H.P., 183
- Krull
  - Hauptideal result, 51
- Kummer sequence, 93
- large vertex, 116
- Laurent series, 374
- Le Bruyn, L., 135, 183, 229, 275, 336, 378
- left orthogonal, 180
- Leray
  - spectral sequence, 121
- Levi subgroup, 234, 259
- lexicographic order, 28
- Lie algebra
  - semi-simple, 222
- Lie derivative, 331
- linear dynamical system, 341
  - completely controllable, 342
  - completely observable, 343
  - equivalent, 343
  - system identification, 344
  - time invariant, 341
- liouville theorem, 367
- local algebra, 141, 165
- local dimension, 142
- local quiver, 189
- locally closed set, 52
- Luna, D., 183
- marked
  - quiver-order, 160
- marked quiver
  - equivalent, 202
  - Euler form, 190
  - representation, 159
- matrix
  - addition, 3
  - adjoint, 82, 337
  - base-change, 3
  - characteristic polynomial, 3, 33
  - companion, 347
  - conjugacy class, 3
  - conjugate, 3
  - control, 342
  - cyclic, 7
  - eigenvalues, 4
  - exponential, 341
  - fundamental, 341
  - Hermitian, 366
  - Jacobian, 91
  - multiplication, 3

- normal, 83
- permutation, 3, 42
- rank, 4
- trace, 13
- tuple, 13
- maximal ideal, 91
- maximal order, 133, 134
- measure
  - of singularity, 160
- microlocalization, 282
- microstructure sheaf, 282
- Milne, J.S., 135
- model
  - Cayley-smooth, 210, 211
- module, 48
  - associated graded, 58
  - automorphism, 48
  - composition factors, 60
  - continuous, 99
  - decomposable, 59
  - extension, 105
  - indecomposable, 52, 78
  - morphism, 48
  - simple, 60
- moduli space, 241
- moduli spaces, 285
- moment map
  - complex, 215, 349
  - real, 82, 84, 87, 337, 338
- monomial
  - equivalent, 29
- morphism
  - closed, 50
  - dominant, 49
  - etale, 91
  - fiber, 9
  - finite, 50, 143
  - flat, 174
  - open, 174
  - section, 8
- Morse function, 86
- multilinear map, 16
- multiplicative group, 93
- multiplicities
  - composition factors, 60
- multiplicity, 64
- Mumford, D., 135
- Nagata-Higman problem, 23, 37
- Nakajima, H., 378
- Nakayama lemma, 294
- necklace
  - algebra, 22
  - infinite algebra, 29
  - relation, 30
    - fundamental, 32
  - word, 22
- necklace algebra, 264
- necklace word, 327
- neighborhood
  - etale, 93
- Ness, L., 88
- Newton function, 33
- Noether
  - normalization, 50
- noncommutative
  - 1-form, 327
  - function, 327
- noncommutative deRham cohomology, 308
- noncommutative differential form, 308
- noncommutative differential form, 309
- noncommutative manifold, 301
  - compact, 301
- noncommutative vectorfield, 331
- nonsingular point, 142
- norm map, 71
- normal domain, 50
- normal space, 144
  - to orbit, 145
- normal variety, 194, 195
- nullcone, 57
- number of parameters, 169, 217
- object
  - acyclic, 95
  - injective, 94
- one parameter subgroup, 54
- operad, 161
  - chain, 163
  - endomorphism, 162
  - homology, 163
  - linear, 162
  - morphism, 162
  - tiny interval, 161
- operator inner product, 337
- optimal corner, 247, 259
- optimal corner type, 260
- orbit, 3
  - map, 52
- order, 133, 134
  - dominance, 73, 74
  - lexicographic, 28
  - maximal, 133, 134
  - quiver, 109
- orthogonal
  - left, 180
- parabolic subgroup, 234, 259
- partition, 26
  - dual, 26
  - transposition, 27
- path algebra, 101
- phase space
  - Calogero, 368
- Poincare duality, 130
- Poincare-Birkhoff-Witt theorem, 279
- point derivation, 142

- pointed set, 95
- Poisson algebra, 280, 331, 372
- Poisson bracket, 371
- Poisson manifold, 372
- Poisson structure, 223
- polarization, 17, 21
- polynomial
  - linear term, 139
  - minimal degree, 139
  - order, 139
- polynomial invariants, 12
- preprojective algebra, 215
  - deformed, 215, 347
- presheaf
  - etale, 92
- primary
  - $\lambda$ -, 374
- prime ideal, 91
- prime ideal spectrum, 91
- principal character, 281
- principal element theorem, 143
- Procesi, C., 44, 88, 135, 183, 229
- proper function, 338, 339
- proper morphism, 246
- pull-back
  - algebra, 151
- quadratic form
  - positive definite, 166
  - radical, 166
  - semidefinite, 166
- quantum Borel algebra, 225
- quantum enveloping algebra, 222
- quantum function algebra, 225
- quantum groups, 222
- quaternion algebra, 362
- Quillen
  - smooth algebra, 102
- Quillen, D., 336
- quiver, 101
  - order
    - marked, 160
  - arrow, 101
  - dual, 215
  - local, 189
  - marked, 158
  - necklace algebra, 111
    - marked, 160
  - opposite, 215
  - order, 109
    - Schurian, 175
  - path algebra, 101
  - reflected, 170
  - representation, 105
  - root, 172
  - shrinking, 200
  - simplified, 201
  - strongly connected, 114
  - strongly connected component, 114
  - tame, 168
  - trace algebra, 109
  - vertex, 101
  - Weyl group, 172
- quiver representation
  - normal space, 189
- quotient
  - algebraic, 66
  - scheme, 70
- quotient variety
  - stratification, 191
- ramification, 134
  - divisor, 134
  - index, 133
- ramification locus, 207
- rank
  - general homomorphism, 236
- Rasmyslov, Y.P., 44
- real moment map, 82, 84, 87
- real points, 82
- reductive group, 62
- Rees ring, 281
- reflected dimension vector, 170
- reflected quiver, 170
- reflection, 172
- reflexion functor, 170
- representable functor, 283
- representation, 47
  - extension, 105
  - generic, 168
  - indecomposable, 165
  - marked quiver, 159
  - Schur, 174
  - type, 64
- representation space
  - restricted, 110
- representation type, 155, 187
  - finite, 168
  - generic semi-simple, 192
  - order, 191
  - successor, 191
- representation variety
  - reduced, 47, 49
  - reduced trace preserving, 68
  - scheme structure, 48, 49
- restitution, 17, 21
- Reynolds operator, 42
- Riedtmann, Ch., 183
- Ringel, C.M., 183
- Roiter, A.V., 183
- root, 170, 172
  - $\lambda$ -Schur, 218
  - imaginary, 172
  - indecomposable, 166
  - real, 172
  - Schur, 174

- system, 166
- root system, 168
- roots
  - fundamental set, 168
  - simple, 114
- scheme
  - affine, 91
- Schofield, A., 183, 275, 294, 336
- Schur
  - lemma, 165
  - representation, 174
  - root, 174
- Schur lemma, 303
- Schur root
  - $\lambda$ -, 218
- section
  - global, 92
- self-extension
  - trace preserving, 145
- semi invariants, 285
- semi-invariant, 241
  - determinantal, 286
  - polynomial, 286
  - standard determinantal, 294
- semi-simple Lie algebra, 222
- semistable representation, 242, 245
- separability locus, 225
- separated filtration, 281
- sequence
  - spectral, 119
- set of roots, 98
- Shatz, S., 135
- sheaf
  - constant, 93
  - etale, 93
  - roots of unity, 93
- Siberskii, K.S., 44
- simple root
  - $\lambda$ -, 349
- simply connected, 129
- singular cube, 163
- singularity
  - measure of, 160
- site
  - etale, 92
- Slodowy, P., 183
- smooth
  - Cof, 149
  - $k$ -, 280
  - $n$ - locus, 159
  - Cayley-, 149
  - Grothendieck-, 149
  - Quillen-, 149
- smooth locus, 160
- smooth point, 142
- spectral sequence, 119
  - coniveau, 128
  - Hochschild-Serre, 124
  - Leray, 121
  - limiting term, 121
- spectrum
  - prime ideal, 91
- splitting
  - of vertex, 201
- stabilizer subgroup, 10, 144
- stable
  - under group action, 48
- stable representation, 245
- stalk, 94
- stratification, 191
- stratum, 10
- string of numbers, 232
- structure constant
  - variety, 225
- subgroup
  - open, 99
- substitution invariant, 36
- sum family, 301
- sum manifold, 301
- super algebras, 313
- support
  - dimension vector, 114
- symmetric
  - group, 12
  - Newton function, 33
  - polynomial, 12
- symplectic
  - action, 371
  - derivation, 332
- symplectic structure, 331
  - canonical, 371
  - noncommutative, 331
- symplectic vectorfield, 331
- tangent cone, 141
- tangent space, 139
- Tannenbaum, A., 378
- Tate
  - field, 124
  - theorem, 124
- tensor
  - pure, 17
- tensor algebra, 311
- test object, 150
- thickening, 281
- Tits form, 166
- topology
  - analytic, 53, 91
  - etale, 91
  - Grothendieck, 91, 92
  - Zariski, 53, 91
- torus knot, 302
  - group, 302
- trace
  - formal, 29

- generated, 40
- infinite algebra, 29
- map, 39
- morphism, 40
- nondegenerate, 23
- preserving representation, 68
- relation, 30
  - fundamental, 32
- trace pairing, 215
- Tsen
  - theorem, 123
- Tsen field, 122
- twisted cubic, 10
- twisted form, 96
  
- unipotent element, 169
- unipotent group, 169
- unitary group, 62
  - multiple, 337
- universal localization, 294
  - universal property, 294
- universal property
  - trace embedding, 44
  
- Van den Bergh, M., 336, 378
- variety
  - irreducible, 49
- vector
  - column, 3
  - dimension, 69
  - dimension, 73, 105
  - row, 17
- vertex
  - admissible, 172
  - dominant form, 258
  - focus, 114
  - good, 116
  - interval, 257
  - large, 116
  - prism, 114
  - sink, 114
  - source, 114
  - splitting, 201
  - starting, 101
  - symmetric group, 258
  - terminating, 101
  
- weight, 215, 232
- weight space, 84
- weight space decomposition, 58, 232
- Westbury, B., 336
- Weyl algebra, 48, 280, 375
- Weyl group
  - of quiver, 172
- Weyl skewfield, 375
- Weyman, J., 183, 336
- wild quiver, 341
- Wilson, G., 375, 377, 378
- witness algebra, 71
  
- word, 321
- wrinkle, 131
  
- Yang-Mills derivation, 317
- Young
  - diagram, 5, 26
  - move, 74
  - symmetrizer, 27
  - tableau, 26, 74
  
- Zariski closed, 241
- Zariski topology, 48, 91
  - closed set, 48
- Zubkov, A., 336